

## Original Article

# Red cell folate and predicted neural tube defect rate in three Asian cities

Timothy J Green PhD<sup>1</sup>, C Murray Skeaff PhD<sup>1</sup>, Bernard J Venn PhD<sup>1</sup>, Jennifer EP Rockell MSc<sup>1</sup>, Joanne M Todd MSc<sup>2</sup>, Geok L Khor PhD<sup>3</sup>, Su Peng Loh PhD<sup>3</sup>, G Duraisamy MD<sup>3</sup>, Siti Muslimatun PhD<sup>4</sup>, Rina Agustina PhD<sup>4</sup>, Xu Ling MD<sup>5</sup> and Xiaoping Xing PhD<sup>5</sup>

<sup>1</sup>Department of Human Nutrition, University of Otago, Dunedin, New Zealand

<sup>2</sup>Fonterra Brands Ltd, Auckland, New Zealand

<sup>3</sup>Department of Nutrition & Health Sciences, Faculty of Medicine & Health Sciences, Universiti Putra Malaysia, Serdang, Malaysia

<sup>4</sup>SEAMEO TROPMED Regional Center for Community Nutrition, University of Indonesia, Jakarta

<sup>5</sup>Peking Union Medical College Hospital, Beijing, China

**Background:** Periconceptional folic acid reduces neural tube defect (NTD) risk. Red blood cell folate concentration is inversely associated with NTD risk. In many countries there is a lack of information on NTD rates. Red cell folate status in women of childbearing age may be a surrogate for NTD rates and may be helpful in identifying countries or regions most likely to benefit from improved folate status.

**Objective:** To predict NTD rates using red cell folate concentrations in women of childbearing age living in three Asian cities

**Design:** Cross-sectional convenience samples of non-pregnant women living in Beijing (n=220), Kuala Lumpur (n=389), and Jakarta (n=129).

**Results:** Red cell folate concentrations were highest ( $p<0.001$ ) in women from Jakarta at 872 nmol/L (95% CI: 833, 910) followed by Kuala Lumpur at 674 nmol/L (95% CI: 644, 704) and lowest in Beijing at 563 nmol/L (95% CI: 524, 601). Accordingly, predicted NTD rates were highest in Beijing at 30/10000 (95% CI: 27, 33), followed by Kuala Lumpur at 24/10000 (95% CI: 22, 25), and lowest in Jakarta at 15/10000 (95% CI: 14, 15).

**Conclusion:** Our red blood cell folate data suggests that of the three cities improving the folate status of women in Beijing would have the greatest impact on NTD rates.

**Key Words:** red cell folate, NTD, fortification, Asia, women

## Introduction

High folate status in women of childbearing age reduces the risk of neural tube defects (NTDs). In a large cohort of Irish women the risk of having an NTD-affected pregnancy was lowest in women with red cell folate concentrations above 905 nmol/L.<sup>1</sup> Taking folic acid during the periconceptional period reduces individual risk and population incidence of NTD.<sup>2-4</sup> Any programme that increases folate status is likely to have the greatest impact on NTD rates in populations with low folate status. For example, in a southern region of China, the background NTD rate was approximately 10 per 10000 births and mean red cell folate concentration in women aged 35 - 44 y was 911 nmol/L, whereas, in a northern region where red cell folate status was lower (508nmol/L), the background NTD rate was much higher, 50 - 60 per 10000 births.<sup>5</sup> Furthermore, the NTD rate in these regions following a public health campaign that encouraged women to take a daily supplement containing 400 µg folic acid per day was 6-7/10000 in compliant women.<sup>6</sup> Thus, despite large pre-supplementation differences in NTD rates between the northern and southern regions, NTD rates post-supplementation were similar. This is suggestive of a folate-responsive NTD

threshold, possibly around a rate of 5-6/10000 below which the prevalence of NTD cannot be reduced further with folic acid. This is consistent with NTD rates in North America which have not fallen below 5/10000 since mandatory fortification.<sup>2,4</sup>

In many countries, information on birth defects is lacking. In the absence of this information, it is possible that red cell folate status in women of childbearing age may be a surrogate for NTD rates and helpful in identifying countries or regions most likely to benefit from improved folate status. Wald et al used this approach to compare predicted and actual decline in NTD rates in North America.<sup>7</sup> We have used a similar approach to predict NTD rates using red cell folate concentrations in women of childbearing age living in three Asian cities, Kuala Lumpur, Jakarta, and Beijing.

**Corresponding Author:** Dr. Bernard J Venn, Department of Human Nutrition, University of Otago, P.O. Box 56, Dunedin, New Zealand.

Tel: +64 3 479 5068; Fax: +64 3 479 7958

Email: Bernard.venn@stonebow.otago.ac.nz

Manuscript received 22 June 2006. Accepted 29 August 2006.

## Methods

### Subjects

Non-pregnant women 18–40 y were recruited using convenience sampling from Beijing, Kuala Lumpur, and Jakarta. Recruitment was via mail, the posting of flyers, the Internet, and word of mouth. Women were excluded if they were breastfeeding or had breastfed within 12 months, or had a serious or chronic illness. In Beijing, 220 women 20–35 y were recruited between February and June 2002. A total of 389 women 18–40 y comprising approximately equal proportions of Malay, Chinese and Indian ethnicities were recruited from worksites and universities in Kuala Lumpur and its suburbs between January and March 2005. In Jakarta, 129 women 18–40 y were recruited in late 2004. Demographic details of the women were collected using questionnaires. Approval to conduct the studies was provided by Ethics Committees within each of the study regions: The Peking Union Medical College Hospital Ethics Committee, Beijing, China; the Ethics Committee of the Faculty of Medicine and Health Sciences, Universiti Putra, Malaysia; Ethical Committee Faculty of Medicine, University of Indonesia, Jakarta.

### Blood collection and laboratory analysis

Blood samples were taken by venipuncture into tubes containing EDTA following an overnight fast. Blood was processed and red cell folate concentrations measured as

described by O’Broin and Kelleher using a microtitre technique with chloramphenicol-resistant *Lactobacillus casei* as the test microorganism.<sup>8</sup> Whole blood standard (National Institute for Biological Standards and Control, UK) with an assigned folate concentration of 29.4 nmol/L was used to generate a standard curve. The intra-assay coefficient of variation was 10.4% for whole blood based upon repeated measurements of pooled samples.

### Analysis

Predicted NTD risk was estimated based on each woman’s red cell folate concentration using the predictive equation of Daly et al  $\exp[1.6463 - (1.2193 \times \ln\{\text{RCF}\})]$ .<sup>1</sup> ANOVA was used to test for differences between cities in red cell folate and predicted NTD rates, with a Bonferroni correction for multiple comparisons.

## Results

Demographic characteristics of the participants are shown in Table 1. In Beijing, around half of the women were single, whereas in Kuala Lumpur and Jakarta over two thirds of the women were single. The majority of women had received secondary education or higher. Mean red cell folate concentrations and predicted NTD rates are shown in Table 2. Red cell folate concentrations were higher ( $p < 0.001$ ) in women from Jakarta than from Beijing or Kuala Lumpur by 198 (95%CI; 140 to 255) and 309 (254 to 365) nmol/L, respectively. Folate concentra-


**Table 1.** Demographic and dietary characteristics

Descriptive	Beijing	Kuala Lumpur	Jakarta
Number of women	220	389	129
Ethnicity	Chinese Malay (27%) Chinese (31%) Indian (32%)		
Age (y)	28 ± 8.1	25 ± 5.5	30 ± 6.8
BMI (kg/m <sup>2</sup> )	20.9 ± 3.4	22.6 ± 4.5	22.5 ± 4.3
Smokers n (%)	2 (1%)	5 (1%)	9 (7%)
Oral contraceptive users	94 (43%)	NA	NA
Ever been pregnant	94 (43%)	NA	90 (69.8%)
Alcohol consumers	11 (5%)	20 (5%)	NA
Multivitamin or B-complex supplement use	NA	20 (5%)	20 (16%)
Marital status			
Single	113 (51%)	293 (76%)	96 (74%)
Married	104 (47%)	89 (23%)	30 (23%)
Widowed/divorced	1 (<1%)	5 (1%)	3 (2%)
Highest Education			
Primary	2 (1%)	10 (2.6%)	28 (21.7%)
Secondary	23 (11%)	62 (16%)	20 (15.5%)
Matriculation	71 (32%)	52 (13%)	60 (46.5%)
Polytechnic/University	122 (55%)	263 (68%)	21 (16.3%)

**Table 2** Red cell folate concentrations<sup>1</sup> and predicted NTD rate

	Beijing	Kuala Lumpur	Jakarta
Red cell folate (nmol/L)	563 (524 to 601) <sup>a</sup>	674 (644 to 704) <sup>b</sup>	872 (833 to 910) <sup>c</sup>
Predicted NTD rate (/10000)	30 (27 to 33) <sup>a</sup>	24 (22 to 25) <sup>b</sup>	15 (14 to 15) <sup>c</sup>

<sup>1</sup>n = 220, 389, and 129 women in Beijing, Kuala Lumpur and Jakarta, respectively; values are means (95%CI). Numbers not sharing a common superscript are significantly different,  $p < 0.001$ .



**Figure 1.** Distribution of red blood cell folate concentrations among women in Kuala Lumpur (n=389), Beijing (n=220) and Jakarta (n=129).

tions in Kuala Lumpur were 112 (58 to 166) nmol/L higher than in Beijing. More details on the Malaysian dietary and blood folate study will be published in the Asia Pacific Journal of Clinical Nutrition (in press). Accordingly, predicted NTD rates were highest in Beijing, followed by Kuala Lumpur and lowest in Jakarta. The percentages of women falling into various categories of red cell folate concentrations are shown in Figure 1. The categories are the same as those used by Daly et al.<sup>1</sup> in developing the equation we used to predict NTD risk. Women from Jakarta are distributed into the higher red cell folate categories, whereas women from Beijing and Kuala Lumpur are represented across a wider range of the categories. Eighteen percent of the women in Beijing, 8% in Kuala Lumpur and no women in Jakarta had red cell folate concentrations below 305 nmol/L, a cut-off value indicative of folate deficiency.<sup>9</sup>

## Discussion

Our results have shown that red cell folate concentrations in women living in three major Asian cities are markedly different, with the highest concentrations in Jakarta and lowest in Beijing. Indeed, eighteen percent of women in Beijing had red cell folate concentrations that would place them at increased risk of folate deficiency anaemia, major health consequences of which would be fatigue and suboptimal cognitive performance.<sup>10</sup> Moreover, for women who become pregnant, low maternal folate status increases the risk of low birth weight<sup>11</sup> and birth defects.<sup>6, 12</sup>

The most likely explanation for the higher folate status in women in Jakarta is the implementation of mandatory folic acid fortification in 2001 of domestic and imported wheat, at a dose of 200 µg per 100 g;<sup>13</sup> neither China nor Malaysia have a mandatory fortification scheme. Mean red cell folate concentrations in Jakarta (872 nmol/L) are slightly higher than those reported in the United States post-fortification (787 nmol/L), which suggests a higher intake of total folate (naturally occurring and added) by women in Jakarta.<sup>14, 15</sup> If consumption of wheat flour is similar between the two countries, the slightly higher red cell folate concentrations in Jakarta are consistent with the higher level of fortification in Jakarta than in the United States (140 µg/100 g). In countries without mandatory fortification, such as Malaysia and China, the main determinant of red cell folate is dietary folate intake.<sup>16</sup>

Although we did not assess folate intake, the differences in red cell folate concentrations between Kuala Lumpur and Beijing will reflect differences in the folate consumption of women in the cities.

We cannot generalize the results to all women living in China, Indonesia, and Malaysia because convenience samples of women living in the three urban centres were recruited that may not be representative of the population. Whether these women have higher or lower folate status compared with other urban-dwelling or rural women is not known. Nevertheless, our red cell folate concentrations in women living in Beijing are consistent with those reported by Hao et al, where a representative sample of Beijing women (n=205) living in urban or rural settings had a mean red cell folate of 508 nmol/L; a difference of 55 nmol/L compared with women in our study.<sup>5</sup>

In many countries NTD incidence rates are unknown, may not include pregnancy terminations, or may not be based on complete ascertainment of all pregnancy outcomes. In the absence of NTD incidence data, it is possible to predict NTD rates using results from a case-control study conducted in Ireland which quantified the relation between red cell folate and the risk of NTDs.<sup>1</sup> Though applying the formula to predict NTD rates in other populations has limitations, because there are factors other than folate status that determine NTD rates, this approach was used by COMA<sup>17</sup> when considering mandatory fortification in the United Kingdom. Furthermore, Wald et al.<sup>7</sup> used this model to compare predicted and actual changes in NTD rates from large randomized trials and mandatory fortification in the United States. Applying the equation of Daly et al.<sup>1</sup> to our red cell folate data suggests that the NTD rate in Beijing is twice that of Jakarta. The applicability of NTD risk estimates found in Irish women has not been confirmed in an Asian setting; however, the totality of evidence is unequivocal that implementing folic acid fortification in areas with low folate status reduces NTD rates.<sup>3, 5, 6, 18</sup>

Our results indicate low folate status amongst women living in Beijing and suggest that of the three cities improving the folate status of women in Beijing would have the greatest impact on NTD rates.

## Acknowledgement

This work was supported by a research grant from Fonterra Brands Limited, New Zealand.

**References**

1. Daly LE, Kirke PN, Molloy A, Weir DG, Scott JM. Folate levels and neural tube defects. Implications for prevention. *JAMA*. 1995;274(21):1698-702.
2. Honein MA, Paulozzi LJ, Mathews TJ, Erickson JD, Wong LY. Impact of folic acid fortification of the US food supply on the occurrence of neural tube defects. *JAMA*. 2001;285(23):2981-6.
3. Liu S, West R, Randell E, Longerich L, O'Connor K S, Scott H, et al. A comprehensive evaluation of food fortification with folic acid for the primary prevention of neural tube defects. *BMC Pregnancy Childbirth*. 2004 Sep 27;4(1):20.
4. Ray JG, Meier C, Vermeulen MJ, Boss S, Wyatt PR, Cole DE. Association of neural tube defects and folic acid food fortification in Canada. *Lancet*. 2002 Dec 21-28;360(9350):2047-8.
5. Hao L, Ma J, Stampfer MJ, Ren A, Tian Y, Tang Y, et al. Geographical, seasonal and gender differences in folate status among Chinese adults. *J Nutr*. 2003 Nov;133(11):3630-5.
6. Berry RJ, Li Z, Erickson JD, Li S, Moore CA, Wang H, et al. Prevention of neural-tube defects with folic acid in China. China-U.S. Collaborative Project for Neural Tube Defect Prevention. *N Engl J Med*. 1999;341(20):1485-90.
7. Wald NJ. Quantifying the effect of folic acid. *The Lancet*. 2001;358(9298):2069-73.
8. O'Briain S, Kelleher B. Microbiological assay on microtitre plates of folate in serum and red cells. *J Clin Pathol*. 1992;45(4):344-7.
9. Gibson RS. Principles of Nutritional Assessment. 2nd ed. Auckland: Oxford University Press; 2005.
10. Herbert V. Folic Acid. *Annu Rev Med*. 1965;16:359-70.
11. Relton CL, Pearce MS, Parker L. The influence of erythrocyte folate and serum vitamin B12 status on birth weight. *Br J Nutr*. 2005 May;93(5):593-9.
12. Canfield MA, Collins JS, Botto LD, Williams LJ, Mai CT, Kirby RS, et al. Changes in the birth prevalence of selected birth defects after grain fortification with folic acid in the United States: findings from a multi-state population-based study. *Birth Defects Res A Clin Mol Teratol*. 2005 Oct;73(10):679-89.
13. Hardinsyah, Suroso. Micronutrient programs in Indonesia. *Food and Nutrition Bulletin*. 2004;25:82-3.
14. Erickson JD, Mulinare J, Yang Q, Johnson CL, Pfeiffer C, Gunter EW, et al. Folate status in women of childbearing age, by race/ethnicity --- United States, 1999-2000. *MMWR*. 2002;51:80-810.
15. Wright JD, Bialostosky K, Gunter EW, Carroll MD, Najjar MF, Bowman BA, et al. Blood folate and vitamin B12 : United States, 1988-94. National Center for Health Statistics. *Vital Health Stat*. 1998;11 (243).
16. Green TJ, Allen OB, O'Connor DL. A three-day weighed food record and a semiquantitative food-frequency questionnaire are valid measures for assessing the folate and vitamin B-12 intakes of women aged 16 to 19 years. *J Nutr*. 1998 Oct;128(10):1665-71.
17. Committee on Medical Aspects of Food and Nutrition. Folic acid and the prevention of disease. Report on health and social subjects. No 50. London: The Stationery Office; 2000.
18. Lopez-Camelo JS, Orioli IM, da Graca Dutra M, Nazer-Herrera J, Rivera N, Ojeda ME, et al. Reduction of birth prevalence rates of neural tube defects after folic acid fortification in Chile. *Am J Med Genet A*. 2005 Jun 1;135(2):120-5.

## Original Article

# Red cell folate and predicted neural tube defect rate in three Asian cities

Timothy J Green PhD<sup>1</sup>, C Murray Skeaff PhD<sup>1</sup>, Bernard J Venn PhD<sup>1</sup>, Jennifer EP Rockell MSc<sup>1</sup>, Joanne M Todd MSc<sup>2</sup>, Geok L Khor PhD<sup>3</sup>, Su Peng Loh PhD<sup>3</sup>, G. Duraisamy MD<sup>3</sup>, Siti Muslimatun PhD<sup>4</sup>, Rina Agustina PhD<sup>4</sup>, Xu Ling MD<sup>5</sup> and Xiaoping Xing PhD<sup>5</sup>

<sup>1</sup>Department of Human Nutrition, University of Otago, Dunedin, New Zealand

<sup>2</sup>Fonterra Brands Ltd, Auckland, New Zealand

<sup>3</sup>Department of Nutrition & Health Sciences, Faculty of Medicine & Health Sciences, Universiti Putra Malaysia, Serdang, Malaysia

<sup>4</sup>SEAMEO TROPMED Regional Center for Community Nutrition, University of Indonesia, Jakarta

<sup>5</sup>Peking Union Medical College Hospital, Beijing, China

## 三大亚洲城市的红血球叶酸及神经管畸形的预测发生率

**背景：**妊娠期间的叶酸摄取有效降低神经管畸形的风险。红血球的叶酸水平及神经管畸形的风险是互为相比的。现今的很多国家都缺乏神经管畸形发生率的相关资料。育龄妇女的红血球叶酸水平也许是可代替神经管畸形发生率的另一种指标，并且可用于确认那一些从提高叶酸水平计划当中最可能受惠的国家或区域。

**目的：**使用三大亚洲城市的育龄妇女之红血球的叶酸水平以预测神经管畸形的发生。

**方法：**采用断面式方法以录取居于北京(人数：220)，吉隆坡(人数：389)及雅加达(人数：129)的未孕妇女参与此项调查。

**结果：**雅加达的妇女拥有最高的( $p<0.001$ )红血球叶酸水平，872 nmol/L (95% CI; 833, 910)，其次是吉隆坡妇女，叶酸水平高达674 nmol/L (95% CI: 644, 704)。北京妇女的叶酸水平为之最低，只达563 nmol/L (95% CI: 524, 601)。有鉴于此，预测北京的神经管畸形的发生率将是最高，达至30/10000 (95% CI: 27, 33)，接着是吉隆坡，24/10000 (95% CI: 22, 25)，而雅加达的发生率将是最低，只是15/10000 (95% CI: 14, 15)。

**結論：**根据我们的红血球叶酸研究结果显示，在这三大城市中，北京将是在提高叶酸水平以降低神经管畸形的发生率的计划中最为受惠的城市，并拥有最显著的效果。

**關鍵詞：**红血球叶酸，神经管畸形，食品强化，亚洲，妇女。