

Original Article

Relationship between dietary patterns and brachial-ankle pulse wave velocity among middle-aged adults in Japan

Shota Moyama MS¹, Kazuhiro Minami PhD², Mihiro Yano PhD³, Masumi Okumura MS³, Susumu Hayashi PhD⁴, Hiroshi Takayama PhD⁵, Akira Yorimoto PhD⁶

¹Department of Nutrition, Hikone Municipal Hospital, Shiga, Japan

²Department of Faculty of Bio-Industry, Tokyo University of Agriculture, Hokkaido, Japan

³Department of Food Science and Nutrition, The University of Shiga Prefecture, Shiga, Japan

⁴Department of dermatology, Hikone Municipal Hospital, Shiga, Japan

⁵Department of Internal Medicine, Takashima Municipal Hospital, Shiga, Japan

⁶Department of Food and Nutrition, Kyoto Women's University, Kyoto, Japan

Background and Objectives: Arterial stiffness is a leading cause of cardiovascular disease (CVD), and it is considered to be affected by dietary intake. However, few studies have examined the relationship between major dietary patterns and brachial-ankle pulse wave velocity (baPWV) among Japanese middle-aged subjects. We studied whether major dietary patterns were associated with baPWV in this population. **Methods and Study Design:** Between 2009 and 2012, 70 Japanese middle-aged subjects (39 men and 31 women) with no history of stroke, coronary heart disease, or cancer were studied. Dietary intake was documented using a validated food-frequency questionnaire, and dietary patterns were generated using factor analysis. Correlational analyses were performed between baPWV and identified dietary patterns and between baPWV and individual risk factors (total cholesterol, triglyceride, low-density lipoprotein cholesterol [LDL-C], high-density lipoprotein cholesterol [HDL-C], non-HDL-C, LDL/HDL ratio, and augmentation index). **Results:** Two dietary patterns were identified: a rice-rich pattern (high in rice, breads, oils and fats, meats, and confectionery) and a vegetable-rich pattern (high in vegetables, fruit, fish, and seaweed). The rice-rich pattern was not associated with baPWV or other risk factors. A significant inverse correlation was found between baPWV and the vegetable-rich dietary pattern ($p < 0.05$). **Conclusions:** Our findings suggest that a dietary pattern characterized by high consumption of vegetables, seafood, seaweed, fruit, and pulse is inversely associated with arterial stiffness measured by baPWV. This diet offers an additional nutritional approach to the prevention and treatment of arterial stiffness.

Key Words: diet, pattern, baPWV, Japan, risk

INTRODUCTION

Cardiovascular disease (CVD) is a major contributor to the burden of disease in most countries, and arteriosclerosis is one of many dominant risk factors for morbidity and mortality. These background factors facilitate arterial stiffness, and consequently, impaired viscoelasticity of the arterial wall is the leading cause of CVD.¹

Pulse wave velocity (PWV) is a known indicator of arterial stiffness,^{2,3} and brachial-ankle PWV (baPWV) is a simple and non-invasive method of measurement with considerably high validity and reproducibility.⁴⁻⁶ A previous study in a Japanese population showed that baPWV can predict all-cause mortality;⁷ therefore, baPWV can be used to assess vascular damage in this population.

Food intake is one of the most important modifiable risk factors for CVD.⁸ In Western countries, the Dietary Approaches to Stop Hypertension (DASH) dietary pattern, which emphasized the consumption of fruit, vegetables, and low-fat dairy and discouraged the consumption of meat, sugar-sweetened beverages, and saturated and to-

tal fat, has been shown to lower blood pressure, affect natriuresis and the renin-angiotensin-aldosterone system, reduce adrenergic tone, and increase vascular relaxation.⁹⁻¹¹ Additionally, the Mediterranean dietary pattern, which emphasizes the consumption of grains, vegetables, legumes and nuts, fruit, olive oil, dairy products, and fish, has been shown to reduce the risk of CVD and mortality, likely due to the anti-inflammatory properties of its foods (e.g., n-3 polyunsaturated fatty acids [PUFAs]).^{12,13} These studies commonly used a "dietary pattern" approach. Dietary intake is a complex exposure variable with a large number of components, each of which differently influ-

Corresponding Author: Shota Moyama, 1882 Hassaka-cho, Hikone, Shiga, 522-8539, Japan.

Tel: +81-749-22-6050; Fax: +81-749-26-0754

Email: shota0531@gmail.com

Manuscript received 15 November 2015. Initial review completed 30 November 2015. Revision accepted 03 January 2016.

doi: 10.6133/apjn.032016.14

ence the risk of disease (some protective, others harmful). Therefore, dietary pattern analysis is an effective method for examining the association between the overall diet and arterial stiffness.¹⁴ However, few studies have examined the association between dietary patterns and arterial stiffness, especially in Japan.

To test whether specific dietary patterns are associated with arterial stiffness, we investigated the risk factors related to arterial stiffness as well as dietary patterns among Japanese middle-aged adults. In addition, we examined how the individual food groups characterizing each dietary pattern relate to arterial stiffness.

MATERIALS AND METHODS

Study population

We studied 70 native Japanese subjects (39 men and 31 women) with no history of previously reported CVD or other serious diseases (including cancer, diabetes, drug abuse, or major psychiatric illness) that may have influenced arterial stiffness.¹⁵ Subjects using hypolipidemic drugs or estrogen-replacement therapy were excluded from the study, because these drugs are known to reduce PWV.¹⁵ Informed consent was obtained from all subjects. The study protocol was approved by the ethics committee of the University of Shiga Prefecture and Hikone Municipal Hospital and carried out in accordance with the principles of the Declaration of Helsinki.

Assessment of dietary intake

Food consumption was assessed using a 97-item food-frequency questionnaire (FFQ).^{16,17} The reproducibility and validity of nutrient and food group intake measurements from the FFQ used in this study have been described in detail elsewhere,^{16,17} and the FFQ was structured to reflect the food habits of the Japanese population. Each food/beverage item provided three to six possible responses related to frequency of consumption, ranging from “never or less than once a month” to “five or more times per day.” Participants indicated how often, on average, they had consumed a given amount of the specified food/beverage during the past month. The 97 individual food items captured by the FFQ were grouped together in

14 non-overlapping food groups (Table 1). The grouping scheme was based on the similarity of nutrient profiles or culinary usage among the foods and was somewhat similar to that used in other studies.^{9,18-22} Both dietitians and study subjects completed the FFQ, and the dietitians either administered the FFQ, or they reviewed the subjects’ responses. The amount of intake of each nutrient and food group was estimated using the FFQ analysis system (System Supply Co., Ltd).

Assessment of baPWV and other relevant variables

baPWV and blood pressure measurements were performed in the supine position after 5 minutes of bed rest using a validated automatic device (BP-203RPE 2 form PWV/ABI; OMRON Healthcare, Tokyo, Japan) in all subjects. Four oscillometric cuffs, each connected to a plethysmographic sensor that determined volume pulse form and an oscillometric pressure sensor that measured blood pressure, were wrapped on both the brachia and ankle, and two electrocardiogram electrodes were placed on each wrist. They were simultaneously pressurized to the approximate value of the patient’s diastolic pressure so that the pulse volume waveforms could be recorded using semiconductor pressure sensors. The distance between baPWV sampling points was calculated automatically according to the subject’s height. The path length from the suprasternal notch to the ankle (L_a) was calculated as $L_a = 0.8129 \times \text{height (in cm)} + 12.328$. The path length from the suprasternal notch to the brachium (L_b) was calculated as $L_b = 0.2195 \times \text{height} - 2.0734$. baPWV was calculated according to the following formula: $\text{baPWV} = (L_a - L_b) / T_{ba}$, where T_{ba} is the time interval between the wave front of the brachial and ankle waveforms.²³ Two simultaneous measurements of baPWV were recorded (on the right and left side), and the average of these readings was used as the representative value for each individual.

Previous studies have found that baPWV is influenced by age, systolic blood pressure (SBP), and gender.^{6,15} Yamashina et al devised nomograms of the relation between baPWV and blood pressure level and suggested that a comparison of measured baPWV with values calcu-

Table 1. Components of the 14 food groups used in the factor analysis[†]

| Food group | Components |
|--------------------|---|
| Rice | Pilaf, curry and rice, rice bowl dishes, sushi, rice balls |
| Breads | Prepared bread, sweet buns |
| Noodles | Wheat noodles, buckwheat noodles, somen, cold noodles, ramen, pan-fried noodles, spaghetti |
| Other carbohydrate | Okonomiyaki, gratin, cereal, pizza |
| Potatoes | Potato salad, french fries, croquette, other potato dishes |
| Dairy foods | Corn soup, cream stew, yogurt, cheese |
| Meats | Beef, chicken, pork, liver, ground meat dishes, processed meat |
| Pulses | Miso soup, tofu, natto, beans |
| Seafood | Saury, tuna, salmon, horse mackerel, bonito, mackerel, eel, capelin, sardine, salted cod roe, herring egg, squid, shrimp, oyster, fish sausage |
| Vegetables | Tomato, carrot, pumpkin, spinach, greens, bell pepper, broccoli, radish, burdock, lotus, cabbage, Chinese cabbage, egg apple, green bean, pickles |
| Seaweed | Mushroom, barilla, brown seaweed |
| Fruits | Citrus, apple, strawberry, persimmon, kiwi, banana, other fruits |
| Confectionery | Rice crackers, peanuts, cakes, steamed bean-jam bun, snack, pudding, ice cream |
| Oils and fats | Fritters, batter, margarine, dressing, mayonnaise |

[†]Each row represents a single food-frequency questionnaire item.

lated from the nomograms may help to avoid underestimating the real risk of atherosclerotic CVD, including the severity of atherosclerotic vascular damage, reflected by measured baPWV in subjects with different ages and blood pressure levels.¹⁵ The ratio with the values calculated from the nomograms and measured baPWV values is defined as $\% \Delta \text{baPWV} \{[(\text{measured value} - \text{calculated value}) / \text{calculated value}] * 100\}$. It has been reported that $\% \Delta \text{baPWV}$ was not significantly associated with age and SBP.²⁴

Total and high-density lipoprotein cholesterol (HDL-C) and triglyceride (TG) concentrations were measured in blood samples. Participants were asked to fast for 12 hours, avoid smoking on the morning of the examination, and avoid heavy exercise 12 hours before the examination. Low-density lipoprotein cholesterol (LDL-C) was calculated using the Friedewald equation.²⁵ Non-HDL-C and the LDL/HDL ratios were calculated using the formula (LDL/HDL ratio = LDL-C/HDL-C, non-HDL-C = total-cholesterol - HDL-C).

Statistical analysis

Statistical Analysis Software (Ekuseru-Toukei 2012, Social Survey Research Information Co., Ltd. Japan) was used to conduct data analyses. To identify dietary patterns in our study population, factor analysis (principal component method) was conducted. The factors were orthogonally transformed using varimax rotation to achieve a structure with independent (non overlapping) factors and greater interpretability. Components with an eigenvalue >1, Scree test results, and the interpretability of the factors were considered when determining the number of factors to retain.¹⁸⁻²² Food groups with absolute factor loadings ≥ 0.30 were considered as significant contributors to the pattern (Table 2). The factor loadings of the food groups by dietary patterns are provided in Table 2. Food groups with positive loadings contribute to the dietary pattern, whereas food groups with negative loadings are inversely associated with the dietary pattern.

We used a simple linear regression model to assess the relation between dietary patterns and arterial stiffness and

Table 2. Factor-loading matrix for dietary patterns[†]

| Foods | Dietary pattern | |
|---------------------|-----------------|----------------|
| | Rice-rich | Vegetable-rich |
| Rice | 0.70 | — |
| Breads | 0.64 | — |
| Fats and oils | 0.59 | — |
| Potatoes | 0.58 | — |
| Dairy products | 0.47 | — |
| Meats | 0.51 | — |
| Other carbohydrates | 0.46 | — |
| Confectionery | 0.64 | — |
| Noodles | 0.62 | — |
| Pulse | — | 0.49 |
| Fruits | — | 0.49 |
| Seaweed | — | 0.62 |
| Seafood | — | 0.66 |
| Vegetables | — | 0.72 |

[†]Absolute values of factor loadings <0.30 were omitted for simplicity. The food groups are presented in descending order of loading values on the rice-rich dietary pattern.

relevant variables. Data are expressed as the mean \pm standard deviation (SD). Differences among the four groups were evaluated using a one-way analysis of variance, and a *p* value <0.05 was considered statistically significant.

RESULTS

We identified two major dietary patterns using factor analysis. The first pattern was characterized by a high positive loading for rice, breads, oils and fats, potatoes, meat, other carbohydrates, confectionery, and noodles; because the largest contributor to this diet was rice, we named this dietary pattern the “rice-rich” pattern. In the second pattern, vegetables, seafood, seaweed, fruit, and pulse were positively correlated with pattern scores; because the largest contributor was vegetables, we named this dietary pattern the “vegetable-rich” pattern. Table 3 shows the characteristics of study subjects according to the quartiles for the rice-rich and vegetable-rich patterns.

Table 3. Characteristics of study subjects across quartile (Q) of dietary pattern score^{†‡}

| | Group | | | | <i>p</i> for trend [§] |
|---------------------------------|-----------------|-----------------|-----------------|-----------------|---------------------------------|
| | Q1 (n=18) | Q2 (n=18) | Q3 (n=17) | Q4 (n=17) | |
| Rice-rich pattern | | | | | |
| Age (y) | 61.9 \pm 9.7 | 58.2 \pm 10.1 | 53.3 \pm 11.2 | 55.2 \pm 9.7 | <0.05 |
| Height (cm) | 165 \pm 8.4 | 162 \pm 7.4 | 161 \pm 9.2 | 164 \pm 7.8 | 0.786 |
| Body weight (kg) | 59.8 \pm 12.2 | 62.1 \pm 11.3 | 61.0 \pm 10.0 | 60.9 \pm 10.6 | 0.996 |
| BMI (kg/m ²) | 21.7 \pm 2.9 | 23.6 \pm 2.9 | 23.3 \pm 3.0 | 22.6 \pm 3.7 | 0.988 |
| Systolic blood pressure (mmHg) | 129 \pm 16.2 | 128 \pm 9.0 | 121 \pm 13.7 | 122 \pm 14.9 | <0.05 |
| Diastolic blood pressure (mmHg) | 79.8 \pm 12.5 | 78.4 \pm 5.7 | 74.1 \pm 11.1 | 75.7 \pm 8.9 | 0.072 |
| Vegetable-rich pattern | | | | | |
| Age (y) | 54.8 \pm 13.0 | 56.6 \pm 9.8 | 55.5 \pm 10.6 | 61.9 \pm 6.9 | 0.059 |
| Height (cm) | 165 \pm 8.2 | 165 \pm 9.7 | 160 \pm 7.6 | 163 \pm 6.9 | 0.368 |
| Body weight (kg) | 64.5 \pm 13.3 | 62.0 \pm 10.2 | 56.6 \pm 10.7 | 60.5 \pm 7.8 | 0.188 |
| BMI (kg/m ²) | 23.7 \pm 4.3 | 22.7 \pm 2.6 | 21.9 \pm 2.9 | 22.9 \pm 2.4 | 0.329 |
| Systolic blood pressure (mmHg) | 127 \pm 18.7 | 124 \pm 10.3 | 120 \pm 12.3 | 128 \pm 12.0 | 0.871 |
| Diastolic blood pressure (mmHg) | 81.4 \pm 9.2 | 76.5 \pm 12.9 | 73.4 \pm 8.7 | 76.7 \pm 7.3 | 0.131 |

[†]Values are shown as mean \pm SD except for number of study subjects.

[‡]Q; quartile of factor score.

[§]*p*-value from linear regression analysis for quantitative variables and chi-square test for qualitative variables.

Table 4. Energy and nutrient intake of study subjects according to the 1st and 4th quartile (Q) of dietary pattern score^{†‡}

| | Rice-rich pattern | | | Vegetable-rich pattern | | |
|-------------------------|-------------------|------------|---------------------------------|------------------------|----------|---------------------------------|
| | Q1 | Q4 | <i>p</i> for trend [§] | Q1 | Q4 | <i>p</i> for trend [§] |
| Energy (kcal/d) | 1738±463 | 2508±421 | <0.001 | 2036±527 | 2239±509 | 0.055 |
| Protein (g) | 33.4±6.4 | 33.9±3.2 | 0.992 | 30.2±4.4 | 37.3±4.6 | <0.001 |
| Fat (g) | 22.9±5.6 | 31.1±4.7 | <0.001 | 25.0±7.3 | 27.9±4.6 | 0.056 |
| Carbohydrate (g) | 144±14.1 | 132±14.3 | <0.05 | 232±108 | 283±79.0 | 0.203 |
| Calcium (mg) | 273±95.3 | 245±75.1 | 0.443 | 456±211 | 632±238 | 0.062 |
| Magnesium (mg) | 144±23.8 | 118±17.9 | <0.01 | 117±21.9 | 150±24.5 | <0.01 |
| Potassium (mg) | 1313±278 | 1084±189.7 | <0.05 | 1045±288 | 1499±328 | <0.001 |
| Saturated fat (g) | 6.6±2.3 | 8.9±2.0 | <0.001 | 7.2±2.7 | 7.7±2.1 | 0.403 |
| Monounsaturated fat (g) | 7.6±2.1 | 10.8±1.8 | <0.001 | 8.6±2.7 | 9.5±1.7 | 0.156 |
| Polyunsaturated fat (g) | 5.4±1.1 | 6.7±1.0 | <0.001 | 5.5±1.4 | 6.6±1.2 | <0.01 |
| Fibre (g) | 6.1±2.0 | 5.5±1.1 | 0.297 | 4.7±1.2 | 8.0±1.9 | <0.001 |

[†]Values are shown as mean±SD which are adjusted for total energy intake (1,000kcal/day), except energy.

[‡]Q; quartile of factor score.

[§]*p*-value from linear regression analysis.

Table 5. Select cardiovascular disease risk factor across quartiles (Q) of dietary pattern score^{†‡}

| | Group | | | | <i>p</i> for trend [§] |
|------------------------|-----------|-----------|-----------|-----------|---------------------------------|
| | Q1 | Q2 | Q3 | Q4 | |
| Rice-rich pattern | | | | | |
| TC (mg/dL) | 210±37.6 | 212±43.4 | 200±30.4 | 214±27.6 | 0.701 |
| TG (mg/dL) | 103±49.5 | 111±52.9 | 94±37.2 | 122±46.8 | 0.662 |
| LDL-C (mg/dL) | 131±27.9 | 134±33.8 | 119±30.2 | 133±23.6 | 0.942 |
| HDL-C (mg/dL) | 58.6±16.4 | 55.9±15.0 | 61.8±11.0 | 57.5±15.0 | 0.466 |
| nonHDL-C (mg/dL) | 152±30.6 | 157±37.3 | 138±31.8 | 158±26.5 | 0.791 |
| LDL/HDL ratio | 2.4±0.7 | 2.5±0.7 | 2.0±0.8 | 2.5±0.7 | 0.719 |
| %ΔbaPWV (%) | 4.9±16.3 | 5.5±10.8 | 2.5±10.6 | 7.3±7.9 | 0.656 |
| Vegetable-rich pattern | | | | | |
| TC (mg/dL) | 207±37.2 | 208±31.4 | 196±36.1 | 224±33.0 | 0.206 |
| TG (mg/dL) | 118±49.1 | 101±47.3 | 118±54.9 | 94±34.8 | 0.353 |
| LDL-C (mg/dL) | 129±32.3 | 126±28.9 | 119±25.7 | 143±26.9 | 0.333 |
| HDL-C (mg/dL) | 54.9±14.4 | 62.2±15.4 | 53.6±14.4 | 62.9±11.7 | 0.107 |
| nonHDL-C (mg/dL) | 154±36.3 | 146±31.4 | 143±30.3 | 161±30.0 | 0.569 |
| LDL/HDL ratio | 2.6±0.9 | 2.1±0.8 | 2.3±0.6 | 2.3±0.6 | 0.263 |
| %ΔbaPWV (%) | 9.9±14.5 | 3.6±11.0 | 4.2±8.1 | 2.3±11.7 | <0.05 |

TC, total cholesterol; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; PWV, pulse wave velocity.

[†]All values are shown as mean±SD.

[‡]Q; quartile of factor score.

[§]*p*-value from linear regression analysis for quantitative variables.

All components presented a linear trend through the quartiles, with increasing or decreasing trend. Persons in the upper quartiles of the rice-rich pattern were more likely to be young and have low systolic blood pressure (*p* for trend <0.05 for all), whereas no correlation was found between the vegetable-rich pattern and any of the risk factors or variables analyzed.

Table 4 shows the nutrient composition (energy adjusted, in 1,000 kcal/d) of each dietary pattern. Persons with high rice-rich pattern scores consumed more total fat, saturated fat, monounsaturated fat, and PUFA (*p* for trend <0.001 for all) and less potassium and magnesium than persons with lower scores (*p* for trend <0.05 and <0.01, respectively). On the other hand, the vegetable-rich pattern generally showed trends opposite that of the rice-rich pattern in that these subjects consumed more protein, magnesium, potassium, PUFA, and fibre (*p* for trend

<0.001 for all, except for magnesium and PUFA [*p*<0.01]). The CVD risk factors across quartiles of dietary pattern scores are summarized in Table 5. Although we observed no association between dietary patterns and the major CVD risk factors (TC, TG, LDL-C, HDL-C, non-HDL-C, and LDL/HDL ratio), %ΔbaPWV significantly decreased in subjects reporting a vegetable-rich dietary pattern (*p* for trend <0.05), however the trend in %ΔbaPWV was not statistically significant in subjects reporting a rice-rich dietary pattern.

DISCUSSION

In this cross-sectional study of Japanese middle-aged adults, we identified two major dietary patterns using factor analysis. The vegetable-rich pattern was characterized by the consumption of a wide variety of foods, such as vegetables, seafood, seaweed, fruit, and pulse. The

rice-rich pattern included rice, breads, oils and fats, potatoes, meats, other carbohydrates, confectionery, and noodles. We observed an inverse association between the vegetable-rich pattern factor score and arterial stiffness derived from baPWV.

Several studies have examined the relationship between a single nutrient or a few nutrients or foods and CVD risk factors. Lopez et al examined the association between a diet rich in vegetables, fruit, and fish and inflammation markers such as C-reactive protein and E-selectin and found an inverse relation after adjusting for age, body mass index, physical activity, smoking status, and average alcohol consumption.¹⁹ Fukuoka et al reported that a fish-based diet effectively prevented an increase in baPWV in low-risk coronary artery disease patients.²⁶ Fish is a major source of omega-3 fatty acids, eicosapentaenoic acid (EPA), and docosahexaenoic acid (DHA), which are considered promising treatments for the prevention of major coronary events, especially non-fatal ones, in the Japanese population.²⁷ The Japan EPA lipid intervention study (JELIS) and the Japan public health center-based (JPHC) study showed that the consumption of fish and n-3 PUFA reduced the incidence of major coronary events.^{27,28} Aatola et al reported that vegetable consumption was the only lifestyle risk factor that was an independent predictor of PWV when adjusted for lifestyle or traditional risk factors.²⁹ In another report, high fruit and vegetable consumption was related to a reduced CVD risk in prospective settings among middle-aged and elderly subjects.³⁰

Our study provides evidence of a significant inverse association between consumption of a vegetable-rich pattern consisting of a variety of foods and arterial stiffness among Japanese middle-aged adults. In our analysis of individual food items, we confirmed the inverse association between arterial stiffness and the consumption of vegetables, seafood, seaweed, fruit, and pulse reported in earlier studies.²⁶⁻³⁰ The individual consumption of vegetables, fish, and seafood was not associated with arterial stiffness, which were the major food items characterizing the vegetable-rich pattern, although these foods are considered elements of a healthy diet. Wahlqvist et al reported that food variety is associated with less macrovascular disease,³¹ and the increased variety among plant foods eaten might be predictive of better arterial compliance. Additionally, consumption of fish rich in n-3 PUFAs, has been previously found to be protective against arterial stiffness and related risk factors,^{27,28} which might explain the beneficial effects of the vegetable-rich pattern on arterial stiffness. Furthermore, we postulate benefits from other constituents of the vegetable-rich pattern including monounsaturated fatty acids, fibre, high-quality protein, low glycemic index, vitamins, minerals, and other bioactive compounds and that the rice-rich pattern included limited amounts of these constituents.

The strengths of the present study included its population-based setting, the use of standardized methods to identify outcomes, and consideration of a broad range of confounding variables. However, several limitations should be considered while interpreting our findings. First, the cross-sectional design made it difficult to examine the causal relationships between dietary patterns and arterial

stiffness; therefore, further prospective or intervention studies are necessary to confirm our findings. Second, several assumptions were made when constructing food groups (e.g., when mixed dishes were aggregated into component food groups and when questions containing nutritionally diverse foods were categorized). However, our dietary patterns and loading scores were similar to those reported in other studies, and associations between dietary patterns and arterial stiffness were as expected; this suggests that our assumptions were valid.^{9,18-22} Third, regardless of our findings, which are based on empirically derived dietary patterns, it is possible that other food combinations may be more strongly related to arterial stiffness and other relevant variables. However, we found no independent relationships between arterial stiffness and food groups that did not have high factor loadings for any dietary pattern.

In conclusion, this is the first study to demonstrate a relationship between dietary patterns and arterial stiffness in Japanese middle-aged adults. Our findings suggest that a dietary pattern characterized by a high consumption of vegetables, seafood, seaweed, and fruit is inversely associated with arterial stiffness. Such a diet offers an additional nutritional approach to the prevention and treatment of arterial stiffness.

ACKNOWLEDGMENTS

This study was supported by The University of Shiga prefecture of Japan and Hikone Municipal Hospital of Japan as part of the "Hikone study" research project.

AUTHOR DISCLOSURES

None of the authors have any personal or financial conflicts of interest.

REFERENCES

1. Mattace-Raso FU, van der Cammen TJ, Hofman A, van Popele NM, Bos ML, Schalekamp MA et al. Arterial stiffness and risk of coronary heart disease and stroke. The Rotterdam Study. *Circulation*. 2006;113:657-63. doi: 10.1161/CIRCULATIONAHA.105.555235.
2. DeLoach S, Townsend R. Vascular stiffness: its measurement and significance for epidemiologic and outcome studies. *Clin J Am Soc Nephrol*. 2008;3:184-92. doi: 10.2215/CJN.03340807.
3. Hansen TW, Staessen JA, Torp-Pedersen C, Rasmussen S, Thijs L, Ibsen H, Jeppesen J. Prognostic value of aortic pulse wave velocity as index of arterial stiffness in the general population. *Circulation*. 2006;113:664-70. doi: 10.1161/CIRCULATIONAHA.105.579342.
4. Yufu K, Takahashi N, Anan F, Hara M, Yoshimatsu H, Saikawa T. Brachial arterial stiffness predicts coronary atherosclerosis in patients at risk for cardiovascular diseases. *Jpn Heart J*. 2004;45:231-42. doi: 10.1536/jhj.45.231.
5. Tsuchikura S, Shoji T, Kimoto E, Shinohara K, Hatsuda S, Koyama H, Emoto M, Nishizawa Y. Brachial-ankle pulse wave velocity as an index of central arterial stiffness. *J Atheroscler Thromb*. 2010;17:2-9. doi: 10.5551/jat.3616.
6. Yamashina A, Tomiyama H, Takeda K, Tsuda H, Arai T, Hirose K, Koji S, Hori S, Yamamoto Y. Validity, reproducibility, and clinical significance of noninvasive brachial-ankle pulse wave velocity measurement. *Hypertens Res*. 2002;25:359-64. doi: 10.1291/hyres.25.359.
7. Turin TC, Kita Y, Rumana N, Takashima N, Kadota A, Matsui K et al. Brachial-ankle pulse wave velocity predicts

- all-cause mortality in the general population: findings from the Takashima study, Japan. *Hypertens Res.* 2010;33:922-5. doi: 10.1038/hr.2010.103.
8. Ignarro LJ, Balestrieri ML, Napoli C. Nutrition, physical activity, and cardiovascular disease: an update. *Cardiovasc Res.* 2007;73:326-40. doi: 10.1016/j.cardiores.2006.06.030.
 9. Lin PH, Allen JD, Li YJ, Yu M, Lien LF, Svetkey LP. Blood pressure-lowering mechanism of the DASH dietary pattern. *J Nutr Metab.* 2012;2012:1-10. doi: 10.1155/2012/472396.
 10. Ai-Solaiman Y, Jesri A, Mountfold WK, Lackland DT, Zhao Y, Egan BM. DASH lowers blood pressure in obese hypertensive beyond potassium, magnesium and fiber. *J Hum Hypertens.* 2010; 24:237-46. doi: 10.1038/jhh.2009.58.
 11. Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM et al. A clinical trial of the effects of dietary patterns on blood pressure. *N Engl J Med.* 1997;336:1117-24. doi: 10.1056/NEJM199704173361601.
 12. Casas R, Sacanella E, Urpi-Sarda M, Chiva-Blanch G, Ros E, Martinez-Gonzalez MA et al. The effects of the Mediterranean diet on biomarkers of vascular wall inflammation and plaque vulnerability in subjects with high risk for cardiovascular disease. A randomized trial. *PLoS One.* 2014;9:e100084. doi: 10.1371/journal.pone.0100084.
 13. Sofi F, Macchi C, Abbate R, Gensini GF, Casini A. Mediterranean diet and health status: an updated meta-analysis and a proposal for a literature-based adherence score. *Public Health Nutr.* 2014;17:2769-82. doi: 10.1017/S1368980013003169.
 14. Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol.* 2002;13:3-9. doi: 10.1097/00041433-200202000-00002.
 15. Yamashina A, Tomiyama H, Arai T, Koji Y, Yambe M, Motobe H, Gluniza Z, Yamamoto Y, Hori S. Nomogram of the relation of brachial-ankle pulse wave velocity with blood pressure. *Hypertens Res.* 2003;26:801-6. doi: 10.1291/hypr.26.801.
 16. Wakai K, Egami I, Kato K, Lin Y, Kawamura T, Tamakoshi A et al. A simple food frequency questionnaire for Japanese diet: part 1. Development of the questionnaire, and reproducibility and validity for food groups. *J Epidemiol.* 1999;9:216-26. doi: 10.2188/jea.9.216.
 17. Egami I, Wakai K, Kato K, Lin Y, Kawamura T, Tamakoshi A et al. A simple food frequency questionnaire for Japanese diet: part 2. Reproducibility and validity for nutrient intakes. *J Epidemiol.* 1999;9:227-34. doi: 10.2188/jea.9.227.
 18. Menotti A, Alberti-Fidanza A, Fidanza F, Lanti M, Fruttini D. Factor analysis in the identification of dietary patterns and their predictive role in morbid and fatal events. *Public Health Nutr.* 2012;15:1232-9. doi: 10.1017/S1368980011003235.
 19. Lopez-Garcia E, Schulze MB, Fung TT, Meigs JB, Rifai N, Manson JE, Hu FB. Major dietary patterns are related to plasma concentrations of the markers of inflammation and endothelial dysfunction. *Am J Clin Nutr.* 2004;80:1029-35.
 20. Nettleton JA, Schulze MB, Jiang R, Jenny NS, Burke GL, Jacobs DR. A priori-defined dietary patterns and markers of cardiovascular disease risk in the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr.* 2008;88:185-94.
 21. Nettleton JA, Steffen LM, Mayer-Davis EJ, Jenny NS, Jiang R, Herrington DM, Jacobs DR. Dietary patterns are associated with biochemical markers of inflammation and endothelial activation in the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr.* 2006;83:1369-79.
 22. Iqbal R, Anand S, Ounpuu S, Islam S, Zhang X, Rangarajan S, Chifamba J, Ai-Hinai A, Keltai M, Yusuf S. Dietary patterns and the risk of acute myocardial infarction in 52 countries. *Circulation.* 2008;118:1929-37. doi: 10.1161/CIRCULATIONAHA.107.738716.
 23. Katakami N, Osonoi T, Takahara M, Saitou M, Matsuoka T, Yamasaki Y, Shimoura I. Clinical utility of brachial-ankle pulse wave velocity in the prediction of cardiovascular events in diabetic patients. *Cardiovasc Diabetol.* 2014;13:128. doi: 10.1186/s12933-014-0128-5.
 24. Minami K, Moyama S, Yoshida R, Teramura Y, Yano H, Yamada A, Hukui T, Takayama H, Hayashi S, Yorimoto A. Effects of exercise focusing on the walking for arterial stiffness in middle age adults. *Walking Res.* 2011;15:93-8.
 25. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem.* 1972;18:499-502.
 26. Fukuoka Y, Nuruki N, Amiya S, Tofuku K, Aosaki S, Tsubouchi H. Effects of a fish-based diet and administration of pure eicosapentaenoic acid on brachial-ankle pulse wave velocity in patients with cardiovascular risk factors. *J Cardiol.* 2014;63:211-7. doi: 10.1016/j.jjcc.2013.08.005.
 27. Yokoyama M, Origasa H, Matsuzaki M, Matsuzawa Y, Saito Y, Ishikawa Y et al. Effects of eicosapentaenoic acid on major coronary events in hypercholesterolaemic patients (JELIS): a randomized open-label, blinded endpoint analysis. *Lancet.* 2007;369:1090-8. doi: 10.1016/S0140-6736(07)60527-3.
 28. Iso H, Kobayashi M, Ishihara J, Sasaki S, Okada K, Kita Y, Kokubo Y, Tsugane S. Intake of fish and n3 fatty acids and risk of coronary heart disease among Japanese. The Japan Public Health Center-Based (JPHC) Study Cohort 1. *Circulation.* 2006;113:195-202. doi: 10.1161/CIRCULATIONAHA.105.581355.
 29. Aatola H, Koivisto T, Hutri-Kahonen N, Juonala M, Mikkilä V, Lehtimäki T, Viikari JSA, Raitakari OT, Kahonen M. Lifetime fruit and vegetable consumption and arterial pulse wave velocity in adulthood. The Cardiovascular Risk in Young Finns Study. *Circulation.* 2010;122:2521-8. doi: 10.1161/CIRCULATIONAHA.110.969279.
 30. Nagura J, Iso H, Watanabe Y, Maruyama K, Date C, Toyoshima H et al. Fruit, vegetable and bean intake and mortality from cardiovascular disease among Japanese men and women: the JACC Study. *Br J Nutr.* 2009;102:285-92. doi: 10.1017/S0007114508143586.
 31. Wahlqvist ML, Lo CS, Myers KA. Food variety is associated with less macrovascular disease in those with type 2 diabetes and their healthy controls. *J Am Coll Nutr.* 1989;8:515-23. doi: 10.1080/07315724.1989.10720321.