

This author's PDF version corresponds to the article as it appeared upon acceptance. Fully formatted PDF versions will be made available soon.

Safety and benefit of pre-operative oral carbohydrate in infants: A multi-center study in China

doi: 10.6133/apjcn.052018.08

Published online: May 20158

Running title: Oral carbohydrates in pre-operative infants

Weiwei Jiang MD¹, Xiang Liu MSc², Fengli Liu MSc³, Shungen Huang MD⁴, Jie Yuan MSc⁵, Yingzuo Shi MSc⁶, Huan Chen MSc¹, Jie Zhang MD¹, Changgui Lu MD¹, Wei Li MSc¹, Qiming Geng BSc¹, Xiaoqun Xu BSc¹, Weibing Tang MD¹

¹Department of Pediatric Surgery, Children's Hospital of Nanjing Medical University, Nanjing, China

²Anhui Provincial Children's Hospital, Hefei, China

³Xuzhou Children's Hospital, Xuzhou, China

⁴Children's Hospital of Soochow University, Suzhou, China

⁵Changzhou Children's Hospital, Changzhou, China

⁶Wuxi Children's Hospital, Wuxi, China

Authors' email addresses and contributions:

Weiwei Jiang, wwjiang@njmu.edu.cn;

Xiang Liu, 84230331@qq.com;

Fengli Liu, 454509892@qq.com;

Shungen Huang, huangshungen@aliyun.com;

Jie Yuan jie, yuan1@aliyun.com;

Yingzuo Shi, shiyingzuo@aliyun.com

; Huan Chen, 1933258744@qq.com;

Jie Zhang, 36816496@qq.com;

Changgui Lu, 45087534@qq.com;

Qiming Geng, gqm080808@163.com;

Xiaoqun Xu, xiaoqunxu@aliyun.com;

Weibing Tang, tangweibingcn@163.com

Authors' contributions:

Weiwei Jiang: analyzed data and wrote the manuscript; Weibing Tang: designed the research; Xiang Liu, Fengli Liu, Shungen Huang, Jie Yuan, Yingzuo Shi, Huan Chen, Jie Zhang, Changgui Lu, and Wei Li: collected data; Qiming Geng: edited the article; Xiaoqun Xu: participated in discussion; Weibing Tang: reviewed the manuscript

Corresponding Author: Dr Weibing Tang, Department of Pediatric Surgery, Children's Hospital of Nanjing Medical University, 72 Guangzhou Road, Nanjing 210008, China. Tel: +86-025-83117231. Email: tangweibingcn@163.com; 68824153@qq.com

ABSTRACT

Background and Objectives: Pre-operative oral carbohydrate administration (POCA) is an important aspect of enhanced recovery after surgery and has many advantages. The objective of this study was to explore the safety and effect of pre-operative oral carbohydrate administration in infants. **Methods and Study Design:** This was a prospective, multi-center, randomized study that randomly assigned 1200 infants into four groups. In the control group (group A), the infants were strictly restricted to 6-h preoperative fasting before anesthesia. In the enhanced recovery after surgery (ERAS) groups (groups B, C, and D), the infants were orally administered a 10% carbohydrate solution (10% glucose water; 5, 10, and 15 mL/kg, respectively) 2 h before anesthesia. Blood glucose, gastric residual volumes, crying ratios, and the length of hospital stay were observed. **Results:** The blood glucose was significantly higher in groups B, C, and D than group A at the time of anesthesia. The gastric residual volume revealed virtually no residue in groups A, B, and C, but 15 infants in group D had a gastric residual volume. The crying ratio was significantly higher in group A. The length of hospital stay was not significantly different between the groups. **Conclusions:** POCA is well-tolerated in infants at a dose of 10 mL/kg.

Key Words: enhanced recovery after surgery, preoperative oral carbohydrate, infants, blood glucose, gastric residual

INTRODUCTION

Kehlet and Wilmore¹ were the first to describe rapid rehabilitation after surgery. Since the turn of the century, the concept of enhanced recovery after surgery (ERAS) or rapid post-operative rehabilitation has been standardized.² ERAS is a combination of pre-, peri-, and post-operative measures in an attempt to accelerate post-operative recovery, and reduce the surgical stress response, post-operative morbidity, length of hospital stay, and complications.³⁻¹⁰ ERAS mainly includes pre-operative oral carbohydrate administration (POCA), limited peri-operative fasting, early enteral intake, early mobilization, and limited use of surgical drains and tubes.¹¹⁻¹⁵

POCA is an important aspect of the ERAS program, which appears to have many advantages.^{16,17} The conventional approach entails long periods of fasting, thus liquid or solid intake is prohibited 6–8 h before a planned procedure. This long fasting time has recently been questioned in favor of breaking the fast by POCA, which is more comfortable for the patients.¹⁸

POCA has gradually been applied to adult surgery. Recently, many articles have been published on POCA in adult surgical populations, such as elective abdominal surgery,¹⁹ cardiac surgery,²⁰ thyroidectomy,²¹ and orthopedic surgery.^{22,23} Despite its many benefits, POCA has not been universally adopted in pediatric surgery. Moreover, POCA data from pediatric populations is limited. The use of POCA in pediatric surgery is still in the initial stages and is in need of additional research.

The objective of this study was to explore the safety and benefit of POCA in the pediatric population, and the effect of POCA on pre- and peri-operative blood glucose in infants.

MATERIALS AND METHODS

Patients

This was a prospective, multi-center, randomized study. We studied consecutive patients who underwent surgery in the Department of Pediatric Surgery at Children's Hospital of Nanjing Medical University, Anhui Provincial Children's Hospital, Xuzhou Children's Hospital, Children's Hospital of Soochow University, Changzhou Children's Hospital, and Wuxi Children's Hospital between 1 March 2014 and 31 January 2017. The inclusion criteria were an infant who could eat normally before the operation, such as a patient with Hirschsprung's disease, sacrococcygeal teratoma, biliary atresia, or undergoing second stage operation for congenital anus atresia. The exclusion criteria were patients with emergency ileus, enterobrosis, or neonatal necrotizing enterocolitis.

The study was approved by the Ethics Committee of Children's Hospital affiliated with Nanjing Medical University (201601005-1.1, 201701025), multi-center research registration number (NCT02776176). Informed consent was obtained from the parents of all children, and the study was conducted in accordance with the Declaration of Helsinki.

Peri-operative care

The 1200 infants were randomly assigned to four groups. In the control group (group A), the patients were strict implementation of preoperative fasting at 6 hours before anaesthesia. In the ERAS groups (groups B–D), the patients were orally administered a 10% carbohydrate solution 2 h before anesthesia; in groups B, C, and D the patients received 10% glucose water at 5, 10, and 15 mL/kg, respectively.

Observation indices

The gastric juice residual was determined by pre-operative gastrointestinal decompression. Blood glucose (2 h before anesthesia, at the time of anesthesia, 2 h post-operatively, and 12 h post-operatively) was determined using a glucose meter (Bayer Healthcare LLC, Mishawaka, IN, USA). Crying ratios (when the duration of crying was >15 min, or the paroxysmal crying time was >1 h) and lengths of hospital stays were also measured.

Statistical analysis

Statistical analysis was performed with SPSS software (version 20.0; SPSS, Inc., Chicago, IL, USA). A Pearson chi-square test was used to compare the crying ratio between the groups. Analysis of variance was used to compare multi-group variables.

RESULTS

Clinical characteristics

The characteristics of the patients in groups A, B, C, and D are shown in Table 1. There were no significant differences with respect to gender, age, or type of disease in groups A, B, C, and D.

Blood glucose

Blood glucose were not significantly different between the groups 2h before anesthesia, 2h post-operatively, and 12h post-operatively ($p>0.05$), but were significantly higher in groups B, C, and D than group A at the time of anesthesia ($p <0.05$) (Figure1).

Gastric residual

Nearly no residue was measured in groups A, B, and C, but there were 15 patients in group D given the 10% oral carbohydrate solution with gastric residual volume.

Crying ratio peri-operatively

The crying ratio was significantly higher in group A than groups B, C, and D. ($p<0.05$) (Figure2).

Hospital stay

The length of hospital stay was not significantly different between the groups (Figure 3).

DISCUSSION

Following traditional surgery, patients often experience long periods of fasting, mobilization is limited, and the hospital stay is long, all of which are now gradually changing based on the concept of ERAS. Traditional thinking is that long periods of fasting reduce the risk of aspiration during surgery; however, this may let the patient feel thirst and hunger and increases anxiety and resistance to some clinically-relevant stressors which are involved in surgical stress.²⁴ Thus, are long periods of fasting before surgery really needed?

The typical gastric emptying time is 1.5 h, so POCA 2 h before surgery might be safe and without risk of pulmonary aspiration.²⁵ POCA is defined as an isotonic liquid rich in carbohydrates administered orally in the evening before surgery and 2 h before anesthesia.²⁷ A number of studies have demonstrated the risk of regurgitation or aspiration was no increase in POCA, so POCA is safe in adults and children.^{16,27,28} In the current study involving infants, there was almost no gastric residual in the POCA-5 mL and -10 mL groups, and 5% of infants had 10% gastric residual in the POCA-15 mL group. At doses of 5 and 10 mL/kg, oral POCA did not increase gastric residuals; as a result, in this study vomiting/aspiration was not observed. Therefore, we think that 10 mL/kg is a safe dose in infants.

In adults, POCA may reduce pre-operative discomfort, such as hunger and thirst. POCA may also reduce postoperative catabolism and insulin resistance, which leads to an alleviation of metabolic stress responses, an reduced postoperative vomiting, pain, and anxiety.²⁹ Moreover, POCA prevents muscle loss of nitrogen and protein, and maintains nitrogen balance.^{30,31} In our study, the rate of patient crying was significantly lower in the POCA groups pre-operatively. Thirst and hunger may be the main cause of crying and dysphoria, so POCA is likely to reduce thirst and hunger and reduce crying. At the same time, a reduction in crying can reduce pre-operative energy consumption. Moreover, pre-operative crying causes the accumulation of large amounts of gas in the gastrointestinal tract. During laparoscopic surgery, gastrointestinal flatulence leads to peritoneal space narrowing and will affect the surgery or interfere with the surgical procedure.

In adult studies, POCA has been shown to alter patients' metabolic state from a long-term fast, avoiding decreased glycogen reserves, which raise the body's ability to mount stressresponse.^{32,33} Moreover, POCA reduces postoperative discomfort, preserves muscle and respiratory function, and accelerates the recovery process; thus, some studies have shown that POCA can reduce the number of days in hospital.^{33,34} However, other studies have demonstrated that POCA had no effect on the length of hospital stays.³⁵⁻³⁷ Based on these different points of view regarding the effect of POCA on the length of hospital stay, we

studied the POCA on the length of hospital stay following pediatric surgery. The results showed that the difference in the length of hospital stay was not significant between the groups.

Blood glucose was significantly higher in the POCA group at the time of anesthesia, but was not significantly different at other time points. Therefore, POCA did not increase the post-operative blood glucose. Peri-operative trauma, stress, and long fasting can cause metabolic disorders, blood glucose fluctuation, and POCA can reduce blood glucose fluctuation before and during surgery, and reduce metabolic abnormalities. These indicators had significant changes in the peri-operative period of adults, but some indicators were not significantly changed in infants. Perhaps it is time for a change, as we have demonstrated that POCA will benefit infants.

In conclusion, POCA in infants is well tolerated at a dose of 10 mL/kg and was not associated with increased risks of vomiting/aspiration during surgery.

ACKNOWLEDGEMENTS

We thank Professor Fei Jian for providing anesthesia support.

AUTHOR DISCLOSURE

The authors declare no conflict of interest. This study was supported by the National Natural Science Foundation of China (81100318), the Jiangsu Provincial key research and development program (BE2017609), Jiangsu youth medical talent project (QNRC2016081), and Children's Hospital of Nanjing Medical University youth medical talent project (ETYYQM2014009).

REFERENCES

1. Kehlet H, Wilmore DW. Evidence-based surgical care and the evolution of fast-track surgery. *Ann Surg.* 2008;248:189-98. doi: 10.1097/SLA.0b013e31817f2c1a.
2. Fearon KC, Ljungqvist O, Von Meyenfeldt M, Revhaug A, Dejong CH, Lassen K et al. Enhanced recovery after surgery: a consensus review of clinical care for patients undergoing colonic resection. *Clin Nutr* 2005;24:466-77. doi:10.1016/j.clnu.2005.02.002
3. H. Kehlet. Fast-track colorectal surgery. *Lancet* 2008;371:791-3. doi: 10.1016/S0140-6736(08)60357-8
4. Auyong DB, Allen CJ, Pahang JA, Clabeaux JJ, MacDonald KM, Hanson NA. Reduced length of hospitalization in primary total knee arthroplasty patients using an updated enhanced recovery after orthopedic surgery (ERAS) pathway. *J Arthroplasty* 2015;30:1705-9. doi: 10.1016/j.arth.2015.05.007.

- 5 Miller TE, Thacker JK, White WD, Mantyh C, Migaly J, Jin J, Roche AM, Eisenstein EL, Edwards R, Anstrom KJ, Moon RE, Gan TJ. Reduced length of hospital stay in colorectal surgery after implementation of an enhanced recovery protocol. *Anesth Analg* 2014;118:1052-61. doi: 10.1213/ANE.000000000000206.
- 6 Kagedan DJ, Ahmed M, Devitt KS, Wei AC. Enhanced recovery after pancreatic surgery: a systematic review of the evidence. *HPB (Oxford)* 2015;17:11-6. doi: 10.1111/hpb.12265.
- 7 Tang J, Humes DJ, Gemmil E, Welch NT, Parsons SL, Catton JA. Reduction in length of stay for patients undergoing oesophageal and gastric resections with implementation of enhanced recovery packages. *Ann R Coll Surg Engl* 2013;95:323-8. doi: 10.1308/003588413X13629960046039.
- 8 Thiele RH, Rea KM, Turrentine FE, Friel CM, Hassinger TE, McMurry TL et al. Standardization of care: impact of an enhanced recovery protocol on length of stay, complications, and direct costs after colorectal surgery. *J Am Coll Surg* 2015;220:430-43. doi: 10.1016/j.jamcollsurg.2014.12.042.
- 9 He F, Lin X, Xie F, Huang Y, Yuan R. The effect of enhanced recovery program for patients undergoing partial laparoscopic hepatectomy of liver cancer. *Clin Transl Oncol* 2015;17:694-701. doi: 10.1007/s12094-015-1296-9.
- 10 Adamina M, Kehlet H, Tomlinson GA, Senagore AJ, Delaney CP. Enhanced recovery pathways optimize health outcomes and resource utilization: a meta-analysis of randomized controlled trials in colorectal surgery. *Surgery* 2011;149:830-840. doi: 10.1016/j.surg.2010.11.003.
- 11 Gustafsson UO, Scott MJ, Schwenk W, Demartines N, Roulin D, Francis N et al. Guidelines for perioperative care in elective colonic surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *World J Surg* 2013;37:259-84. doi:10.1007/s00268-012-1772-0
- 12 Mortensen K, Nilsson M, Slim K, Schäfer M, Mariette C, Braga M, Carli F, Demartines N, Griffin SM, Lassen K. Consensus guidelines for enhanced recovery after gastrectomy: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *Br J Surg* 2014;101:1209-29. doi: 10.1002/bjs.9582.
- 13 Cerantola Y, Valerio M, Persson B, Jichlinski P, Ljungqvist O, Hubner M et al. Guidelines for perioperative care after radical cystectomy for bladder cancer: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *Clin Nutr* 2013;32:879-87. doi: 10.1016/j.clnu.2013.09.014.
- 14 Lassen K, Coolen MME, Slim K, Carli F, de Aguiar-Nascimento JE, Schäfer M et al. Guidelines for perioperative care for pancreaticoduodenectomy: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *World J Surg* 2013;37:240-58. doi: 10.1007/s00268-012-1771-1
- 15 Nygren J, Thacker J, Carli F, Fearon KC, Norderval S, Lobo DN, Ljungqvist O, Soop M, Ramirez J. Guidelines for perioperative care in elective rectal/pelvic surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *World J Surg* 2013;37:285-305. doi: 10.1007/s00268-012-1787-6
- 16 Gawecka A, Mierzevska-Schmidt M. Tolerance of, and metabolic effects of, preoperative oral carbohydrate administration in children - a preliminary report. *Anaesthesiol Intensive Ther.* 2014;46:61-4. doi: 10.5603/AIT.2014.0013.
- 17 Tsutsumi R, Kakuta N, Kadota T, Oyama T, Kume K, Hamaguchi E, Niki N, Tanaka K, Tsutsumi YM. Effects of oral carbohydrate with amino acid solution on the metabolic status of patients in the

- preoperative period: a randomized, prospective clinical trial. *J Anesth.* 2016;30:842-9. doi: 10.1007/s00540-016-2217-y.
- 18 Awad S, Varadhan KK, Ljungqvist O, Lobo DN. A meta-analysis of randomised controlled trials on preoperative oral carbohydrate treatment in elective surgery. *Clin Nutr* 2013;32:34-44. doi: 10.1016/j.clnu.2012.10.011.
 - 19 Peixe-Machado PA, de Oliveira BD, Dock-Nascimento DB, de Aguilar-Nascimento JE. Shrinking preoperative fast time with maltodextrin and protein hydrolysate in gastrointestinal resections due to cancer. *Nutrition.* 2013;29:1054-9. doi: 10.1016/j.nut.2013.02.003.
 - 20 Breuer JP, von Dossow V, von Heymann C, Griesbach M, von Schickfus M, Mackh E et al. Preoperative oral carbohydrate administration to ASA III-IV patients undergoing elective cardiac surgery. *Anesth Analg.* 2006;103:1099-108. DOI:10.1213/01.ane.0000237415.18715.1d.
 - 21 Lauwick SM, Kaba A, Maweja S, Hamoir EE, Joris JL. Effects of oral preoperative carbohydrate on early postoperative outcome after thyroidectomy. *Acta Anaesthesiol Belg.* 2009;60:67-73.
 - 22 Harsten A, Hjartarson H, Toksvig-Larsen S. Total hip arthroplasty and perioperative oral carbohydrate treatment: a randomised, double-blind, controlled trial. *Eur J Anaesthesiol.* 2012;29:271-4. doi: 10.1097/EJA.0b013e3283525ba9.
 - 23 Ljunggren S, Hahn RG. Oral nutrition or water loading before hip replacement surgery: a randomized clinical trial. *Trials.* 2012;13:97. doi: 10.1186/1745-6215-13-97.
 - 24 Longchamp A, Harputlugil E, Corpataux JM, Ozaki CK, Mitchell JR. Is Overnight Fasting before Surgery Too Much or Not Enough? How Basic Aging Research Can Guide Preoperative Nutritional Recommendations to Improve Surgical Outcomes: A Mini-Review. *Gerontology.* 2017;63:228-37. doi: 10.1159/000453109.
 - 25 Nygren J, Thorell A, Jacobsson H, Larsson S, Schnell PO, Hylen L, Ljungqvist O. Preoperative gastric emptying. Effects of anxiety and oral carbohydrate administration. *Ann Surg* 1995;222:728-734.
 - 26 Alfonsi P, Slim K, Chauvin M, Mariani P, Faucheron JL, Fletcher D. Working Group of Société française d'anesthésie et réanimation (SFAR); Société française de chirurgie digestive (SFCD). French guidelines for enhanced recovery after elective colorectal surgery. *J Visc Surg* 2014;151:65-79. doi:10.1016/j.jvisc Surg.2013.10.006
 - 27 Brady M, Kinn S, Stuart P. Preoperative fasting for adults to prevent perioperative complications. *Cochrane Database Syst Rev* 2003;4:CD004423. doi:10.1002/14651858.CD004423
 - 28 Bilku DK, Dennison AR, Hall TC, Metcalfe MS, Garcea G. Role of preoperative carbohydrate loading: a systematic review. *Ann R Coll Surg Engl* 2014;96:15-22. doi: 10.1308/003588414X13824511650614.
 - 29 Ljungqvist O, Nygren J, Thorell A. Modulation of postoperative insulin resistance by preoperative carbohydrate loading. *Proc Nutr Soc* 2002;61:329-336. DOI:10.1079/ PNS2002168.
 - 30 Can MF, Yagci G, Dag B, Ozturk E, Gorgulu S, Simsek A, Tufan T. Preoperative administration of oral carbohydrate-rich solution: Comparison of glucometabolic responses and tolerability between patients with and without insulin resistance. *Nutrition.* 2009;25:72-7. doi: 10.1016/j.nut.2008.07.021.

- 31 Hausel J, Nygren J, Lagerkranser M, Hellström PM, Hammarqvist F, Almström C, Lindh A, Thorell A, Ljungqvist O. A carbohydrate-rich drink reduces preoperative discomfort in elective surgery patients. *Anesth Analg*. 2001;93:1344-50.
- 32 Ljungqvist O, Nygren J, Hausel J, Thorell A. Preoperative nutrition therapy – novel developments. *Scand J Nutr* 2000;44: 3-7.
- 33 Noble SE, Watson DS, Huong H, Davison B, Hainsworth PJ, Horgan AF. Pre-operative oral carbohydrate loading in colorectal surgery: a randomized controlled trial. *Colorectal Dis*. 2006;8:563-9. doi:10.1111/j.1463-1318.2006.00965.x
- 34 Dock-Nascimento DB, Aguilar-Nascimento JE, Linetzky Waitzberg D. Ingestion of glutamine and maltodextrin two hours preoperatively improves insulin sensitivity after surgery surgery: a randomized, double blind, controlled trial. *Rev Col Bras Cir*. 2012;39:449-55.
- 35 Yuill KA, Richardson RA, Davidson HI, Garden OJ, Parks RW. The administration of an oral carbohydrate-containing fluid prior to major elective upper-gastrointestinal surgery preserves skeletal muscle mass postoperatively-a randomised clinical trial. *Clin Nutr*. 2005;24:32-7. doi:10.1016/j.clnu.2004.06.009
- 36 Mathur S, Plank LD, McCall JL, Shapkov P, McIlroy K, Gillanders LK, Merrie AE, Torrie JJ, Pugh F, Koea JB, Bissett IP, Parry BR. Randomized controlled trial of preoperative oral carbohydrate treatment in major abdominal surgery. *Br J Surg*.2010;97:485-94. doi: 10.1002/bjs.7026.
- 37 Järvelä K, Maaranen P, Sisto T. Pre-operative oral carbohydrate treatment before coronary artery bypass surgery. *Acta Anaesthesiol Scand*. 2008;52:793-7. doi: 10.1111/j.1399-6576.2008.01660.x.

Table 1. Clinical characteristics

Characteristics	Group A (n=300)	Group B (n=300)	Group C (n=300)	Group D (n=300)	<i>p</i> value
Gender					0.369
Male	138	146	134	154	
Female	162	154	166	146	
Age					0.474
≤28 days	96	100	84	88	
28 days–1 year	204	200	216	212	
Disease					0.338
Hirschsprung's disease	148	162	157	152	
Sacrococcygeal teratoma	30	16	28	20	
Biliary atresia	34	30	35	44	
Congenital anal atresia	88	92	80	84	




Figure 1. Blood glucose was shown 2 h before anesthesia, at the time of anesthesia, 2 h post-operatively, and 12 h post-operatively. Blood glucose was significantly higher in groups B, C, and D than group A at the time of anesthesia. (* $p < 0.05$ groups B, C, and D vs group A groups).




Figure 2. The crying ratio was significantly higher in group A than groups B, C, and D.




Figure 3. The length of hospital stay was not significantly different between the groups.