

This author's PDF version corresponds to the article as it appeared upon acceptance. Fully formatted PDF versions will be made available soon.

Auscultation-assisted bedside postpyloric placement of feeding tube in critically ill patients: a prospective, observational study

doi: 10.6133/apjcn.201905/PP.0008

Published online: May 2019

Running title: Bedside postpyloric placement of feeding tube

Jianguo Xiao MD¹, Zhi Mao MD¹, Ming Hua MD^{1,2}, Tengfei Chen MD^{1,3}, Hui Liu MD¹, Pan Hu MM¹, Sheng Tang MD¹, Hongjun Kang MD¹, Feihu Zhou MD, PhD¹

¹Department of Critical Care Medicine, The General Hospital of PLA, Beijing, China

²Department of Emergency, 149 Clinical Department of the Eighty-Second Hospital of PLA, Lianyungang, China

³Department of Critical Care Medicine, Beijing Hospital of Traditional Chinese Medicine (affiliated to Capital Medical University), Beijing, China

⁴National Clinical Research Center for Kidney Diseases, The General Hospital of PLA, Beijing, China

Authors' email addresses and contributions:

Jianguo Xiao: xiaojg301@126.com; Zhi Mao: maozhi@126.com; Ming Hua: seshua@139.com; Tengfei Chen: 824053093@qq.com; Hui Liu: weichu1977@sohu.com; Pan Hu: hupan0215@163.com; Sheng Tang: ts00307@163.com; Hongjun Kang: doctorklbd@126.com; Feihu Zhou: feihuzhou301@126.com

The roles of the authors in this study were as follows: FH-Z, JG-X and Z-M conceived the idea. JG-X, Z-M, M-H, and TF-C wrote the manuscript. S-T coordinated the study. M-H, TF-C, P-H collected data. JG-X and M-H analyzed the data. Z-M and P-H performed statistical analysis. HJ-K and H-L supervised the whole study process. FH-Z and HJ-K assisted with study design and revised the manuscript. JG-X and Z-M contributed equally to this work. All authors read and approved the final manuscript. JG-X and Z-M contributed equally to this work

Corresponding Author: Dr Feihu Zhou, Department of Critical Care Medicine and National Clinical Research Center for Kidney Diseases, Chinese People's Liberation Army General Hospital, 28 Fu-Xing Road, Beijing 100853, People's Republic of China. Tel: 86-10-66938148. Fax: 86-10-88219862. Email: feihuzhou301@126.com; xiaojg301@126.com

ABSTRACT

Background and Objectives: To assess the efficacy and safety of auscultation-assisted bedside postpyloric feeding tube (ABPFT) placement in early enteral nutritional support for critically ill patients. **Methods and Study Design:** A prospective observational study was conducted and 92 critically ill patients who met the inclusion criteria undergoing ABPFT placement after the intravenous injection of 10 mg of metoclopramide were included. Abdominal X-ray was performed to confirm the location of the catheter tip. End points investigated were the success rate of tube placement, rate of jejunal tube placement, duration of the procedure, length of insertion, and number of attempts. Operational-related adverse events or complications were also documented and evaluated. **Results:** The total success rate of postpyloric feeding tube implantation was 97.8% (90/92), among which, 89.1% (82/92) of the tubes were placed proximal to the jejunum. The first-attempt success rate was 91.3% (84/92) and the mean attempt per individual patient was 1.11 ± 0.38 times. The mean operation time was 28.6 ± 17.7 minutes, and the mean insertion length of tube was 106.3 ± 9.6 cm. A total of 27 adverse events occurred in 19.6% (18/92) patients and there was no serious adverse events or complications during the study period. **Conclusions:** Assistance by auscultation can significantly improve the success rate of nasal feeding tube placement. This simple, safe and fast approach is feasible for the application among health practitioners in the intensive care unit.

Key Words: blind bedside, postpyloric placement, auscultation, nasojejunal tube, enteral nutrition

INTRODUCTION

Enteral nutritional support is one of the crucial treatment modalities for critically ill patients, which can significantly improve the prognosis of these patients compared with parenteral nutrition.¹ Jejunal nasogastric tube feeding, as a well-established treatment operation, has become an essential nutritional approach in patients with diseases such as severe acute pancreatitis, esophageal fistula, and severe gastroparesis. In addition, considering the risk factors for enteral feeding intolerance often seen in critically ill patient such as advanced age, mechanical ventilation, sedation, muscle relaxation, supine position, large volume feeding, previous gastroesophageal reflux, unconsciousness, and high-level paraplegia,^{2,3} various practice guidelines recommend postpyloric feeding to be considered in these circumstances⁴⁻⁸

to achieve the target feeding rate as soon as possible, and reduce the risk of aspiration pneumonia.⁹⁻¹²

At present, X-ray and endoscopy are the most commonly used guiding approaches for the placement of jejunum feeding tubes, with a success rate of almost 100%.¹³⁻¹⁵ However, it often requires transferring the patients to the radiology department or endoscopic center, which would not only increase preparation time for the procedure and workload of healthcare staff, but is also limited by the disease severity of critically ill patients and their tolerance to transport.¹⁶ Therefore, bedside approaches for jejunum nasogastric feeding tube placement are gaining momentum recently, such as electromagnetic navigation and ultrasound guidance.¹⁷⁻¹⁹ However, the shortcomings of these techniques are their uncertain success rate and dependency on the device and equipment. Recently, many have reported a bedside blind placement of jejunum tube but with various success rate and unsatisfactory reliability.²⁰⁻²⁵ The latest study found that only 5.5% of ICU patients had a transnasal feeding tube,²⁶ indicating a lack of a universal and safe method for the bedside placement of transnasal jejunal tube. The purpose of this prospective study was to evaluate the safety and efficacy of auscultation-assisted bedside postpyloric feeding tube (ABPFT) placement.

MATERIALS AND METHODS

Study design

This study is a single-center and prospective observational study approved by the ethics committee of the General Hospital of PLA (No. S2017-054-01) and registered at <http://www.chiicr.org.cn> (ChiCTR-OOC-17012108).²⁷ The patients were fully informed of the content of the study prior to the start of the study and informed consent forms were signed by the patient or the patient's next of kin (authorized family member). The study design is consistent with requirements of an observational study in clinical epidemiology.²⁸

Patients

Critically ill patients admitted in the ICU who had enteral nutrition support since July 24, 2017 were enrolled into this study if they had one of the following comorbidities or risk factors for gastric feeding intolerance: 1) severe acute pancreatitis; 2) mechanical ventilation ≥ 48 hours; 3) history of gastroesophageal reflux disease; 4) impairment of consciousness (Glasgow score ≤ 8); 5) high-level paraplegia; 6) need for continuous use of sedatives or muscle relaxants; 7) intolerance to gastric feeding; 8) esophageal fistula. Patients were excluded if they met one of the following criteria: 1) the presence of gastrointestinal bleeding,

obstruction or lower gastrointestinal fistula and enteral nutrition was not possible; 2) nasopharyngeal tumors, bleeding, surgery and other conditions making nasal cannulation impossible; 3) contraindications for metoclopramide; 4) history of esophageal varices confirmed by endoscopy or computed tomography (CT) scan; 4) progressively deteriorating disease without effective control.

Materials and medications

A total of two types of nasogastric tubes were used in this study: a 10-F convolute nasogastric tube (Flocare Bengmark, CH10-145 cm, NUTRICIA, The Netherlands) and a Frey Trelumina feeding tube (CH16/9-150 cm, FRESSENIUS KABI, Germany). Other equipment used were 20ml syringes, stethoscopes, measuring rulers, paraffin oil, and metoclopramide (injection).

Technique of auscultation-assisted bedside postpyloric placement of feeding tube.

Different from other bedside blind jejunal feeding tube placement methods, this study introduces an auscultation-assisted approach that is characterized by intermittent injection of a small amount of gas (5mL/time) to the lumen while simultaneously checking the catheter tip position via auscultation by stethoscopes. The specific auscultation sites and their corresponding anatomic locations were shown in Figure 1.

Ethics approval and consent to participate

All subjects provided informed consent. This study was approved by the medical ethics committee of the General Hospital of PLA (No. S2017-054-01) and registered at <http://www.chiicr.org.cn> (ChiCTR-OOC-17012108)

Preparation

There is no special requirement for patient position, although the supine position is most commonly adopted with head elevation of 30 to 45 degrees. A slow intravenous injection of metoclopramide (10mg) was given 10 minutes prior to the procedure. Sedative agents may also be given when indicated. One of the two types of feeding tubes was randomly selected, and the tube and guidewire were fully lubricated with paraffin oil. The length from the earlobe through the tip of the nose to the subxyphoid (U) and to the umbilicus (D) were measured (recorded as L1 and L2, respectively), as well as the length from D to R (recorded as L3).

The procedure

The procedure was performed by intensive care physicians or nurses who have undergone standardized training previously, and all of them had previous experience of blind insertion of jejunum tube by other bedside methods.

The feeding tube with guide wire was inserted through the nasal cavity, and then carefully forwarded with the same approach as nasogastric tube placement. After the feeding tube is confirmed to be in the stomach (approximately the length of L1) by Whoosh test²⁹ or aspiration of gastric fluid, the tube was further advanced to the length L2. A sufficient suction was then done with a syringe to maximally empty the gas and fluid within the stomach, before rapid gas injection of 5mL with simultaneous auscultation with a stethoscope at point M to feel the strength of the sound.

Subsequently, the jejunal tube was slowly inserted with a gentle force and the resistance was felt by the operator. When inserted at the length of (L2 + L3), auscultation was performed at the five auscultation points (U, D, L, R and M). The 5 mL of gas was injected with the same rate and complete suction afterwards. If the point with the strongest sound was located at R or between R and D, the operator should continue to advance the feeding tube slowly and repeat auscultation every 3-5 cm until the strongest sound reached point R while the sound at point M was abruptly turned down, indicating that the feeding tube has passed through the pylorus into the duodenal bulb or its descending part. If the strongest sound was located at U or L and the insertion resistance significantly reduced, the tube should be retracted to the length of L1 and the above procedure should be repeated.

When the feeding tube is advanced beyond the pylorus, the operator should continue advancing the tube slowly while repeating auscultation every 5-10 cm to identify whether the strongest part of sound is consistent with the shape of the duodenum consistent until the inserted length reaches 100-110cm (calculated based on patient height or empirically [L2 + 40] cm). The feeding tube is then fixed to the alar area with tape followed by a bedside X-ray. If the tip of the feeding tube has reached the Treitz ligament, the guidewire could be removed slowly and then enteral feeding could also be given. If X-ray result is indeterminant about whether the feeding tube has passed the pylorus, the operator should retract the feeding tube back to L2 and repeat the above operation. The detail of procedure steps is shown in Figure 2.

Outcomes and endpoints

Clinical parameters prior to intubation were collected, including baseline characteristics, disease diagnosis, presence or absence of mechanical ventilation, use of vasopressors or

sedatives, and results of disease severity assessment such as acute physiology and chronic health evaluation (APACHE) II, sequential organ failure assessment (SOFA), and nutrition risk in the critically ill (NUTRIC) scores.

The intubation process was performed under supervision. The variables documented were: the duration of the procedure, the length of intubation, the number of attempts (number of X-ray examinations), the adverse events from intubation or metoclopramide. Meanwhile, the changes in vital signs during intubation, including heart rate (HR), respiratory rate (RR), mean arterial pressure (MAP), pulse oximetry (SpO₂), and electrocardiographic were also documented.

The primary endpoint was the success rate of the feeding tube insertion into the pylorus, defined as tube reaching or exceeding the first part of the duodenum, as judged by bedside radiographs by intensive care physicians and experienced radiologists. Secondary endpoints included the rate at which the tube was placed further to the Treitz ligament, duration of the procedure, length of insertion, and number of attempts. The failure of the procedure was defined as three unsuccessful attempts to place the feeding tube into the pylorus and other approaches must be used.

Safety endpoints included catheter-related adverse events or vital signs events, the latter is defined as over 20% change in preoperative HR, RR or MAP, or SpO₂ below 90% during the procedure.

Statistical analysis

All statistical analyses were conducted using SPSS21.0 software (IBM Corp., Armonk, NY, USA). Quantitative variables were expressed as median or mean \pm standard deviation, and qualitative variables were expressed as the number of cases or percentages.

RESULTS

From July 2017 to March 2018, a total of 683 critically ill patients were admitted to the department, and 92 of them were eventually enrolled in the study according to the inclusive and exclusive criteria. Among which, 81 (88%) had APACHE II scores >8 , 63 (68.5%) needed mechanical ventilation, 28 (30.4%) required chronic renal replacement therapy (CRRT), and 15 (16.3%) required vasoactive drugs for blood pressure maintenance. The baseline characteristics of the included patients were presented in Table 1.

Endpoints

Ninety (97.8%) out of the 92 patients had successful placements of transnasal jejunal feeding tube distal to the pylorus with assistance of auscultation, and 82 of which had feeding tube placed proximal to the jejunum (Table 2). The first-attempt success rate in tube placement was 91.3% (84/92), with an average of 1.11 ± 0.38 attempts per patient, a mean operation time of 28.6 ± 17.7 minutes, and a mean catheter length of 106.3 ± 9.6 cm. Two patients who had failed tube placement both had history of radical resection of esophageal cancer. The feeding tube was then successfully placed under the guidance of a bedside upper endoscopy on the next day, and 1 patient was found with severe gastroptosis under endoscopy.

Complications and adverse events

A total of 27 adverse events occurred in 19.6% (18/92) of included patients, including vital sign fluctuations, nausea, mucosal injury, and pain due to intolerance to intubation that required additional sedation and analgesic agents (Table 3). The vital sign fluctuation presented mainly with heart rate and respiration rate changes that would typically returned to the baseline level after the procedure. No significant changes of blood pressure and SpO₂, nor serious adverse events such as aspiration or misplacement into the trachea or drug reactions to metoclopramide were found.

DISCUSSION

The most challenging part of transnasal jejunal feeding tube placement is ensuring the tube to pass through the pylorus. Some of the traditional assistive methods, including pushing a tube with external force through the pylorus under direct vision (endoscope), or with indirect assistive approaches (such as chest X-ray and electromagnetic navigation) to determine the location of the tip of the tube, can be used with a relatively high success rate. This indicates that if assisted with direct or indirect monitoring or guidance, jejunal feeding tube can become a relatively easy procedure for clinicians. However, the above approaches have substantial dependency on equipment or exposure to radiation damage, which would pose considerable risks for critically ill patients.

Therefore, many recent studies are investigating the role of portable equipment (such as ultrasound and electrocardiogram devices) on the assistance of feeding tube placement^{18,19} or blind bedside transnasal jejunal tube,²⁰⁻²⁵ aiming at minimizing the equipment dependence of this procedure and facilitate the popularization of the technique. However, the success rates of the above methods were not satisfactory. A recent study investigating blind bedside

placement in 127 critically ill patients reported a success rate of 81.9% for postpyloric feeding tube placement.³⁰ Therefore, the accuracy and reliability of these instruments, as well as the pH test and vacuum test, are still not sufficient to satisfactorily guide the tube placement procedure.

In this study, transnasal jejunum tube was placed with the bedside approach assisted by abdominal auscultation. The success rate of postpylorus tube placement (97.8%, 90/92), the first-attempt success rate (91.3%), and the mean number of attempts per patient (1.11 ± 0.38) were all significantly more favorable compared with previous studies, indicating that the auscultation-assisted technique can accurately determine the position of the tip of the feeding tube and serve as a reliable "eye" for the blind procedure, which can significantly improve the success rate. Two patients with failed intubation both had history of radical esophageal cancer resection and one of them had severe gastropnoxis, which made the accurate localization through auscultation almost impossible. Therefore, this approach may not be suitable for patients with anatomical changes of the upper gastrointestinal tract after surgery.

Another reason for the high success rate of this study may be the adequate decompression of the gastrointestinal tract prior to the procedure and prompt aspiration of the gastric air during the procedure, so that the volume of the gastric cavity could be minimized, and the tube would not easily be reverted in the stomach. These facilitated a smooth passage of the tube through the pylorus. In addition, when the tube wall is too soft, it may not be able to overcome the friction of the stomach wall and would be easily reverted. The two types of feeding tube used in this study are more tenacious and can better transduce force, which might have contributed to the higher success rate in this study.

Multiple previous studies reported that a blind transnasal tube placement is safe,^{20,21,30-32} but some also found that the blind jejunal tube placement could lead to more serious complications such as perforation of the alimentary tract and misplacement into the trachea.^{33,34} The mean APACHE II score in patients included in this study was 16.5 ± 8.0 (up to 37). In addition, the proportion of patients requiring mechanical ventilation, CRRT, or vasoactive drugs was 68.5%, 30.4%, and 16.3%, respectively. Although the overall risk was high, all patients successfully underwent the operation. The overall incidence of adverse events was 27 in 18 patients (19.6%) consisting mainly of minor life events such as nausea, mucous membrane bleeding, tachycardia and shortness of breath that quickly alleviated after the procedure. This indicates that it is safe to avoid forced advance of the tube via the auscultation-assisted approach, which can be carried out in a wider range of critically ill patients.

In the previous study by Lv et al,³⁰ the insertion length of the feeding tube was 95.6 ± 9.3 cm, and in 33.9% of the patients the tube was managed to reach the proximal jejunum. In comparison, the mean insertion length of tube in our study was 106.3 ± 9.6 and 89.1% of the patients the tube was delivered distal to the Treitz ligament, and the incidence of adverse events was relatively low. Our findings indicate that the deeper placement of jejunal feeding tube is safe.

Early enteral nutrition in critically ill patients may be associated with shorter ICU length of stay and better prognosis.⁶ Many studies suggest that jejunal nutrition can reduce feeding intolerance regurgitation more effectively and thus reduce the incidence of aspiration.^{2,3,5,7} Therefore, ABPFT can provide a reliable way for the implementation of early enteral nutrition, and significantly reduce operating time and cost compared with other methods such as navigations by endoscopy, fluoroscopy or magnet.

All operators of this study were both doctors and nurses, and they all had previous experience of blind nasojejunal intubation. It should also be taken into account that the success rate of this procedure may also depend on the operator's experience and technique. Therefore, for beginners, basic skills training may be needed. And whether this method can still maintain a high success rate may need to be further validated by future studies with multicentral nature and larger patient volume.

Conclusion

Auscultation-assisted bedside postpyloric feeding tube placement is a simple, safe and lowcost method that allows feasible bedside procedure at any time without the need for special additional equipment. It is potentially a more effective way of early enteral nutritional support for critically ill patients. The method is easy to learn and master, and is ready for application and popularization among ICU medical personnel.

CONFLICT OF INTEREST AND FUNDING DISCLOSURE

The authors declare no conflict of interest.

The study was financially supported by The Open Project of National Clinical Research Center for Geriatric Diseases, NCRCG-PLAGH-2017008.

REFERENCES

1. Joint Standards Task Force of A.S.P.E.N and the American Dietetic Association Dietitians in Nutrition Support Dietetic Practice Group, Russell M, Stieber M, Brantley S, Freeman AM, Lefton J, Malone

- AM, Roberts S, Skates J, Young LS. American Society for Parenteral and Enteral Nutrition and American Dietetic Association: standards of practice and standards of professional performance for registered dietitians (generalist, specialty, and advanced) in nutrition support. *J Am Diet Assoc.* 2007;107:1815-22. doi:10.1016/j.jada.2007.08.032.
2. Sajid MS, Harper A, Hussain Q, Forni L, Singh KK. An integrated systematic review and meta-analysis of published randomized controlled trials evaluating nasogastric against postpyloris (nasoduodenal and nasojejunal) feeding in critically ill patients admitted in intensive care unit. *Eur J Clin Nutr.* 2014;68:424-32. doi:10.1038/ejcn.2014.6.
 3. Schlein K. Gastric versus small bowel feeding in critically ill adults. *Nutr Clin Pract.* 2016;31:514-22. doi:10.1177/0884533616629633.
 4. Taylor BE, McClave SA, Martindale RG, Warren MM, Johnson DR, Braunschweig C et al. Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *Crit Care Med.* 2016;44:390-438. doi:10.1097/CCM.0000000000001525.
 5. Kreymann KG, Berger MM, Deutz NE, Hiesmayr M, Jolliet P, Kazandjiev G et al. ESPEN guidelines on enteral nutrition: intensive care. *Clin Nutr.* 2006;25:210-23. doi: 10.1016/j.clnu.2006.01.021.
 6. Reintam Blaser A, Starkopf J, Alhazzani W, Berger MM, Casaer MP, Deane AM, et al. Early enteral nutrition in critically ill patients: ESICM clinical practice guidelines. *Intensive Care Med.* 2017;43:380-98. doi:10.1007/s00134-016-4665-0.
 7. Critical Care Nutrition. Strategies to optimize delivery and minimize risks of EN: small bowel feeding vs gastric. In: Canadian Clinical Practice Guidelines. 2015/05 [cited 2015/11/13]; Available from: <http://criticalcarenutrition.com/docs/CPGs%202015/5.3%202015.pdf>.
 8. Academy of Nutrition and Dietetics, Evidence Analysis Library. Critical illness guideline. 2012. [cited 2014/11/15]; Available from: <https://www.andeal.org/topic.cfm?menu=5302&cat=4800>.
 9. Acosta-Escribano J, Fernández-Vivas M, Grau Carmona T, Caturla-Such J, Garcia-Martinez M, Menendez-Mainer A, Solera-Suarez M, Sanchez-Payá J. Gastric versus transpyloric feeding in severe traumatic brain injury: a prospective, randomized trial. *Intensive Care Med.* 2010;36:1532-39. doi:10.1007/s00134-010-1908-3.
 10. Metheny NA, Stewart BJ, McClave SA. Relationship between feeding tube site and respiratory outcomes. *JPEN J Parenter Enteral Nutr.* 2011;35:346-55. doi:10.1177/0148607110377096.
 11. Deane AM, Dhaliwal R, Day AG, Ridley EJ, Davies AR, Heyland DK. Comparisons between intragastric and small intestinal delivery of enteral nutrition in the critically ill: a systematic review and meta-analysis. *Crit Care.* 2013;17:R125. doi:10.1186/cc12800.
 12. Alhazzani W, Almasoud A, Jaeschke R, Lo BW, Sindi A, Altayyar S, Fox-Robichaud AE. Small bowel feeding and risk of pneumonia in adult critically ill patients: a systematic review and meta-analysis of randomized trials. *Crit Care.* 2013;17:R127. doi: 10.1186/cc12806.

13. Zhang L, Huang YH, Yao W, Chang H, Guo CJ, Lin SR. Transnasal esophagogastroduodenoscopy for placement of nasoenteric feeding tubes in patients with severe upper gastrointestinal diseases. *J Dig Dis.* 2012;13:310-15. doi: 10.1111/j.1751-2980.2012.00594.x.
14. Zhu Y, Yin H, Zhang R, Ye X, Wei J. Endoscopy versus fluoroscopy for the placement of postpyloric nasoenteric tubes in critically ill patients: a meta-analysis of randomized controlled trials. *J Crit Care.* 2016;33:207-12. doi: 10.1016/j.jcrc.2016.01.022.
15. Fang JC, Hilden K, Holubkov R, DiSario JA. Transnasal endoscopy vs fluoroscopy for the placement of nasoenteric feeding tubes in critically ill patients. *Gastrointest Endosc.* 2005;62:661-6. doi:10.1016/j.gie.2005.04.027.
16. Schwebel C, Clec'h C, Magne S, Minet C, Garrouste-Orgeas M, Bonadona A, et al. Safety of intrahospital transport in ventilated critically ill patients: a multicenter cohort study. *Crit Care Med.* 2013;41:1919-28. doi: 10.1097/CCM.0b013e31828a3bbd.
17. Powers J, Fischer MH, Ziemba-Davis M, Brown J, Phillips DM. Elimination of radiographic confirmation for small-bowel feeding tubes in critical care. *Am J Crit Care.* 2013;22:521-27. doi: 10.4037/ajcc2013755.
18. Mathus-Vliegen EMH, Duflou A, Spanier MBW, Fockens P. Nasoenteral feeding tube placement by nurses using an electromagnetic guidance system (with video). *Gastrointest Endosc.* 2010;71:728-36. doi: 10.1016/j.gie.2009.10.046.
19. Hernandez-Socorro CR, Marin J, Ruiz-Santana S, Santana L, Manzano JL. Bedside sonographic-guided versus blind nasoenteric feeding tube placement in critically ill patients. *Crit Care Med.* 1996;24:1690-4.
20. Gatt M, MacFie J. Bedside postpyloric feeding tube placement: a pilot series to validate this novel technique. *Crit Care Med.* 2009;37:523-7. doi: 10.1097/CCM.0b013e3181959836.
21. Kohata H, Okuda N, Nakataki E, Itagaki T, Onodera M, Imanaka H, Nishimura M. A novel method of post-pyloric feeding tube placement at bedside. *J Crit Care.* 2013;28:1039-41. doi: 10.1016/j.jcrc.2013.06.018.
22. Lai CW, Barlow R, Barnes M, Hawthorne AB. Bedside placement of nasojejunal tubes: a randomised-controlled trial of spiral-VS straightended tubes. *Clin Nutr.* 2003;22:267-70.
23. Duggan S, Egan SM, Smyth ND, Feehan SM, Breslin N, Conlon KC. Blind bedside insertion of small bowel feeding tubes. *Ir J Med Sci.* 2009;178:485-9. doi: 10.1007/s11845-009-0351-3.
24. Rollins CM. Blind bedside placement of postpyloric feeding tubes by registered dietitians: success rates, outcomes, and cost effectiveness. *Nutr Clin Pract.* 2013;28:506-9. doi: 10.1177/0884533613486932.
25. Lee AJ, Eve R, Bennett MJ. Evaluation of a technique for blind placement of post-pyloric feeding tubes in intensive care: application in patients with gastric ileus. *Intensive Care Med.* 2006;32:553-6. doi:10.1007/s00134-006-0095-8.

26. Bendavid I, Singer P, Theilla M, Themessl-Huber M, Sulz I, Mouhieddine M, Schuh C, Mora B, Hiesmayr M. NutritionDay ICU: a 7 year worldwide prevalence study of nutrition practice in intensive care. *Clin Nutr.* 2017;36:1122-9. doi: 10.1016/j.clnu.2016.07.012.
27. Chinese Clinical Trial Registry. Accessed 2017. Available from: <http://www.chictr.org.cn/showproj.aspx?proj=20676>.
28. Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, STROBE Initiative. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Int J Surg.* 2014;12:1495-9. doi: 10.1016/j.ijso.2014.07.013.
29. Dawson J. Nasogastric tube incidents and the use of the 'whoosh test'. *Crit Care.* 2007;11:419. doi: 10.1186/cc6083.
30. Lv B, Hu L, Chen L, Hu B, Zhang Y, Ye H, Sun C, Zhang X, Lan H, Chen C. Blind bedside postpyloric placement of spiral tube as rescue therapy in critically ill patients: a prospective, tricentric, observational study. *Crit Care.* 2017;21:248-55. doi: 10.1186/s13054-017-1839-2.
31. Rollins CM. Blind bedside placement of postpyloric feeding tubes by registered dietitians: success rates, outcomes, and cost effectiveness. *Nutr Clin Pract.* 2013;28:506-9. doi: 10.1177/0884533613486932.
32. Lee AJ, Eve R, Bennett MJ. Evaluation of a technique for blind placement of post-pyloric feeding tubes in intensive care: application in patients with gastric ileus. *Intensive Care Med.* 2006;32:553-6. doi:10.1007/s00134-006-0095-8.
33. Stayner JL, Bhatnagar A, McGinn AN, Fang JC. Feeding tube placement: errors and complications. *Nutr Clin Pract.* 2012;27:738-48. doi: 10.1177/0884533612462239.
34. van Kerckhoven G, Kant KM. Dedicated NAVA enteral feeding tube placement complicated by a hydropneumothorax but producing a correct EAdi signal. *Intensive Care Med.* 2012;42:271-2. doi: 10.1007/s00134-015-3981-0

Table 1. General characteristics and clinical data of patients enrolled[†]

Characteristics	Values (n=92)
Age (years)	53.1±13.2
Gender, male	69 (75.0)
Height (cm)	169.1±7.7
Weight (kg)	69.2±11.4
BMI (kg/m ²)	24.2±3.1
Primary diagnosis	
Severe acute pancreatitis	31 (33.7)
Gastrointestinal	22 (23.9)
Neurologic	17 (18.5)
Multiple trauma	8 (8.7)
Sepsis	6 (6.5)
Respiratory	4 (4.3)
Cardiovascular	2 (2.2)
Others	2 (2.2)
Preexisting diseases	
Hypertension	24 (26.1)
Previous gastrointestinal surgery	11 (12.0)
Coronary heart disease	9 (9.8)
Gastroesophageal reflux disease	5 (5.4)
APACHE II score	16.5±8.0
SOFA score	6.5±4.1
NUTRIC score	4.1±2.3
Mechanical ventilation	63 (68.5)
CRRT	28 (30.4)
Vasopressors	15 (16.3)
Sedatives & Analgesics	36 (39.1)

BMI: body mass index; APACHE II: acute physiology and chronic health evaluation II; SOFA: sequential organ failure assessment; NUTRIC: nutrition risk in the critically ill; CRRT: continuous renal replacement therapy.

[†]Data is presented as mean ± standard deviation or number (percentage).

Table 2. Primary and second endpoints of the procedures[†]

Endpoints	Values (n=92)
Primary endpoint	
Postpyloric placement	90 (97.8)
Secondary endpoints	
Placed at the proximal jejunum	82 (89.1)
Time to insertion (min)	28.6±17.7
Number of attempts	1.11±0.38
Success in first attempt	84 (91.3)
Length of insertion (cm)	106.3±9.6

[†]Data is presented as mean ± standard deviation or number (percentage).

Table 3. Major procedure-associated events during the insertion[†]

Procedure-associated events	Values (n=27)
Vital signs events [‡]	7 (25.9)
Increased sedation or analgesia during procedure	7 (25.9)
Nausea	6 (22.2)
Nasal or pharyngolaryngeal mucosa bleeding	3 (11.1)
Nasal or pharyngolaryngeal pain	2 (7.4)
Vomiting	1 (3.7)
Abdominal distention	1 (3.7)

[†]Data is presented as number (percentage).

[‡]Defined as vital sign fluctuate over the range of ±20% or pulse oxygen saturation drop to less than 90% during the insertion .




Figure 1. Auscultation sites and anatomic locations. U: the subxyphoid, corresponding to the gastric fundus; D: the umbilical cord, corresponding to the junction between the great curvature of the stomach and the antrum; M: the middle point between point U and point D, corresponding to the middle of the gastric cavity; R: the intersection point between the horizontal line through M and the midline of the right clavicle, corresponding to the bulbar and descending part of the duodenum; L: the intersection point between the horizontal line through M and the midline of the left clavicle, corresponding to the junction of stomach body and fundus on the greater curvature side.




Figure 2. Steps of auscultation-assisted bedside postpyloric feeding tube (ABPFT) placement. U: the subxyphoid; D: the umbilical cord; M: the middle point between point U and point D; L: the intersection point between the horizontal line through M and the midline of the left clavicle; L1: the length from the earlobe through the tip of the nose to U; L2: the length from the earlobe through the tip of the nose to D; L3: the length from D to R.