

Original Article

Increased coffee, tea, or other sugar-sweetened beverage consumption in adolescents is associated with less satisfactory dietary quality, body fatness and serum uric acid profiles over the past 18 years in Taiwan

Ya-Hui Shih MS¹, Hsin-Chuan Wu PhD², Wen-Harn Pan PhD^{1,3,4}, Hsing-Yi Chang DrPH^{1,3}

¹Institute of Population Health Sciences, National Health Research Institutes, Miaoli, Taiwan

²Department of Food Nutrition, College of Human Science and Technology, Chung Hwa University of Medical Technology, Tainan, Taiwan

³National Yang-Ming University, Taipei, Taiwan

⁴Institute of Biomedical Sciences of Academia Sinica, Taipei, Taiwan

Background and Objectives: Taiwan has a high density of convenience stores and beverage shops, which makes sugar-sweetened beverages (SSBs) very accessible to teenagers. This study examined the changes and the association between SSBs and biomarkers and nutrient intake, for teenagers over the course of 18 years using a national representative sample. **Methods and Study Design:** This cross-sectional study used data from the Nutrition and Health Survey in Taiwan (NAHSIT). Complete data for teenagers aged 13 to 19 years including a 24-hour dietary recall, anthropometric and clinical measurements, and SSBs from two periods were analyzed (1993-1996: N=1820; 2010-2011: N=2513). SAS callable SUDAAN was used for statistical analysis, adjusting for the sampling scheme. Log transformation was used for non-normal variables before linear models were used. Coffee or tea and SSB (excluding tea or coffee) consumers were categorized as non-drinkers, low (below), or high (above) consumers based on median intake during 1993-1996. **Results:** Intake of coffee or tea increased significantly in the 18 years of this study ($p<0.01$), whereas intake of SSBs (excluding coffee or tea) decreased significantly ($p<0.05$). Intake was significantly higher among second survey and those with high total energy intakes ($p<0.01$). For both coffee/tea and SSB, the high-intake groups had higher serum uric acid and intake of carbohydrates ($p<0.05$), lower intake of protein and phosphorus and lower dietary diversity score ($p<0.05$). **Conclusions:** Consumption of coffee or tea increased in adolescents during the 18 years. High intakes of SSB, coffee or tea was associated with high serum uric acid values and worse dietary quality.

Key Words: sugar-sweetened beverage, 24-hour dietary recall, serum uric acid, adolescents

INTRODUCTION

Energy intake from sugar-sweetened beverages (SSBs) among adolescents trended upwards between 1977 and 1996 in the United States,¹ but in both the United States and Australia this trend appeared to be declining in the last 10–20 years.²⁻⁴ Research performed in the United States has found that, although consumption of soft drinks has decreased, consumption of sweetened coffee, sweetened tea, sports drinks, and energy drinks has increased.^{2,4} Despite the decline in soft drink consumption, sugar from SSBs still remains the greatest source of sugar in the diets of adolescents, representing almost 20% of total sugar intake according to research conducted in Australia.⁵ The 2015 World Health Organization (WHO) guidelines strongly recommend that free sugar intake should represent only 10% of the total daily energy intake for children and adults. It also conditionally recommends that this intake should be reduced to 5% of the total daily energy intake, with particular attention focused on sugar intake

from SSBs.⁶ Excessive consumption of SSBs leads to increased risk of the metabolic syndrome,^{7,8} weight problems, obesity, gout, cardiovascular disease, and diabetes in children, adolescents, and adults.⁹ High SSB intake can also result in nutritional imbalances in adolescents.¹⁰⁻¹⁵

Taiwan has a very high concentration of convenience stores and beverage shops; at the end of 2017, there were a total of 10,662 convenience stores (including four major chains)^{16,17} and 21,346 beverage shops (including juice shops, beverage shops, coffee shops and alcoholic beverage

Corresponding Author: Dr Hsing-Yi Chang, Institute of Population Health Sciences, National Health Research Institutes, 35 Keyan Road, Zhunan Town, Miaoli County 350, Taiwan. Tel: +886-37-246166 ext 36333; Fax: +886-37-586261. Email: hsiingyi@nhri.org.tw

Manuscript received 31 January 2019. Initial review completed 11 February 2019. Revision accepted 26 February 2019. doi: 10.6133/apjcn.201906_28(2).0020

ages shop),¹⁸ or approximately one convenience store per 2211 persons^{16,17} and one beverage shop per 1104 persons.¹⁸ Beverages, excluding alcoholic beverages, were the greatest source of income (32.5%) to these convenience stores in 2015.¹⁶ Regulations in Taiwan prohibit the sale of SSBs in elementary and junior high schools, as well as the sale of soft drinks in senior high schools. Despite these provisions, the high density of convenience stores and beverage shops near schools means that SSBs remain very accessible for teenagers. Past research suggests that the closer teenagers are to a convenience store, the higher their chance of purchasing SSBs.¹⁹ In this 18 years, Taiwanese teenagers spent less on original foods and decreased energy intakes. Instead, they spent more money on processed food and ultra-processed food,²⁰ including SSB since the number of convenience stores and beverage shops has increased over time, and information about SSB intake was scarce, we examined the time trend of SSB intake using national representative samples. Since sweetened tea and coffee become very popular, especially in beverage stores, tea and coffee and other SSB are studied. We also explored the demographic factors associated with tea/coffee and other SSB intake, as well as the consequences of tea/coffee and other SSB intake on biomarkers. Figure 1 shows the concept diagram of this study.

METHODS

Study population

This study analyzed data from the 1993–1996 and 2010–2011 Nutrition and Health Survey in Taiwan (NAHSIT), and was approved by the Institute Review Board of the National Health Research Institutes. The study population for the 1993–1996 survey included those aged 3 years and older residing in Taiwan. The 1993–1996 survey incorporated a multi-stage stratified cluster sampling scheme to

select individuals for face-to-face interviews. The survey divided the whole of Taiwan into 7 strata according to the unique dietary patterns of the residents, the urbanization index, and geographical location. The four groups with unique dietary patterns were: (1) the Hakka people, who consume a lot of preserved food and lard; (2) 80% of residents of mountainous areas, who are aborigines; (3) people living on the eastern coast, isolated from the western areas; and (4) people living on the Peng-Hu islands, who consume a high amount of salt. After these four groups were identified, the remaining area was stratified by the degree of urbanization and geographic location into (5) metropolitan cities, (6) provincial class I townships, and (7) provincial class II townships. Three townships or districts were selected based on the selection probability proportional to its population sizes (PPS) in each of the strata, and each township or district was surveyed in the same season to account for seasonal variation. Each township had a designated number of individuals to be interviewed by age and gender and interviews were conducted in a household setting. The response rate was 74%. Details of the sampling design have been described elsewhere.²¹

The study population for the 2010–2011 survey included students at public and private high schools. A two-stage stratified cluster sampling was adopted for this survey. It first stratified the whole of Taiwan into 5 strata, including 2 northern regions, 1 central region, 1 southern region, and 1 eastern region. Four schools were selected from each stratum with PPS. Twenty students (10 male and 10 female) of each grade from each school were sampled randomly. The response rate was 87% for those in junior high school and 92% for those in senior high school. Further details of the survey sample and sampling methods have been provided elsewhere.²² All data collection methods were carried out in accordance with relevant




Figure 1. Conceptual diagram of study.

guidelines and regulations. All measurement protocols were approved by the Internal Review Board of National Health Research Institutes.

Informed consent was obtained after students were selected. Parental consent was also obtained for students younger than 18 years of age. These two surveys resulted in nationally representative samples. In the present study, we included junior and senior high school students aged 13 to 19 years, who could provide complete data for 24-hour dietary recall, anthropometric and clinical measurements, and SSB consumption. After excluding excessively high (≥ 37656 kJ) or low (< 836.8 kJ) total energy intake ($N=10$)²³, a total of 4333 participants were included in the analysis (1993–1996: $N=1820$; 2010–2011: $N=2513$). Ethical approval was not required in 1993–1996. Participants joined the survey if they agreed verbally. The 2010–2011 survey was approved by the National Health Research Institutes Research Ethics Committee.

Availability of data and material

The 1993–1996 Nutrition and Health Survey in Taiwan can be download directly at the following website for members: <https://srda.sinica.edu.tw/search/fsciiitem/1321>.

Dietary assessments

We used NAHSIT questionnaire data including 24-hour dietary recall, medical history, physical activity, and parental data. SSB and daily nutrient intake data were obtained from the 24-hour dietary recall information. For the purposes of this study, SSBs included sweetened coffee, sweetened tea, carbonated drinks (soft drinks), fruit drinks (including reconstituted juice, fermented fruit and vegetable juice, and other beverages), energy drinks (including sports drinks), and other drinks (cocoa or chocolate drinks, lactic drinks, iced drinks, and scented teas). SSBs did not include 100% juice or fruit wine. We noted that non-sweetened coffee or tea was an available option, although most teens tended to buy sweetened coffee or tea. For the purposes of this study, we therefore chose to consider coffee or tea separately from SSBs (excluding coffee or tea). Daily nutrient intake, were estimated based on the 24-hour dietary recall information and food composition tables from the website of the Food and Drug Administration, Ministry of Health and Welfare,²⁴ and included both macronutrients (carbohydrates, protein, total fats, saturated fatty acids, and cholesterol) and micronutrients (calcium, phosphorus, iron, sodium, niacin, vitamin B1, vitamin B2, vitamin C, and vitamin E). Individual intake was adjusted for total energy intake to determine % energy for macronutrients and mg/4184 kJ for micronutrients. In addition, based on other local research into SSBs,²⁵ we also examined the consumption of sweets, including all types of candy, chocolate, and cookies, but not bread or pastries. Adolescents who ate breakfast 7 days per week were considered regular breakfast eaters.

Other variables included age, sex, smoking status (non-smokers and ever-smokers, including former smokers, occasional smokers, and daily smokers), alcohol drinking status (non-drinkers and ever-drinkers, including anyone who had ever consumed alcohol), and physical activity level. Physical activity was measured in weekly metabolic equivalents (MET) based on the research by Ridley et

al.²⁶

Dietary quality

We used Dietary Diversity Score (DDS) to evaluate the dietary quality of adolescents. DDS has been widely used to assess the overall quality of food and nutrient intakes.^{27–29} Since there are different food groups in different countries, each country can develop their own DDS. This index used six major food groups that obtained from 24-hr recall. The six major food groups are grains, dairy, beans/fish/eggs/meat, vegetables, fruits, oil and nuts.^{28–30} We used two methods, one is giving 1 point for taking half portion of each food group daily (DDS1), the other is giving 1 point for taking 1 portion of each food group daily (DDS2). Both indices ranged 0 to 6 points.

Anthropometric and clinical measurements

NAHSIT data included a health examination component, which collected anthropometric data and biochemistry according to 8-hour fasting blood samples. Body mass index (BMI) was calculated as body weight in kilograms divided by the square of body height in meters. Waist circumference (WC) measurements were performed at the midpoint between the lower edge of the rib cage and the top of the iliac crest. Fasting blood was drawn during the morning, immediately after the individuals arrived for health check-ups. Blood samples were centrifuged and stored on dry ice, then delivered to a central laboratory and frozen at -70°C on the same day for further analysis. Blood profiles included high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), total cholesterol (TC), triglycerides (TG), fasting blood glucose, and serum uric acid.

Statistical analyses

SAS version 9.4 (SAS Institute Inc.) and SUDAAN version 11.0 were used to perform statistical analyses accounting for the unequal probability of sample selection. Each individual was weighted to reflect representativeness in the population at the time of the interview. Estimated daily intakes for coffee, tea, and SSB (excluding coffee or tea) in grams had a non-normal distribution and were categorized into three categories based on total intake during 1993–1996. These groups were: coffee or tea: non-intake (0 g), low intake (>0 g to < 402 g), and high intake (≥ 402 g); and SSB (excluding coffee or tea): non-intake (0 g), low intake (>0 g to < 379 g), and high intake (≥ 379 g). Low and high groups were determined based on relation to median of the 1993–1996 survey: low = lower than median, high = higher than median. Descriptive statistics were provided as percentages or means \pm standard deviations.

Comparisons between SSB intake groups and between survey dates were tested using the chi-squared test (categorical variables) or t test (continuous variables). Stacked bar charts were used to present patterns of SSBs in different surveys by sex. Because SSB intake was categorized into three groups, multinomial logistic regression was used to identify potential risk factors. Trend analysis was used to examine whether dependent variables increased or decreased in different SSB groups. Multiple regression was used to analyze the association between SSB and

anthropometry, biomarkers, and nutrients after adjusting for survey date, age, sex, MET, and total energy. Because nutrient intake had been adjusted for total energy intake, analysis of the relationship between SSB and nutrients did not account for total energy intake. To ensure that the regression fulfilled the criteria of normality, residual analysis was performed, including histogram, box plot, Q-Q plot, and residual plots. Only the residuals of LDL-C, carbohydrate, fat, and SFA remained normal in the regression. All others were log-transformed after excluding outliers. All results of multiple regression included residuals that were normally distributed, with skewness close to 0 and the absolute kurtosis less than 2. Two-sided p values < 0.5 were considered significant.

RESULTS

Table 1 shows the demographic characteristics of the adolescents in the two surveys by gender. The average age for both genders was 15.1 years in 2010–2011, slightly lower than the average of 15.5 years in 1993–1996. The BMI and WC of both genders increased significantly over the course of 18 years ($p < 0.05$). The proportion of females in the ever-drink alcohol group increased from 12.9% to 20.7% in 18 years ($p < 0.05$). The proportion of total energy intake greater than 1993–1996 median (7464.3 kJ) increased from about 68% to about 82% in high-intake groups of boys and from about 29% to about 56% ($p < 0.001$) in girls over the course of 18 years. The consumption of sweets almost doubled for both girls and boys ($p < 0.001$).

Figure 2 shows the changes in SSB consumption over the course of 18 years according to beverage type. Average intake of SSBs was 371.2 g (SE=16.7) in 1993–1996, which increased to 413.4 g (SE=22.9) 2010–2012, alt-

hough this did not reach statistical significance ($p=0.126$; data not shown). The intake of coffee or tea increased from 206.3 g to 311 g ($p < 0.001$), and the intake of soda decreased from 76.4 g to 27.2 g ($p < 0.001$). Over the 18 years, coffee and tea became a significant type of SSB for adolescents; in 1993–1996, coffee and tea formed approximately 56% of the total SSB intake, while in 2010–2011 this increased to 75% (data not shown).

Table 2 shows the clinical outcome, dietary quality and nutrient consumption of the adolescents in the two surveys by gender. BMI, WC, and fasting glucose increased significantly in 18 years ($p < 0.05$), whereas uric acid decreased significantly ($p < 0.001$). TG also increased significantly in boys ($p < 0.05$). Both DDS1 and DDS2 increased significantly ($p < 0.01$), indicating improvement of dietary quality. Most nutrient intakes increased ($p < 0.05$), except carbohydrates and vitamin C decreased ($p < 0.01$). Intakes of calcium and vitamin B-2 decreased significantly in girls ($p < 0.01$).

Table 3 shows the characteristics of SSB consumers. Over 50% of teens drank coffee or tea in 2010–2011 (boys 56.3% and girls 49.4%), and consumption for both genders increased significantly from 1993–1996. The proportion of teenagers consuming SSBs (excluding coffee or tea), on the other hand, decreased. For low-intake consumers, intake of both coffee or tea and SSB (excluding coffee or tea) decreased significantly among boys ($p < 0.05$), although only SSB (excluding coffee or tea) intake decreased among girls ($p=0.015$).

Table 4 shows the results of multinomial logistic regression of factors associated with different patterns of SSB intake. In terms of high vs. non-intake of coffee or tea, high intake was significantly associated with the following survey groups: 2010–2011 survey date (OR=1.66,




Figure 2. Trends in consumption patterns of SSBs in two different surveys.

Table 1. Modifications of biochemical parameters

Variable [†]	Boys (N= 2158)					Girls (N=2175)				
	1993~1996		2010~2011		<i>p</i> value*	1993~1996		2010~2011		<i>p</i> value*
	N	Weighted %	N	Weighted %		N	Weighted %	N	Weighted %	
Sample size (n)	918		1240			902		1273		
Age (years) (mean±SE)	15.5±0.1		15.1±0.1		0.008	15.5±0.1		15.1±0.1		0.005
Age, group					0.001					0.002
Young (junior high school)	568	49.3	718	64.0		558	48.7	747	64.9	
Older (senior high school)	352	50.7	526	36.0		345	51.3	527	35.1	
BMI (kg/m ²) (mean±SE)	20±0.4		21.9±0.2		0.001	20.4±0.1		20.9±0.1		0.011
WC (cm) (mean±SE)	67.5±0.7		77.5±0.5		<0.001	63.7±0.3		75±0.4		<0.001
Smoking status (%)					0.023					0.400
Non-smokers	524	80.1	821	89.0		679	97.0	914	95.3	
Ever-smokers	125	19.9	139	11.0		24	3.0	60	4.7	
Alcohol drinking status (%)					0.089					0.026
Non-drinkers	564	73.9	760	80.6		667	87.1	765	79.3	
Ever-drinkers	224	26.1	201	19.4		118	12.9	208	20.7	
MET (kJ/ kg/ h), group (%) [‡]					0.283					0.125
Low (<127.6)	264	38.7	423	43.4		474	63.0	647	69.3	
High (≥127.6)	515	61.3	536	56.6		295	37.0	316	30.7	
Energy intake (%) [‡]					<0.001					<0.001
Low (<7464.3kJ)	361	31.9	215	17.8		634	70.9	549	43.7	
High (≥7464.3kJ)	557	68.0	1025	82.2		268	29.1	724	56.3	
Dietary behavior (%)										
Sweet intake (>0 g/day)	385	41.4	1070	85.5	<0.001	427	47.8	1126	86.2	<0.001
Breakfast (7 days/week)	638	74.3	865	71.0	0.276	590	68.9	779	62.4	0.144

BMI: body mass index; WC: waist circumference; SSB: sugar-sweetened beverage; MET: metabolic equivalent task per week.

[†]Missing value not shown.

[‡]Based on median from 1993-1996.

**p* value based on t test for continuous variables and *p* value based on chi-square test for categorical variables between the surveys.

Table 2. Modifications of biochemical parameters

Variable	Boys (mean±SE)			Girls (mean±SE)		
	1993~1996	2010~2011	<i>p</i> value*	1993~1996	2010~2011	<i>p</i> value*
Sample size (n)	918	1240		902	1273	
Anthropometry and biochemical data						
BMI, kg/m ²	20.0±0.4	21.9±0.2	0.001	20.4±0.1	20.9±0.1	0.011
WC, cm	67.5±0.7	77.5±0.5	<0.001	63.7±0.3	75.1±0.4	<0.001
HDL-C, mg/dL	55.6±1.5	52.2±0.5	0.056	59.6±1.8	58.4±0.5	0.538
LDL-C, mg/dL	86.2±2.0	88.7±0.9	0.280	94.4±2.1	92.1±1.3	0.374
TC, mg/dL	155±2.0	156±0.9	0.949	168±2.8	164±1.5	0.264
TG, mg/dL	66.7±2.0	73.0±1.7	0.038	70.0±2.4	69.3±1.0	0.789
Fasting blood glucose, mg/dL	80.8±0.9	96.1±0.4	<0.001	79.7±0.9	92.9±0.4	<0.001
Serum uric acid, mg/dL	7.4±0.1	6.7±0.1	<0.001	5.5±0.1	4.8±0.0	<0.001
Dietary quality (range 0-6) [†]						
DDS1	4.2±0.1	4.5±0.0	0.003	4.1±0.0	4.5±0.0	<0.001
DDS2	3.7±0.1	4.1±0.0	<0.001	3.7±0.1	4.0±0.1	0.001
Nutrient consumption						
Carbohydrates (% energy)	55.9±0.9	51.7±0.52	0.002	56.4±0.7	51.8±0.5	<0.001
Protein (% energy)	15.5±0.4	15.7±0.2	0.681	15.5±0.4	15.4±0.1	0.818
Total fat (% energy)	28.8±0.9	32.9±0.4	0.002	28.6±0.5	33.4±0.4	<0.001
SFA (% energy)	10±0.3	11.3±0.2	0.005	9.9±0.2	11.6±0.2	<0.001
Cholesterol (mg/4184 kJ)	180±5.1	187±3.5	0.249	183±11.4	199±6.1	0.238
Calcium (mg/4184 kJ)	220±10.4	214±4.5	0.625	273±11.5	229±5.1	0.006
Phosphorus (mg/4184 kJ)	508±13.8	553±5.5	0.013	543±13.8	558±5.0	0.381
Iron (mg/4184 kJ)	6.5±0.2	7.2±0.1	0.011	6.8±0.4	7.6±0.2	0.067
Sodium (g/4184 kJ)	1.5±0.0	1.9±0.0	<0.001	1.7±0.0	2.0±0.0	<0.001
Niacin (mg/4184 kJ)	8.1±0.3	9.1±0.1	0.009	8.0±0.5	9.0±0.2	0.083
Vitamin B-1 (mg/4184 kJ)	0.6±0.0	0.6±0.0	0.025	0.6±0.0	0.6±0.0	0.037
Vitamin B-2 (mg/4184 kJ)	0.6±0.0	0.6±0.0	0.156	0.7±0.0	0.6±0.0	0.004
Vitamin C (mg/4184 kJ)	64.4±3.1	48.1±1.6	0.001	80.5±5.8	58.2±1.7	0.004
Vitamin E (mg α-TE/4184 kJ)	3.1±0.1	3.6±0.1	<0.001	3.9±0.2	4.0±0.1	0.734

SSB: sugar-sweetened beverage; BMI: body mass index; WC: waist circumference; HDL-C: high-density lipoprotein cholesterol; LDL-C: low-density lipoprotein cholesterol; TC: total cholesterol; TG: triglyceride; DDS: dietary diversity score; SFA: Saturated fatty acids.

[†]DDS1: participant ate 0.5 serving per group per day; DDS2: participant ate 1 serving per group per day.

**p* value based on *t* test for continuous variables and *p* value based on chi-square test for categorical variables between the surveys

Table 3. Changes of SSB consumption of teens aged 13-19 years between two surveys by sex

Variable [†]	Boys (mean±SE)			Girls (mean±SE)		
	1993-1996	2010-2011	<i>p</i> value*	1993-1996	2010-2011	<i>p</i> value*
Sample size (n)	918	1240		902	1273	
Coffee or tea						
Intake (g/day)	232±10.8	365±21.4	<0.001	176±15.7	254±18.4	0.009
Intake by group (g/day) [‡]						
No intake (0 g)	0.0	0.0	--	0.0	0.0	--
Low (0< to <402 g)	301±12.2	263±6.1	0.018	281±19.2	248±9.7	0.149
High (≥402 g)	773±45.5	858±24.9	0.191	677±21.3	725±19.5	0.131
Distribution of SSB group [‡] (weighted %)			0.002			0.030
No intake (0 g)	57.8	43.7		62.7	50.6	
Low (0< to <402 g)	20.1	19.9		19.4	21.8	
High (≥402 g)	22.2	36.4		17.8	27.6	
SSB (excluding coffee or tea)						
Intake (g)	208±19.7	117±13.8	0.004	114±5.4	87.3±9.2	0.032
Intake group (g/day) [‡]						
No intake (0 g)	0.0	0.0	--	0.0	0.0	--
Low (0< to <379 g)	258±17.0	198±15.5	0.028	235.3±11.5	187±11.8	0.015
High (≥379 g)	754±36.5	704±23.1	0.275	644.9±48.8	694±37.5	0.441
Distribution of SSB group [‡] (weighted %)			0.022			0.084
No intake (0 g)	62.0	73.8		71.4	75.3	
Low (0< to <379 g)	15.9	13.4		17.2	16.6	
High (≥379 g)	22.2	12.8		11.4	8.1	

SSB: sugar-sweetened beverage.

[†]Missing value not shown.

[‡]Based on median from 1993-1996.

**p* value based on *t* test for continuous variables and *p* value based on chi-square test for categorical variables between the surveys.

p=0.012), senior high (OR=1.46, *p*=0.002), ever-smokers (OR=2.36, *p*=0.004), and high total intake (OR=1.84, *p*=0.009). Those with high MET were associated with the non-intake group (OR=0.76, *p*=0.046). A positive trend in appeared in the second survey among senior high students (vs. junior high students), ever-smokers, and total energy intake (*p* values for trend=0.035, 0.002, 0.011, and 0.014, respectively), indicating that the higher these variables, the higher the chance of being in the high coffee or tea intake group. Regarding SSB (excluding coffee or tea) high vs. non-intake, high total energy intake (OR=2.38, *p*<0.001) was associated with the high SSB group. Teens in the 2010-2011 survey were negatively associated with high SSB intake (OR=0.44, *p*=0.001). The positive trend appeared for boys and total energy intake (*p* values for trend=0.014 and <0.001, respectively) and a negative trend appeared in the second survey compared to the first survey (*p* values for trend=0.002).

The relationships between the different patterns of SSB intake and anthropometry, biomarkers, dietary quality and nutrients are presented in Table 5. For coffee or tea intake, after controlling for age, sex, MET, and total energy intake, BMI (β =0.02, *p*=0.041), WC (β =0.02, *p*=0.013) and serum uric acid (β =0.04, *p*=0.022) were significantly higher in the high intake group compared to the non-intake group. DDS were lower in high intake group than non-drinkers (β =-0.33 to -0.31, *p*<0.01). DDS1 in low intake group also significantly lower than the non-intake group. (β =-0.21, *p*=0.046). Because nutrients were already converted to nutrient density (per 4184 kJ), the variables controlled were survey date, age, sex, and MET. The results showed that the group with a high intake of coffee or tea had higher carbohydrate intake (β =2.62, *p*=0.003), lower protein intake (β =-0.09, *p*=0.003), lower

phosphorus intake (β =-0.06, *p*=0.041), and lower vitamin E (β =-0.09, *p*=0.008) intake than the group with no intake of coffee or tea.

The group with a high intake of SSBs (excluding coffee or tea) had a higher serum uric acid (β =0.05, *p*=0.012), a higher intake of carbohydrates (β =6.09, *p*<0.001), and vitamin C (β =0.49, *p*<0.001), but had a lower DDS (β =-0.31 to -0.24, *p*<0.05), a lower intake of protein (β =-0.13, *p*<0.001), total fat (β =-3.79, *p*<0.001), SFA (β =-0.90, *p*=0.027), calcium (β =-0.19, *p*=0.023), phosphorus (β =-0.12, *p*<0.001), and vitamin B1 (β =-0.12, *p*=0.010) than the group with no intake of SSBs (excluding coffee or tea).

DISCUSSION

This study compared adolescent SSB intake over 18 years using two surveys of national representative samples. We found that tea or coffee, as a variety of SSB, became a major source of sugar consumption for Taiwanese adolescents over the course of 18 years. Both high- and low-SSB intake were significantly associated with survey date and total energy intake. Consuming higher amounts of SSB or coffee or tea was associated with higher values of serum uric acid and higher intake of carbohydrates, as well as lower intake of protein and phosphorus and had a lower DDS.

We found that adolescents in Taiwan consumed more than 400 g of SSBs per day in 2010-2011, which is almost double the amount consumed by 10-17 years old in 13 other countries, which had average intakes of 271 mL/day for boys and 250 mL/day for girls. SSBs in this case included hot beverages (coffee, tea, and other hot beverages), juices, regular soft beverages (sugared and artificially sweetened, carbonated and non-carbonated

Table 4. The characteristics of participating chefs and cooks (n=90)

Variable	Low vs Non-intake [†]			High vs Non-intake [†]			<i>p</i> -trend*
	OR	95% CI	<i>p</i> value	OR	95% CI	<i>p</i> value	
Coffee or tea							
Survey year (ref. 1993-1996)							
2010-2011	1.22	0.85-1.73	0.247	1.66	1.18-2.92	0.012	0.035
Age group (ref. young)							
Older (senior high school)	0.96	0.72-1.28	0.739	1.46	1.19-1.79	0.002	0.002
Boy (Ref. girl)	0.97	0.69-1.37	0.871	0.89	0.63-1.24	0.434	0.647
Smoking status (ref. non-smokers)							
Ever-smokers	1.76	1.06-2.92	0.033	2.36	1.41-3.96	0.004	0.011
Alcohol drinking status (ref. non-drinkers)							
Ever-drinkers	0.82	0.41-1.63	0.534	1.16	0.89-1.51	0.243	0.481
MET, group (ref. low)							
High (≥ 127.6 kJ/kg/h) [‡]	0.88	0.64-1.20	0.386	0.76	0.58-0.99	0.046	0.124
Energy intake, group (ref. low)							
High (≥ 7464.3 kJ) [‡]	1.27	0.68-2.35	0.417	1.84	1.21-2.80	0.009	0.014
Dietary behavior							
Sweet intake (>0 g/day) (ref. no sweet intake)	1.02	0.61-1.69	0.946	1.27	0.78-2.06	0.294	0.351
Breakfast (7 days/week) (ref. <7 days/week)	1.02	0.66-1.58	0.920	0.78	0.48-1.27	0.286	0.530
SSB (excluding coffee or tea)							
Survey year (ref. 1993-1996)							
2010-2011	0.78	0.61-1.01	0.061	0.44	0.30-0.64	0.001	0.002
Age group (ref. young)							
Older (senior high school)	1.14	0.63-2.04	0.635	1.32	0.82-2.12	0.224	0.455
Boy (Ref. girl)	0.84	0.51-1.39	0.466	1.48	1.00-2.18	0.050	0.014
Smoking status (ref. non-smokers)							
Ever-smokers	0.84	0.49-1.44	0.498	0.88	0.37-2.14	0.762	0.750
Alcohol drinking status (ref. non-drinker)							
Ever-drinker	1.06	0.71-1.59	0.740	1.00	0.51-1.97	1.000	0.883
MET, group (ref. low)							
High (≥ 127.6 kJ/kg/h) [‡]	1.18	0.79-1.78	0.374	0.89	0.46-1.72	0.711	0.511
Energy intake (%), group (ref. low)							
High (≥ 7464.3 kJ) [‡]	1.42	1.00-2.01	0.052	2.38	1.73-3.27	<0.001	<0.001
Dietary behavior from FFQ							
Sweet intake (>0 g/day) (ref. no sweet intake)	1.02	0.83-1.24	0.868	1.43	0.87-2.34	0.138	0.314
Breakfast (7 days/week) (ref. <7 days/week)	1.16	0.69-1.93	0.540	0.69	0.44-1.07	0.089	0.157

MET: metabolic equivalent task per week; SSB: sugar-sweetened beverage.

[†]Intake group in coffee or tea (g): Non-intake, 0; Low, >0 to <402; High, ≥ 402 . Intake group in SSB (excluding coffee or tea) (g): Non-intake, 0; Low, >0 to <379; High, ≥ 379 . The values are based on the median from 1993-1996.

[‡]Based on median from 1993-1996.

*Tests for linear trend.

soft drinks, energy drinks, sports drinks, and other sugared or artificially sweetened soft drinks), as well as other beverages. Thirty-one Eastern Asia in general has the lowest SSB intake (0.2 servings/day which was about 45 g/day) among 187 countries (including sodas, fruit drinks, sports/energy drinks, pre-sweetened iced tea, and homemade SSB), according to a 2010 survey. In contrast to this, Taiwanese men consumed 1.02 servings/day and women consumed 0.94 servings/day, equivalent to 231 g/day to 213 g/day.³² This suggests that Taiwanese consumption of SSB was high regardless age. Some countries have recently shown declines in SSB consumption. A 2005–2008 Australian survey found SSB consumption decreased among 4 to 18-year-olds over the course of 2–3 years.³ In the United States, the number of adolescents consuming SSBs decreased from 87% in 1999–2000 to 77% in 2007–2008, and the energy taken from SSBs also decreased from 363 kcal/day to 286 kcal/day (1518.8 kJ/day to 1196.6 kJ/day).² In addition to standard SSBs,

coffee and tea consumption was high among Taiwanese adolescents; the proportion drinking coffee and tea already exceeded 50% in 1993–1996 and this increased to 70% in 2010–2011. One possible reason for this increase might be the increased availability of tea and coffee at beverage shops, which sell various sweetened teas, such as black or green tea, and mixed tea, such as pearl milk tea, jelly milk tea, and fruit tea. Along with these teas, some shops also sell coffee, all at a price affordable to teens; these beverage shops rarely sell soda drinks, limiting the availability of that type of SSB.

Many other studies have observed that SSB consumption increases with age, with adolescents consuming more than younger children.^{5,10,11,33,34} This agrees with our findings that senior high school students drank more coffee or tea than those in junior high. This may possibly be a factor of the older students' greater disposable income and longer commute to school, which increases both their opportunity and ability to eat in restaurants and purchase

Table 5. Regression analysis of clinical outcomes, DDS and daily nutrient intake associated with different pattern of SSB consumption

Variables [†]	Coffee or tea						SSB (excluding coffee or tea)					
	Low*		<i>p</i> value	High*		<i>p</i> value	Low*		<i>p</i> value	High*		<i>p</i> value
	Beta	95% CI		Beta	95% CI		Beta	95% CI		Beta	95% CI	
Anthropometry and biochemical data [‡]												
Log-BMI, kg/m ²	0.01	-0.03 to 0.04	0.706	0.02	0.00 to 0.04	0.041	-0.01	-0.03 to 0.02	0.509	-0.01	-0.03 to 0.01	0.178
Log-WC, cm	0.00	-0.01 to 0.02	0.621	0.02	0.00 to 0.03	0.013	-0.01	-0.02 to 0.01	0.210	-0.01	-0.02 to 0.00	0.114
Log-HDL-C, mg/dL	0.03	-0.05 to 0.10	0.460	-0.01	-0.07 to 0.06	0.827	-0.02	-0.07 to 0.03	0.337	-0.03	-0.08 to 0.02	0.250
LDL-C, mg/dL	1.56	-3.46 to 6.58	0.504	2.20	-1.66 to 6.06	0.233	4.10	-0.12 to 8.32	0.055	4.87	-1.07 to 10.80	0.098
Log-TC, mg/dL	0.01	-0.03 to 0.05	0.456	0.01	-0.01 to 0.04	0.272	0.02	-0.01 to 0.05	0.267	0.02	-0.02 to 0.06	0.337
Log-TG, mg/dL	-0.02	-0.09 to 0.05	0.597	0.01	-0.05 to 0.07	0.779	0.01	-0.08 to 0.10	0.730	-0.00	-0.12 to 0.12	0.995
Log-Fasting blood glucose, mg/dL	0.01	-0.02 to 0.04	0.511	0.01	-0.02 to 0.04	0.512	0.01	-0.01 to 0.03	0.415	0.00	-0.01 to 0.01	0.772
Log-Serum uric acid, mg/dL	0.01	-0.03 to 0.04	0.727	0.04	0.01 to 0.08	0.022	-0.01	-0.04 to 0.03	0.734	0.05	0.01 to 0.09	0.012
Dietary quality (range 0-6) ^{‡§}												
DDS1	-0.21	-0.43 to -0.00	0.046	-0.33	-0.46 to -0.21	<0.001	0.03	-0.15 to 0.22	0.695	-0.24	-0.42 to -0.06	0.014
DDS2	-0.26	-0.52 to 0.00	0.053	-0.31	-0.50 to -0.13	0.004	0.02	-0.17 to 0.20	0.825	-0.31	-0.49 to -0.13	0.003
Nutrient consumption [¶]												
Carbohydrates (% energy)	0.40	-2.85 to 3.65	0.791	2.62	1.11 to 4.13	0.003	-0.11	-3.32 to 3.09	0.938	6.09	4.59 to 7.59	<0.001
Log-Protein (% energy)	-0.03	-0.07 to 0.02	0.195	-0.09	-0.14 to -0.04	0.003	0.00	-0.04 to 0.04	0.884	-0.13	-0.17 to -0.08	<0.001
Total fat (% energy)	-0.08	-3.29 to 3.12	0.955	-1.23	-2.98 to 0.52	0.149	0.10	-3.26 to 3.47	0.947	-3.79	-5.31 to -2.28	<0.001
SFA (% energy)	0.23	-0.98 to 1.45	0.677	0.05	-0.86 to 0.96	0.908	0.44	-0.60 to 1.47	0.371	-0.90	-1.66 to -0.13	0.027
Log-Cholesterol (mg/4184 kJ)	-0.04	-0.16 to 0.09	0.511	-0.04	-0.13 to 0.05	0.375	-0.09	-0.25 to 0.07	0.227	-0.11	-0.27 to 0.04	0.133
Log-Calcium (mg/4184 kJ)	0.04	-0.09 to 0.17	0.518	-0.07	-0.19 to 0.05	0.210	-0.03	-0.19 to 0.12	0.645	-0.19	-0.36 to -0.03	0.023
Log-Phosphorus (mg/4184 kJ)	0.02	-0.03 to 0.06	0.463	-0.06	-0.02 to -0.00	0.041	0.01	-0.04 to 0.06	0.536	-0.12	-0.17 to -0.08	<0.001
Log-Iron (mg/4184 kJ)	0.04	-0.07 to 0.14	0.426	0.07	-0.01 to 0.15	0.093	-0.00	-0.06 to 0.06	0.884	-0.03	-0.10 to 0.04	0.300
Log-Sodium (g/4184 kJ)	-0.10	-0.21 to 0.01	0.064	-0.08	-0.18 to 0.03	0.130	0.01	-0.14 to 0.15	0.903	-0.03	-0.13 to 0.06	0.449
Log-Niacin (mg/4184 kJ)	0.04	-0.02 to 0.09	0.152	0.02	-0.05 to 0.10	0.489	-0.00	-0.09 to 0.08	0.985	0.02	-0.09 to 0.13	0.670
Log-Vitamin B-1 (mg/4184 kJ)	-0.06	-0.15 to 0.03	0.170	-0.07	-0.16 to 0.03	0.138	0.01	-0.08 to 0.10	0.818	-0.12	-0.20 to -0.04	0.010
Log-Vitamin B-2 (mg/4184 kJ)	0.04	-0.04 to 0.13	0.279	0.05	-0.06 to 0.15	0.346	0.01	-0.06 to 0.08	0.752	-0.01	-0.07 to 0.05	0.740
Log-Vitamin C (mg/4184 kJ)	0.06	-0.06 to 0.18	0.260	0.01	-0.17 to 0.20	0.895	0.14	-0.05 to 0.34	0.125	0.49	0.28 to 0.70	<0.001
Log-Vitamin E (mg α -TE/4184 kJ)	-0.07	-0.15 to 0.02	0.124	-0.09	-0.14 to -0.03	0.008	-0.09	-0.25 to 0.07	0.226	-0.11	-0.27 to 0.04	0.120

SSB: sugar-sweetened beverage; BMI: body mass index; WC: waist circumference; HDL-C: high-density lipoprotein cholesterol; LDL-C: low-density lipoprotein cholesterol; TC: total cholesterol; TG: triglyceride; DDS: dietary diversity score; SFA: Saturated fatty acids.

[†]The log transformation is used for most variables except for LDL-C, carbohydrates, total fat and SFA.

[‡]The multiple regression model was adjusted for survey year, age, gender, physical activity performed per week (MET), daily energy intake (in kilojoule).

[§]DDS1: participant ate 0.5 serving per food group per day; DDS2: participant ate 1 serving per food group per day.

[¶]The multiple regression model was adjusted for survey date, age, gender and physical activity performed (MET) per week.

*Intake group in coffee or tea (g): Non-intake, 0; Low, >0 to <402; High, \geq 402. Intake group in SSB (excluding coffee or tea) (g): Non-intake, 0; Low, >0 to <379; High, \geq 379.

SSBs. Two surveys of 14–18 year old in the US found that smokers had higher intake of SSBs, which accords with our findings that students in the ever-smoker group had high intakes of SSBs.^{35,36} As has been observed in males in many other studies, high total energy intake was associated with high SSB intake.^{31,32,37–39}

Adolescents consuming high amounts of SSB or coffee or tea had high serum uric acid values. High consumption of coffee or tea was also associated with higher BMI and WC. Fructose might contribute to this effect. Usually, sucrose or high-fructose corn syrup (HFCS) are added to SSBs; the main contents of these two sweeteners are fructose and glucose. Fructokinase has no negative feedback preventing it from continuing to phosphorylate substrate; as a result, fructose turns into acetyl-CoA and glyceraldehyde, which then accumulate and increase *de novo* lipogenesis. Eventually, TG are formed. In fructokinase, a large amount of adenosine triphosphate (ATP) is consumed and generates uric acid.⁴⁰ The effects of SSB consumption on increasing uric acid values have been observed in a 1999–2004 survey in United States⁴¹ and Taiwan.⁴² Uric acid is an independent risk factor for all-cause mortality.^{43,44}

The average DDS2 was around 3.7, indicating relative good dietary quality.²⁹ Steyn et al has shown that when using 6 food groups to calculate DDS, those scored 3 had over 70% micronutrients intakes. That was sufficient. But that study was on children under 9 years old. There has not been appropriate cut-off for adolescents. Nevertheless, we found that higher intakes of SSB would reduce DDS. In other words, SSB intakes reduced dietary quality.^{28,29} Only three nutrients were associated with SSB intakes. It was possible due to intakes of ultra-processed food.²⁰ Past study has shown that high school students have increased more ultra-processed food intakes, and resulted in decrease in many micronutrients.

We found that intakes of carbohydrates increased as both coffee or tea and SSB (excluding coffee or tea) intake increased, and intakes of protein and phosphorus decreased as both pattern of SSB increased. The more SSBs consumed, the higher the sugar (carbohydrate) intake.^{11,14,15,45} Furthermore, consuming large amounts of SSBs was associated with the intake of other nutrients.^{10–15,45} We found that high consumption of SSBs (excluding coffee or tea) was associated with high vitamin C intake. The increase in vitamin C intake may be due to consumption of fruit drinks or to beverage shops adding fruits such as plums to beverages. A 1994–1996 survey in the United States found that adolescents consuming fruit-flavored drinks had a better chance of fulfilling their daily recommended vitamin C intake, but drinking soda did not help.¹² Canada had similar findings in their 2004 survey.⁴⁶ Our study included soft drinks, energy drinks, and fruit drinks; we believe the vitamin C may have come from fruit drinks.

The strength of this study is that it used national representative samples to examine the change in adolescent SSB intake over the course of 18 years. In addition, 24-hour dietary recall was conducted during a face-to-face interview. These surveys were performed periodically to reflect social changes in a timely manner. However, there were some limitations to this study. The cross-sectional

nature of the design could not infer a causal relationship. There were some differences in measurement and units of certain nutrients; therefore, some nutrients, such as vitamin A, vitamin D, magnesium, potassium, and zinc, were not included in the analysis. Intake information according to 24-hour recall was used; normally, this might not reflect the usual intake of SSBs. However, we obtained repeated measurements on a subsample (N=325) of participants in 2010–2011. The correlation on total SSBs was 0.48 (0.42 for boys and 0.57 for girls). All reached statistical significance, and the high correlation implied that they consumed SSBs regularly. Furthermore, correlation coefficients could be used to estimate the variance between individuals in dietary data with repeated measurements.⁴⁷ The variance between individuals can be estimated as $S_b = S\sqrt{\gamma}$, where S is the total variance. When variance between individuals is reduced, the chance of reaching statistical significance increases. This calculation is, however, beyond the scope of this study. Based on the high correlation of our repeated subsample measurements, we believe our data reflects the usual intake of SSBs in teens.

Conclusions

Coffee or tea has become a major source of SSBs in adolescents in Taiwan. Over 50% of adolescents consumed coffee or tea in 2010–2011. High intake of SSBs (including coffee or tea) was associated with higher uric acid. This suggests that adolescent consumption of SSBs is not trivial and is a habit requiring great effort to change. To promote healthier teen choices regarding SSBs, public health education should take greater care in educating parents on providing a healthy diet with low sugar intake. Schools should teach students how to choose beverages intelligently, encouraging children and teens to choose non-sweetened drinks or water instead of SSBs.

ACKNOWLEDGEMENTS

Data analyzed came from the “Nutrition and Health Survey in Taiwan (NAHSIT)” funded by the Department of Health (DOH FN8202, DOH-83-FS-41, DOH-84-FS-11, DOH-85-FS-11, DOH-86-FS-11) in 1993–1996 and by the Food and Drug Administration, Department of Health, Executive Yuan (99TFDA-FS-408, 100TFDA-FS-406) in 2010–2011. In 1993–1996 the project was carried out by the Institute of Biomedical Sciences, Academia Sinica and the Department of Biochemistry, College of Medicine, National Taiwan University under the direction of Dr Wen-Harn Pan and Dr Po-Chao Huang. The Office of Nutrition Survey, Center for Survey Research, Academia Sinica was responsible for data distribution. In 2010–2011 the project was carried out by the Division of Preventive Medicine and Health Services Research, Institute of Population Health Sciences, National Health Research Institutes (NHRI) under the direction of Dr Wen-Harn Pan. The Office of Nutrition Survey, Division of Preventive Medicine and Health Services Research, Institute of Population Health Sciences, National Health Research Institutes (NHRI) was responsible for data distribution. The assistance provided by the institutes and aforementioned individuals is greatly appreciated. The views expressed herein are solely those of the authors.

AUTHOR DISCLOSURES

The authors declare that there are no competing interests.

Part of this study was supported by the National Health Re-

search Institutes (grant number PH-103-SP-12) and the Ministry of Science and Technology (grant number MOST 105-3011-F-400-001).

REFERENCES

- Nielsen SJ, Siega-Riz AM, Popkin BM. Trends in food locations and sources among adolescents and young adults. *Prev Med.* 2002;35:107-13. doi: 10.1006/pmed.2002.1037.
- Han E, Powell LM. Consumption patterns of sugar-sweetened beverages in the United States. *J Acad Nutr Diet.* 2013;113:43-53. doi: 10.1016/j.jand.2012.09.016.
- Jensen BW, Nichols M, Allender S, de Silva-Sanigorski A, Millar L, Kremer P, Lacy K, Swinburn B. Consumption patterns of sweet drinks in a population of Australian children and adolescents (2003-2008). *BMC Public Health.* 2012;12:771. doi: 10.1186/1471-2458-12-771.
- Kit BK, Fakhouri TH, Park S, Nielsen SJ, Ogden CL. Trends in sugar-sweetened beverage consumption among youth and adults in the United States: 1999-2010. *Am J Clin Nutr.* 2013;98:180-8. doi: 10.3945/ajcn.112.057943.
- Louie JC, Moshtaghan H, Rangan AM, Flood VM, Gill TP. Intake and sources of added sugars among Australian children and adolescents. *Eur J Nutr.* 2016;55:2347-55. doi: 10.1007/s00394-015-1041-8.
- World Health Organization. *Guideline: Sugars intake for adults and children.* Geneva, Switzerland: World Health Organization; 2015.
- Chan TF, Lin WT, Huang HL, Lee CY, Wu PW, Chiu YW et al. Consumption of sugar-sweetened beverages is associated with components of the metabolic syndrome in adolescents. *Nutrients.* 2014;6:2088-103. doi: 10.3390/nu6052088.
- Mirmiran P, Yuzbashian E, Asghari G, Hosseinpour-Niazi S, Azizi F. Consumption of sugar sweetened beverage is associated with incidence of metabolic syndrome in Tehranian children and adolescents. *Nutr Metab (Lond).* 2015;12:25. doi: 10.1186/s12986-015-0021-6.
- Malik VS, Hu FB. Fructose and cardiometabolic health: what the evidence from sugar-sweetened beverages tells us. *J Am Coll Cardiol.* 2015;66:1615-24. doi: 10.1016/j.jacc.2015.08.025.
- Fiorito LM, Marini M, Mitchell DC, Smiciklas-Wright H, Birch LL. Girls' early sweetened carbonated beverage intake predicts different patterns of beverage and nutrient intake across childhood and adolescence. *J Am Diet Assoc.* 2010;110:543-50. doi: 10.1016/j.jada.2009.12.027.
- Libuda L, Alexy U, Buyken AE, Sichert-Hellert W, Stehle P, Kersting M. Consumption of sugar-sweetened beverages and its association with nutrient intakes and diet quality in German children and adolescents. *Br J Nutr.* 2009;101:1549-57. doi: 10.1017/S0007114508094671.
- Ballev C, Kuester S, Gillespie C. Beverage choices affect adequacy of children's nutrient intakes. *Arch Pediatr Adolesc Med.* 2000;154:1148-52. doi: 10.1001/archpedi.154.11.1148.
- Lyons J, Walton J, Flynn A. Food portion sizes and dietary quality in Irish children and adolescents. *Public Health Nutr.* 2015;18:1444-52. doi: 10.1017/S1368980014001979.
- Bowman SA. Beverage choices of young females: changes and impact on nutrient intakes. *J Am Diet Assoc.* 2002;102:1234-9. doi: 10.1016/S0002-8223(02)90273-7.
- Kant AK, Graubard BI. Contributors of water intake in US children and adolescents: associations with dietary and meal characteristics: National Health and Nutrition Examination Survey 2005-2006. *Am J Clin Nutr.* 2010;92:887-96. doi: 10.3945/ajcn.2010.29708.
- Ministry of Economic Affairs. *Industrial Production Indexes in April 2018 2016/6/27* [cited: 2019/03/06]; Available from: https://www.moea.gov.tw/Mns/dos/bulletin/Bulletin.aspx?kind=9&html=1&menu_id=18808&bull_id=5108.
- DeAeth D. Taiwan second highest in world for ratio of convenience stores per population 2018/4/3 [cited: 2019/03/06]; Available from: <https://www.taiwannews.com.tw/en/news/3397167>.
- Ministry of Finance. *Monthly statistics of finance: January 2018.* 2018/10/01 [cited: 2019/03/06]; Available from: <http://web02.mof.gov.tw/njswww/WebProxy.aspx?sys=100&funid=defjspf2>.
- Hearst MO, Pasch KE, Laska MN. Urban v. suburban perceptions of the neighbourhood food environment as correlates of adolescent food purchasing. *Public Health Nutr.* 2012;15:299-306. doi: 10.1017/S1368980011002114.
- Chen YC, Huang YC, Lo YTC, Wu HJ, Wahlqvist ML, Lee MS. Secular trend towards ultra-processed food consumption and expenditure compromises dietary quality among Taiwanese adolescents. *Food Nutr Res.* 2018;62. doi: 10.29219/fnr.v62.1565.
- Chang HY, Pan WH, Yeh WT, Tsai KS. Hyperuricemia and gout in Taiwan: results from the Nutritional and Health Survey in Taiwan (1993-96). *J Rheumatol.* 2001;28:1640-6. doi: 10.1007/s00394-015-1041-8.
- Tu SH, Hung YT, Chang HY, Hang CM, Shaw NS, Lin W et al. Nutrition and Health Survey of Taiwan Elementary School Children 2001-2002: research design, methods and scope. *Asia Pac J Clin Nutr.* 2007;16(Suppl 2):507-17. doi: 10.6133/apjcn.2007.16.s2.02.
- Ferrari P, Slimani N, Ciampi A, Trichopoulou A, Naska A, Lauria C et al. Evaluation of under- and overreporting of energy intake in the 24-hour diet recalls in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Public Health Nutr.* 2002;5:1329-45. doi: 10.1079/PHN2002409.
- Food composition tables. Food and Drug Administration, Ministry of Health and Welfare; 2018. [cited date: 2019/03/06]; Available from: <https://consumer.fda.gov.tw/Food/TFND.aspx?nodeID=178>.
- Tseng AT, Tsao HC, Sarukura N, Huang KC, Wang MF, Huang PC, Hsu TF, Yamamoto S. Development of a food composition database of refined monosaccharides and disaccharides in snacks, desserts, and beverages. *Nutritional Sciences Journal.* 2010;35:11. doi: 10.6691/NSJ.
- Ridley K, Ainsworth BE, Olds TS. Development of a compendium of energy expenditures for youth. *Int J Behav Nutr Phys Act.* 2008;5:45. doi: 10.1186/1479-5868-5-45.
- Kant AK, Schatzkin A, Harris TB, Ziegler RG, Block G. Dietary diversity and subsequent mortality in the First National Health and Nutrition Examination Survey Epidemiologic Follow-up Study. *Am J Clin Nutr.* 1993;57:434-40. doi: 10.1093/ajcn/57.3.434.
- Mirmiran P, Azadbakht L, Esmailzadeh A, Azizi F. Dietary diversity score in adolescents - a good indicator of the nutritional adequacy of diets: Tehran lipid and glucose study. *Asia Pac J Clin Nutr.* 2004;13:56-60.
- Steyn NP, Nel J, Labadarios D, Maunder EMW, Kruger HSJN. Which dietary diversity indicator is best to assess micronutrient adequacy in children 1 to 9 y? *Nutrition.* 2014;30:55-60. doi: 10.1016/j.nut.2013.06.002.
- Lee MS, Huang YC, Wahlqvist ML. Chewing ability in conjunction with food intake and energy status in later life affects survival in Taiwanese with the metabolic syndrome. *J Am Geriatr Soc.* 2010;58:1072-80. doi: 10.1111/j.1532-5415.2010.02870.x.

31. Guelinckx I, Iglesia I, Bottin JH, De Miguel-Etayo P, Gonzalez-Gil EM, Salas-Salvado J et al. Intake of water and beverages of children and adolescents in 13 countries. *Eur J Nutr.* 2015;54(Suppl 2):69-79. doi: 10.1007/s00394-015-0955-5.
32. Singh GM, Micha R, Khatibzadeh S, Shi P, Lim S, Andrews KG et al. Global, regional, and national consumption of sugar-sweetened beverages, fruit juices, and milk: a systematic assessment of beverage intake in 187 countries. *PLoS One.* 2015;10:e0124845. doi: 10.1371/journal.pone.0124845.
33. Lee HS, Kwon SO, Lee Y. Weight status and dietary factors associated with sugar-sweetened beverage intake among Korean children and adolescents - Korea National Health and Nutrition Examination Survey, 2008-2011. *Clin Nutr Res.* 2013;2:135-42. doi: 10.7762/cnr.2013.2.2.135.
34. Stern D, Piernas C, Barquera S, Rivera JA, Popkin BM. Caloric beverages were major sources of energy among children and adults in Mexico, 1999-2012. *J Nutr.* 2014;144:949-56. doi: 10.3945/jn.114.190652.
35. Terry-McElrath YM, O'Malley PM, Johnston LD. Energy drinks, soft drinks, and substance use among US secondary school students. *J Addict Med.* 2014;8:6. doi: 10.1097/01.ADM.0000435322.07020.53.
36. Park S, Blanck HM, Sherry B, Foti K. Problem behavior, victimization, and soda intake in high school students. *Am J Health Behav.* 2013;37:414-21. doi: 10.5993/AJHB.37.3.14.
37. Bleich SN, Wolfson JA. U.S. adults and child snacking patterns among sugar-sweetened beverage drinkers and non-drinkers. *Prev Med.* 2015;72:8-14. doi: 10.1016/j.ypmed.2015.01.003.
38. Yamada M, Murakami K, Sasaki S, Takahashi Y, Okubo H. Soft drink intake is associated with diet quality even among young Japanese women with low soft drink intake. *J Am Diet Assoc.* 2008;108:1997-2004. doi: 10.1016/j.jada.2008.09.033.
39. Mathias KC, Slining MM, Popkin BM. Foods and beverages associated with higher intake of sugar-sweetened beverages. *Am J Prev Med.* 2013;44:351-7. doi: 10.1016/j.amepre.2012.11.036.
40. Perheentupa J, Raivio K. Fructose-induced hyperuricaemia. *Lancet.* 1967;2:528-31. doi: 10.1016/S0140-6736(67)90494-1.
41. Nguyen S, Choi HK, Lustig RH, Hsu CY. Sugar-sweetened beverages, serum uric acid, and blood pressure in adolescents. *J Pediatr.* 2009;154:807-13. doi: 10.1016/j.jpeds.2009.01.015.
42. Lin WT, Huang HL, Huang MC, Chan TF, Ciou SY, Lee CY et al. Effects on uric acid, body mass index and blood pressure in adolescents of consuming beverages sweetened with high-fructose corn syrup. *Int J Obes (Lond).* 2013;37:532-9. doi: 10.1038/ijo.2012.121.
43. Chen JH, Chuang SY, Chen HJ, Yeh WT, Pan WH. Serum uric acid level as an independent risk factor for all-cause, cardiovascular, and ischemic stroke mortality: a Chinese cohort study. *Arthritis Rheum.* 2009;61:225-32. doi: 10.1002/art.24164.
44. Zhang W, Iso H, Murakami Y, Miura K, Nagai M, Sugiyama D, Ueshima H, Okamura T, Epoch-Japan G. Serum uric acid and mortality from cardiovascular disease: EPOCH-JAPAN study. *J Atheroscler Thromb.* 2016;23:1365-6. doi: 10.5551/jat.Er31591.
45. Vartanian LR, Schwartz MB, Brownell KD. Effects of soft drink consumption on nutrition and health: a systematic review and meta-analysis. *Am J Public Health.* 2007;97:667-75. doi: 10.2105/AJPH.2005.083782.
46. Danyliw AD, Vatanparast H, Nikpartow N, Whiting SJ. Beverage intake patterns of Canadian children and adolescents. *Public Health Nutr.* 2011;14:1961-9. doi: 10.1017/S1368980011001091.
47. Delcourt C, Cubeau J, Balkau B, Papoz L. Limitations of the correlation coefficient in the validation of diet assessment methods. *CODIAB-INSERM-ZENECA Pharma Study Group. Epidemiology.* 1994;5:518-24.