

This author's PDF version corresponds to the article as it appeared upon acceptance. Fully formatted PDF versions will be made available soon.

Effects of fructose from apple and honey on serum uric acid in young Chinese: randomized crossover trials

doi: 10.6133/apjcn.202112/PP.0003

Published online: December 2021

Running title: Serum uric acid and dietary fructose

Yinyin Cheng BSc^{1†}, Hui Zhang BSc^{1†}, Yong Zhu BSc⁴, Zhe Xue MD¹, Mengyao Yan MD¹, Hui Wang BSc², Shuben Sun MD³, Xiaohong Zhang PhD^{1,3}

¹Department of Preventive Medicine, Zhejiang Key Laboratory of Pathophysiology, School of Medicine, Ningbo University, Zhejiang, China

²Department of Colorectal Surgery, Shaoxing People's Hospital, Shaoxing, Zhejiang, China

³The Affiliated Hospital of Medical School, Ningbo University, Zhejiang, People's Republic of China

⁴Ningbo Academy of Agricultural Sciences, Ningbo, Zhejiang, China

[†]Both authors contributed equally to this manuscript

Authors' email addresses and contributions:

Yinyin Cheng: ChengYinYin411526@163.com

Contribution: carried out the trials, performed the data analysis and contributed to the writing of the manuscript

Hui Zhang: zhzhanghai2020@163.com

Contribution: performed the blood sample collection and data analysis and contributed to the writing of the manuscript

Yong Zhu: zjzsy@163.com

Contribution: detected the content of sugar in foods

Zhe Xue: xuezhe95@163.com

Contribution: collected the blood samples and obtained the anthropometric parameters

Mengyao Yan: yanmy922118@163.com

Contribution: collected the blood samples

Hui Wang: 269413012@qq.com

Contribution: collected the blood samples

Shuben Sun: nbhappyben@163.com

Contribution: collected blood samples

Xiaohong Zhang: zhangxiaohong1@nbu.edu.cn

Contribution: designed the study and revised the paper

Corresponding Author: Dr Xiaohong Zhang, Department of Preventative Medicine, Medicine School; Zhejiang Key Laboratory of Pathophysiology, School of Medicine, Ningbo University, 818 Fenghua Road, Ningbo, Zhejiang Province, 315211, China; The Affiliated Hospital of Medical School, Ningbo University, 247 Renmin Road, Ningbo, Zhejiang, 315020, People's Republic of China. Tel: +86-0574-87609591. Fax: +86 0574 87608638. Email: zhangxiaohong1@nbu.edu.cn

Not Proof Read

ABSTRACT

Background and Objectives: Overconsumption of drinks containing fructose increases the risk for hyperuricemia and gout. Comparative analysis evaluating the indicators of serum uric acid (SUA) load caused by natural food-derived fructose and pure fructose in sweeteners is lacking. We aimed to uncover the effect of fructose from apple and honey and pure fructose powder on the SUA concentration of healthy young Chinese individuals. **Methods and Study Design:** Two randomized crossover trials were performed. The participants were randomly assigned to consume apple or honey (test food) or pure fructose powder (reference food); one week later, the groups' dietary intervention was switched. Blood samples were collected at 0, 30, 60, and 120 min after meal to measure the SUA and blood glucose concentrations. **Results:** At 30 and 60 min, the SUA concentration in participants consuming apple or honey was lower than in those consuming fructose powder. At 120 min, the SUA concentration of participants consuming apple returned to baseline. The areas under the curve (AUC) within 2 h (2h-AUCs) of SUA exhibited the trend of fructose >honey >apple. The 2h-AUC ratio between test food and reference food was determined using the uric acid index to assess the efficiency of food-derived fructose in increasing the SUA concentration. The uric acid index of honey was higher than that of apple. Men had higher postprandial SUA concentration than women. **Conclusions:** Food-derived fructose caused a lighter load on uric acid metabolism than pure fructose reagent. Uric acid index can be useful for distinguishing fructose-containing foods.

Key Words: serum uric acid, uric acid index, fructose, apple, honey

INTRODUCTION

Uric acid in blood at low concentration exerts an anti-oxidative effect according to the human and animal data; however, in excessive amounts, it crystallizes and acts as a pro-oxidant, thereby inducing gout.¹ To date, chronic hyperuricemia is accepted as an independent risk factor for gout,² rheumatoid arthritis,³ nonalcoholic fatty liver disease, diabetes,⁴⁻⁶ and even some types of cancers.^{7,8}

Uric acid is not only derived from the breakdown of purines, but also from fructose metabolism. The hepatic metabolism of fructose leads to the rapid depletion of ATP and increased production of uric acid.⁹⁻¹² In the modern food industry, a sweetener rich in fructose is often used in processed foods, including sweetened foods and beverages, since fructose is the sweetest naturally occurring carbohydrate.¹³ High fructose corn syrup (HFCS) is a man-

made flavored syrup containing a high proportion of fructose and is widely used in manufactured foods. Participants consuming HFCS-sweetened beverage reportedly had higher serum uric acid (SUA) concentration than those consuming sucrose-sweetened beverage. Data from previous prospective cohort studies reported a positive association between the intake of fructose-rich beverages and an increased risk of gout or hyperuricemia in men and women.^{14,15} A previous meta-analysis showed that the risk of hyperuricemia is positively correlated with the intake of fructose (odds ratio: 1.85; 95% confidence interval: 1.66–2.07).¹⁶ Moreover, a larger body of evidence, including from animal experiments,^{17,18} epidemiological studies, and clinical trials, supports that sugary drinks with HFCS also increase the risk of obesity, hypertension, insulin resistance, fatty liver, and dyslipidemia. Therefore, limiting the intake of HFCS is largely recommended.¹⁹⁻²¹

Notably, fruits and honey are the main sources of natural fructose.²² The Brisighella Heart Study based on the Mediterranean cohort showed that people who consumed large amounts of fruits had elevated SUA concentration and increased incidence of hypertension and diabetes,²³ low-density lipoprotein oxidation,²⁴ arterial stiffness,²⁵ impaired cognitive function,²⁶ and heart failure.²⁷ The overconsumption of fructose even from fruits is suggested to be harmful. However, increasing the intake of fruits is recommended by health organizations worldwide as they are low energy-dense foods that are rich in fiber and micronutrients.²⁸ Therefore, a method for assessing the capacity of fruits to increase the SUA concentration is warranted. Honey is a sweet and viscous food substance. It is often used for cooking and baking, or as a spread on bread in Western countries; it is also used in some commercial beverages such as tea drink in China.

This study aimed to compare the effect of fructose from apple, honey, and pure fructose powder on the SUA concentration of healthy young Chinese individuals and explore a method to evaluate the load on uric acid metabolism caused by natural food-derived fructose.

MATERIALS AND METHODS

Study design

The study was performed at the School of Medicine, Ningbo University, from September to October 2020. Two randomized crossover trials, apple trial and honey trial, were carried out: apple or honey was used as test food, while a commercial pure fructose powder was used as a reference food. In the apple trial, the participants were randomly assigned to consume apple (test food) or pure fructose powder (reference food) at the first stage, and received an alternative dietary treatment at the second stage after a washout period of one week. In the

honey trial, the test food was honey, while the reference food was pure fructose powder; the participants received the dietary intervention following the same principle. If the sample size is 16 persons, an 80% power is required to detect a difference of 1 standard deviation (SD) in plasma uric acid response to treatment at an α level of 0.05, since 20% of the participants is expected to drop out from the study. Therefore, the sample size exceeds 20 persons.²⁹

Participants

College students were recruited from Ningbo University. Participants 1) aged 18–26 years, 2) with a fasting blood glucose concentration of <6.1 mmol/L, and 3) with a body mass index of <24 kg/m² according to the recommendation of the China Nutrition Society were included in this study. Participants with 1) menstruation and 2) chronic diseases, including diabetes, hypertension, cancers, digestive disorders, gout, and hyperuricemia, which was defined as a fasting SUA of >416.5 μ mol/L (men) or 357 μ mol/L (women) on two tests were excluded.³⁰ Written informed consent was obtained before the trials were conducted. The experiment was approved by the Ethics Committee of School of Medicine, Ningbo University (Ningbo, China), and registered in the Chinese Clinical Trial Registry at <http://www.chictr.org.cn/> (registration number: ChiCTR2000036443).

Trial foods

The same batch of red Fuji apples and Guanshengyuan honey produced by a local manufacturer were used as test foods. Fructose powder with a purity of $\geq 99\%$ was purchased from Shandong Xiwang Sugar Co. (Shandong, China) and was used as the reference food. At the beginning of trial, honey, fructose powder dissolved in 50 mL warm water, and fresh apple pulp were provided to the participants.

Measurement of fructose in foods using HPLC

HPLC reagents and conditions

The amount of fructose, glucose, and sucrose in foods was measured using high-performance liquid chromatography (HPLC) according to the China National Standard protocol GB 5009.8-2016. A Waters 2695 HPLC apparatus equipped with 2414 differential refractive index detector (Waters Corp, USA) was used. Glucose, fructose, and sucrose standard substances were purchased from Chem Service (West Chester, USA). Zinc acetate and potassium ferrocyanide of analytical grade were purchased from Sinopharm Chemical Reagent Co., Ltd (Shanghai, China). A 20 μ L sample was injected into a Kromasil 100A NH2

column at a temperature of 40 °C (250 × 4.6 mm, 5 µm; Kromasil) and detected using a refractive index detector at a temperature of 35 °C. The elution was performed using acetonitrile and water (75:25, v/v) at a flow rate of 1.0 mL/min.

Sample preparation

Three randomly chosen red Fuji apples were equally divided into three layers (inner, middle, and outer) after removal of peels and seeds. Equal amounts (5 g) of flesh from each layer of the three apples were mixed with the solution (2.5 mL of zinc acetate solution [Zn(CH₃COO)₂·2H₂O], 2.5 mL of potassium ferricyanide solution {K₄[Fe(CN)₆]·3H₂O}, and 45 mL of ultra-pure water) and shaken for 60 min. After centrifugation at 9,500 rpm for 5 min, the mixtures were filtered through a 0.22-µm nylon filter (Millipore). Honey (30 g) from three individual bottles was directly added to sterile tubes.

Calibration curve

A series of dilutions with concentrations of 0.4, 2.0, 5.0, 10.0, and 20.0 mg/mL were prepared and then detected by HPLC to establish the calibration curve of fructose, glucose, and sucrose. The calibration curve was plotted with the peak area on the vertical (Y) axis and the standard concentrations on the horizontal (X) axis using linear regression.

Study protocol

The participants were prohibited from drinking alcohol, consuming oversized animal meat, eating desserts, or staying up late the day before the test. All participants fasted from 10:00 pm the day before the test. On the actual test day, they arrived at the scene at 08:00 am. The height, body weight, and blood pressure of the participants were recorded. Then, the dietary treatment was administered orally. During the dietary intervention, all participants were asked to consume their food within 10 min. Three milliliters of venous blood samples were collected at 0, 30, 60, and 120 min in order to analyze the uric acid concentration. Blood pressure was also measured at 120 min.

Statistical analysis

All data were analyzed using SPSS 24.0 software (Chicago, IL, USA). The parameters obtained at 0 min of dietary intervention, uric acid index, and 2h-AUC of SUA are expressed as mean ± Standard deviation (SD), while other data were presented as mean ± standard error of means (SEM). The differences in the SUA and blood glucose concentrations between 0

min and 30 min or between 30 and 120 min after a specific treatment were analyzed using repeated measures of the general linear model, while the values obtained at 0 min were adjusted as covariates to avoid the interference of differences between individuals; the sequence of dietary intervention and participants as random effects were adjusted. The 2h-AUC of serum uric acid and blood glucose after administering the dietary treatment were calculated using the trapezoidal rule (GraphPad Prism 8.0.1, San Diego, CA, USA). The normality of variables such as 2h-AUC of uric acid, 2h-AUC of blood glucose, and the changes in blood pressure (both diastolic and systolic) from 0 min to 120 min was assessed using the Shapiro-Wilk test. Normalized and non-normalized data were analyzed using one-way analysis of variance and non-parametric test, respectively. Independent-samples t-test was used to determine the uric acid index. A two-tailed p value of <0.05 was considered statistically significant.

RESULTS

Quantification of fructose with HPLC

The result of HPLC showed that 1 g of Fuji apple provides 0.110 g of fructose (0.096 g of free fructose, 0.041 g of glucose, and 0.0311 g of sucrose comprising 0.014 g of glucose and 0.014 g of fructose), while 1 g of honey has 0.489 g of free fructose and 0.442 g of glucose. Considering the feasibility of intake, the dose of fructose consumed was 25 g, which is equivalent to 222 g of flesh from a medium-sized apple or 51 g of honey.

Anthropometric characteristics of the participants

A flow chart of the study is presented in Figure 1. Twenty-nine and thirty-five college students aged 22–26 years were enrolled in the apple trial ($n=29$) and honey trial ($n=35$), respectively. Prior to the administration of the dietary intervention, seven (men) and three (two men and one woman) volunteers dropped out of the apple trial and honey trial, respectively, due to unrelated personal reasons. Four participants (three men and one woman) in the apple trial and 11 participants (eight men and three women) in the honey trial were excluded owing to the occurrence of hyperuricemia. The anthropometric characteristics of participants at baseline are summarized in Table 1; no difference was found in all measured parameters at baseline among the participants.

SUA concentration of participants

The changes in SUA concentration of all participants consuming apple, honey, and fructose powder are depicted in Figures 2A and Table 2. As shown in Figures 2A, the SUA concentration of participants after consumption of apple rapidly peaked during the first 30 min, but then gradually decreased to the lowest at 120 min; meanwhile, the SUA concentration of participants consuming honey continuously increased. Of note, the curves of SUA over time after fructose powder and apple consumption had the same shape but different amplitude. After comparing SUA concentration at different time points, results showed that the fructose powder intake induced a significant increase in SUA concentration at 30 and 60 min compared with that at 0 min (fructose(apple trial): $p_{30\text{min}} < 0.001$, $p_{60\text{min}} < 0.001$; fructose(honey trial): $p_{30\text{min}} < 0.001$, $p_{60\text{min}} < 0.001$). However, no significant difference was found between the SUA concentrations at 30 and 60 min after consuming apple and honey and the SUA concentration at 0 min (apple: $p_{30\text{min}} = 0.075$, $p_{60\text{min}} = 0.148$; honey: $p_{30\text{min}} = 0.057$, $p_{60\text{min}} = 0.052$). In our study, the difference between SUA test food and SUA reference food (i.e., SUA test food – SUA reference food) of each participant at the same time points after meal was calculated to compare the body's response to fructose derived from different sources. As shown in Table 2, all the values of the SUA test food and SUA reference food at 30 min and 60 min for the apple trial and honey trial were negative. Results of the statistical analysis showed that the SUA concentration of women at 30 min and 60 min after consuming apple were significantly lower than those of fructose ($p_{30\text{min}} = 0.01$, $p_{60\text{min}} = 0.02$).

Moreover, men had a stronger response to fructose intake than women, with a higher concentration of SUA at the same time point compared with women, as illustrated in Figure 2B and Figure 2C. The difference between SUA men and SUA women (SUA men–SUA women) was used to quantitatively evaluate the effect of gender on uric acid metabolism. As shown in Table 3, the SUA concentrations of men at 60 min and 120 min were higher than that of women, and a significant difference in SUA was observed between women and men at 120 min after fructose intake in the honey trial ($p = 0.04$).

Then, the 2h-AUC of SUA after dietary fructose loading was measured. As shown in Table 4 and Figure 2D and 2E, male and female participants consuming apple and honey had lower 2h-AUC of SUA than those consuming fructose powder. The 2h-AUC of SUA of men consuming apple and honey was significantly higher than that of women (p value for apple intake: was 0.005, p value for honey intake: 0.015). In terms of SUA concentration, men exhibited a stronger response to fructose-containing foods than women. The gender-related

difference in fructose treatment almost disappeared when the participants consumed fructose powder.

In this study, a new metabolic parameter was introduced, uric acid index, which was defined as the ratio of 2h-AUC_{test food} to 2h-AUC_{reference food}, to assess the uric acid metabolic load caused by specific food-derived fructose. Both test food and reference food provided 25 g of fructose. Therefore, the uric acid index was calculated and compared using the independent samples t-test. The uric acid index of honey was higher than that of apple, although no significant difference in the uric acid index was observed between apple and honey ($p=0.153$) as depicted in Figure 3F and Table 5. For either men or women, no significant difference was also found in the uric acid index of apples and honey ($p_{\text{women}}=0.294$, $p_{\text{men}}=0.419$).

Blood glucose measurement

As depicted in Figure 3A, the blood glucose concentration increased significantly at 30 min after the intake of honey and apple, while it only slightly increased after consumption of fructose powder. The highest blood glucose concentration ($5.55 \mu\text{mol/L}$) was observed in the apple-treated group at 30 min, and no difference was observed in the blood glucose concentration between the honey group and apple group ($p=0.151$). Relatively, the blood glucose concentration of participants who consumed fructose powder was the lowest. At 60 min, the blood glucose concentration of participants who consumed honey was the highest ($p<0.001$), while those of participants who consumed apple and fructose powder had similar blood glucose concentration. The 2h-AUC of blood glucose was also calculated. The 2h-AUC of honey was significantly higher than that of apple and fructose ($p<0.01$), as shown in Figure 3B.

Blood pressure evaluation

The effect of food containing fructose on blood pressure is shown in Figure 3C and 3D. No significant difference was observed in the change in diastolic or systolic blood pressure after the ingestion of foods, although the consumption of apples tended to reduce the systolic and diastolic blood pressure.

DISCUSSION

Earlier studies have shown that purine-rich foods such as meat, seafood, and alcohol are important dietary factors contributing to the development of hyperuricemia.³¹ In recent

decades, the consumption of fructose-rich drinks has gained increased attention in relation to the overproduction of uric acid as a byproduct of fructose metabolism.³² Our found that acute intake of fructose-containing foods such as apples, honey, and fructose powder induced a significant increase in SUA concentration in all participants, regardless of the source of fructose. This finding is consistent with that of a previous study, which indicated that the consumption of apples and apple juice containing fructose increased the SUA concentration in an American population;²⁹ in that study, the amount of fructose consumed (26.5 g) was similar to that consumed in our study (25 g). Fructose is a monosaccharide, with a metabolic pathway in the human body that differs greatly from that of glucose, although it is an isomer of glucose. During glucose metabolism, hexokinase catalyzes glucose into glucose-6-phosphate, which then modulates the activity of hexokinase via negative feedback, thereby preventing the overproduction of glucose metabolites. However, fructose metabolism lacks this negative feedback mechanism. When a large amount of fructose was consumed, its rapid metabolism led to the breakdown of adenosine triphosphate (ATP) and phosphate to generate adenosine diphosphate and adenosine monophosphate (AMP). The rapid decrease in phosphate stimulates the production of AMP deaminase, which in turn catalyzes the transformation of AMP to hypoxanthine nucleotide and ultimately to uric acid.³³ The intracellular concentration of uric acid increased and was released into the circulation; as a result, the SUA concentration peaked at 15 min to 1 h after ingestion.³⁴ Animal and human studies revealed that increase in the SUA concentrations mediated the adverse effects of fructose ingestion in our body's metabolism.^{35,36} A previous clinical trial showed that consumption of fructose-containing beverages for 10 weeks significantly elevated the postprandial SUA concentration compared with isocaloric intake of glucose.³⁷ The ratio of fructose to glucose (F:G) in beverages has a great influence on SUA concentration; that is, consumption of a large amount of fructose increases the SUA concentration. A previous study conducted in male participants showed that an F80:G20 drink induced a higher serum concentration of uric acid than a G50:F50 drink.³⁸ Therefore, it is necessary to limit the ordinary intake of fructose in soft drinks to maintain health. Some scholars even warned that gouty participants should avoid consuming fruits.³⁹

Our data demonstrated that the ingestion of pure fructose rapidly increased the uric acid concentration and resulted in the highest uric acid 2h-AUC; moreover, the 2h-AUC of SUA after apple intake was the lowest among the three dietary treatments. Fruits are enriched with bioactive compounds such as vitamin C and polyphenol, and robust epidemiological evidence showed the beneficial effects of fruit consumption on health.⁴⁰ The slow response of the

human body to apple intake may be related to its plentiful bioactive substances, including fiber, vitamin C, and antioxidants such as polyphenols.^{34,38,41} Polyphenols were found to interfere with intestinal sugar transporters, and result in glucose and fructose absorption.⁴² Fibers derived from natural fruits can slow down the digestion of fructose in the small intestine,⁴³ while vitamin C enhances the urinary urate excretion.⁴⁴ We initially proposed the definition of uric acid index and pointed out that it is more optimal for the general population and gouty patients to choose foods based on the uric acid index rather than completely eradicating fruits from their diet. In our study, honey also induced a slow increase in the SUA concentration like apple; in contrast to pure fructose, the slow increase in the SUA concentration may be associated with the presence of phenols in honey, which may impede the intestinal sugar transporters. Our data showed that the uric acid index of honey is higher than that of apple, suggesting that apples may be safer for gouty participants to consume than honey.

Moreover, men exhibited a more rapid increase in the SUA concentration after fructose consumption than women, and the SUA concentration of men did not return to baseline at 120 min. By contrast, the SUA concentration of women returned to baseline by 120 min. The incidence of hyperuricemia and gout in men was four times higher than that in women.⁴⁵ Postmenopausal women who did not undergo hormone replacement therapy had high concentrations of SUA;⁴⁶ meanwhile, exogenous estrogen intervention decreased the SUA concentrations in postmenopausal women. An epidemiological study in Taiwan conducted in 5,896 participants (2,960 women and 2,936 men) reported that the SUA concentration in women increased with age.⁴⁷ The abovementioned studies suggested that estrogen has a positive effect on uric acid metabolism. Moreover, women had stronger capacity to handle the load of dietary fructose. However, there are some limitations in this study. First, this study included healthy young adults, and the response of individuals with gout or compromised kidney function remains unclear. Second, this was a short-term trial; hence, long-term intervention trials are warranted to assess the health effect of fructose from different sources. Finally, due to financial and logistical constraints, two trials were carried out in two groups of participants; however, the serum uric acid responses of these healthy young adults to the same amount of fructose were similar, which strengthens the generalizability of our results.

Conclusion

This study was the first to demonstrate that the consumption of the same amount of fructose derived from different food sources triggered the elevation of SUA concentration to different

extents, exhibiting the trend of fructose powder > honey > apple. A gender difference was found in the metabolism of fructose; the SUA concentration of men exhibited a more rapid increase after fructose consumption compared with that of women. Hence, it is meaningful to establish fructose intake limits by gender. Uric acid index may be a useful tool to guide hyperuricemia or gout patients in choosing appropriate foods that they can consume.

ACKNOWLEDGEMENTS

We appreciate all the volunteers who have participated in the study and the staff at the Ningbo University.

CONFLICT OF INTEREST AND FUNDING DISCLOSURE

The authors declare no conflicts of interest.

This work was supported by the National Natural Science Foundation of China (NSFC; Grant No. 81872620, 81673163), Ningbo Natural Science Foundation (2018A610370), Zhejiang Key laboratory of pathology (201802). This study was sponsored by the K. C. Wong Magna Fund in Ningbo University. This study was supported by Fundamental Research Program of Public Good in Zhejiang Province (LGF18H060006), Zhejiang Provincial Medical and Health Science and Technology plan (2018KY819), and the scientific research projects of Shaoxing University (2019SK003).

REFERENCES

1. Plummer RW, Smith WB, Cone IF, Flint J. Uric acid. The chemistry, physiology and pathology of uric acid and the physiologically important purin bodies, with a discussion of the metabolism in gout. *JAMA*. 1906;12:906. doi: 10.1001/jama.1906.02510390064027.
2. Meneses-Leon J, Denova-Gutierrez E, Castanon-Robles S, Granados-Garcia V, Talavera JO, Rivera-Paredes B et al. Sweetened beverage consumption and the risk of hyperuricemia in Mexican adults: a cross-sectional study. *BMC Public Health*. 2014;14:445. doi: 10.1186/1471-2458-14-445.
3. Bae J, Chun BY, Park PS, Choi BY, Kim MK, Shin MH, Lee YH, Shin DH, Kim SK. Higher consumption of sugar-sweetened soft drinks increases the risk of hyperuricemia in Korean population: The Korean Multi-Rural Communities Cohort Study. *Semin Arthritis Rheum*. 2014;43:654-61. doi: 10.1016/j.semarthrit.2013.10.008.
4. Cho SK, Chang Y, Kim I, Ryu S. U-Shaped Association Between Serum Uric Acid Level and Risk of Mortality: A Cohort Study. *Arthritis Rheumatol*. 2018;70:1122-32. doi: 10.1002/art.40472.
5. King C, Lanaspá MA, Jensen T, Tolan DR, Sanchez-Lozada LG, Johnson RJ. Uric Acid as a Cause of the Metabolic Syndrome. *Contrib Nephrol*. 2018;192:88-102. doi: 10.1159/000484283.

6. Krishnan E, Pandya BJ, Chung L, Hariri A, Dabbous O. Hyperuricemia in young adults and risk of insulin resistance, prediabetes, and diabetes: a 15-year follow-up study. *Am J Epidemiol.* 2012;176:108-16. doi: 10.1093/aje/kws002.
7. Sirota JC, McFann K, Targher G, Johnson RJ, Chonchol M, Jalal DI. Elevated serum uric acid levels are associated with non-alcoholic fatty liver disease independently of metabolic syndrome features in the United States: Liver ultrasound data from the National Health and Nutrition Examination Survey. *Metabolism.* 2013;62:392-9. doi: 10.1016/j.metabol.2012.08.013.
8. Hu L, Hu G, Xu BP, Zhu L, Zhou W, Wang T, Bao H, Cheng X. U-Shaped association of serum uric acid with all-cause and cause-specific mortality in US adults: a cohort study. *J Clin Endocrinol Metab.* 2020;105:e597-609. doi: 10.1210/clinem/dgz068.
9. Fox IH, Kelley WN. Studies on the mechanism of fructose-induced hyperuricemia in man. *Metabolism.* 1972;21:713-21. doi: 10.1016/0026-0495(72)90120-5.
10. Heuckenkamp PU, Zöllner N. Fructose-induced hyperuricaemia. *Lancet.* 1971;1:808-9. doi: 10.1016/S0140-6736(71)91259-1.
11. Mayes PA. Intermediary metabolism of fructose. *Am J Clin Nutr.* 1993;58:754S-65S. doi: 10.1093/ajcn/58.5.754S.
12. Perheentupa J, Raivio K. Fructose-induced hyperuricaemia. *Lancet.* 1967;290:528-31. doi: 10.1016/S0140-6736(67)90494-1.
13. Hanover LM, White JS. Manufacturing, composition, and applications of fructose. *Am J Clin Nutr.* 1993;58:724S-32S. doi: 10.1093/ajcn/58.5.724S.
14. Choi HK, Ford ES. Prevalence of the metabolic syndrome in individuals with hyperuricemia. *Am J Med.* 2007;120:442-7. doi: 10.1016/j.amjmed.2006.06.040.
15. Jamnik J, Rehman S, Blanco Mejia S, de Souza RJ, Khan TA, Leiter LA, Wolever TM, Kendall CW, Jenkins DJ, Sievenpiper JL. Fructose intake and risk of gout and hyperuricemia: a systematic review and meta-analysis of prospective cohort studies. *BMJ Open.* 2016;6:e013191. doi: 10.1136/bmjopen-2016-013191.
16. Li R, Yu K, Li C. Dietary factors and risk of gout and hyperuricemia: a meta-analysis and systematic review. *Asia Pac J Clin Nutr.* 2018;27:1344-56. doi: 10.6133/apjcn.201811_27(6).0022.
17. Ishimoto T, Lanaspa MA, Rivard CJ, Roncal-Jimenez CA, Orlicky DJ, Cicerchi C et al. High-fat and high-sucrose (western) diet induces steatohepatitis that is dependent on fructokinase. *Hepatology.* 2013;58:1632-43. doi: 10.1002/hep.26594.
18. Johnson RJ, Sanchez-Lozada LG, Andrews P, Lanaspa MA. Perspective: A Historical and Scientific Perspective of Sugar and Its Relation with Obesity and Diabetes. *Adv Nutr.* 2017;8:412-22. doi: 10.3945/an.116.014654.
19. Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *Am J Clin Nutr.* 2013;98:1084-102. doi: 10.3945/ajcn.113.058362.

20. Stanhope KL, Bremer AA, Medici V, Nakajima K, Ito Y, Nakano T et al. Consumption of fructose and high fructose corn syrup increase postprandial triglycerides, LDL-cholesterol, and apolipoprotein-B in young men and women. *J Clin Endocrinol Metab.* 2011;96:E1596-605. doi: 10.1210/jc.2011-1251.
21. Yang Q, Zhang Z, Gregg EW, Flanders WD, Merritt R, Hu FB. Added sugar intake and cardiovascular diseases mortality among US adults. *JAMA Intern Med.* 2014;174:516-24. doi: 10.1001/jamainternmed.2013.13563.
22. Park YK, Yetley EA. Intakes and food sources of fructose in the United States. *Am J Clin Nutr.* 1993;58:737S-47S. doi: 10.1093/ajcn/58.5.737S.
23. Cicero AF, Rosticci M, Bove M, Fogacci F, Giovannini M, Urso R, D'Addato S, Borghi C, Brisighella Heart Study G. Serum uric acid change and modification of blood pressure and fasting plasma glucose in an overall healthy population sample: data from the Brisighella heart study. *Ann Med.* 2017;49:275-82. doi: 10.1080/07853890.2016.1222451.
24. Cicero AF, Rosticci M, Cagnati M, Urso R, Scapagnini G, Morbini M, Grandi E, D'Addato S, Borghi C, Brisighella Heart Study G. Serum uric acid and markers of low-density lipoprotein oxidation in nonsmoking healthy subjects: data from the Brisighella Heart Study. *Pol Arch Intern Med.* 2014;124:661-68. doi: 10.20452/pamw.2548.
25. Cicero AF, Rosticci M, Fogacci F, Grandi E, D'Addato S, Borghi C, Brisighella Heart Study G. High serum uric acid is associated to poorly controlled blood pressure and higher arterial stiffness in hypertensive subjects. *Eur J Intern Med.* 2017;37:38-42. doi: 10.1016/j.ejim.2016.07.026.
26. Cicero AF, Desideri G, Grossi G, Urso R, Rosticci M, D'Addato S, Borghi C, Brisighella Heart Study G. Serum uric acid and impaired cognitive function in a cohort of healthy young elderly: data from the Brisighella Study. *Intern Emerg Med.* 2015;10:25-31. doi: 10.1007/s11739-014-1098-z.
27. Cicero AF, Rosticci M, Parini A, Baronio C, D'Addato S, Borghi C. Serum uric acid is inversely proportional to estimated stroke volume and cardiac output in a large sample of pharmacologically untreated subjects: data from the Brisighella Heart Study. *Intern Emerg Med.* 2014;9:655-60. doi: 10.1007/s11739-013-1016-9.
28. Wiseman M. The second World Cancer Research Fund/American Institute for Cancer Research expert report. Food, nutrition, physical activity, and the prevention of cancer: a global perspective. *Proc Nutr Soc.* 2008;67:253-6. doi: 10.1017/S002966510800712X.
29. White SJ, Carran EL, Reynolds AN, Haszard JJ, Venn BJ. The effects of apples and apple juice on acute plasma uric acid concentration: a randomized controlled trial. *Am J Clin Nutr.* 2018;107:165-72. doi: 10.1093/ajcn/nqx059.
30. Becker MA, Schumacher HR, Wortmann RL, MacDonald PA, Eustace D, Palo WA, Streit J, Joseph-Ridge N. Febuxostat compared with allopurinol in patients with hyperuricemia and gout. *N Engl J Med.* 2005;353:2450-61. doi: 10.1056/NEJMoa050373.
31. Teng GG, Tan CS, Santosa A, Saag KG, Yuan JM, Koh WP. Serum urate levels and consumption of common beverages and alcohol among Chinese in Singapore. *Arthritis Care Res (Hoboken).* 2013;65:1432-40. doi: 10.1002/acr.21999.

32. Bombback AS, Derebail VK, Shoham DA, Anderson CA, Steffen LM, Rosamond WD, Kshirsagar AV. Sugar-sweetened soda consumption, hyperuricemia, and kidney disease. *Kidney Int.* 2010;77:609-16. doi: 10.1038/ki.2009.500.
33. Lanaspá MA, Sánchez-Lozada LG, Choi YJ, Cicerchi C, Kanbay M, Roncal-Jiménez CA et al. Uric acid induces hepatic steatosis by generation of mitochondrial oxidative stress. *J Biol Chem.* 2012;287:40732-44. doi: 10.1074/jbc.M112.399899.
34. Le MT, Frye RF, Rivard CJ, Cheng J, McFann KK, Segal MS, Johnson RJ, Johnson JA. Effects of high-fructose corn syrup and sucrose on the pharmacokinetics of fructose and acute metabolic and hemodynamic responses in healthy subjects. *Metabolism.* 2012;61:641-51. doi: 10.1016/j.metabol.2011.09.013.
35. Nakagawa T, Hu H, Zharikov S, Tuttle KR, Short RA, Glushakova O et al. A causal role for uric acid in fructose-induced metabolic syndrome. *Am J Physiol Renal Physiol.* 2006;290:F625-31. doi: 10.1152/ajprenal.00140.2005.
36. Pérez-Pozo SE, Schold J, Nakagawa T, Sánchez-Lozada LG, Johnson RJ, Lillo JL. Excessive fructose intake induces the features of metabolic syndrome in healthy adult men: role of uric acid in the hypertensive response. *Int J Obes (Lond).* 2010;34:454-61. doi: 10.1038/ijo.2009.259.
37. Cox CL, Stanhope KL, Schwarz JM, Graham JL, Hatcher B, Griffen SC et al. Consumption of fructose-sweetened beverages for 10 weeks reduces net fat oxidation and energy expenditure in overweight/obese men and women. *Eur J Clin Nutr.* 2012;66:201-8. doi: 10.1038/ejcn.2011.159.
38. Akhavan T, Anderson GH. Effects of glucose-to-fructose ratios in solutions on subjective satiety, food intake, and satiety hormones in young men. *Am J Clin Nutr.* 2007;86:1354-63. doi: 10.1093/ajcn/86.5.1354.
39. Buja LM. Medical education today: all that glitters is not gold. *BMC Med Educ.* 2019;19:110. doi: 10.1186/s12909-019-1535-9.
40. Du H, Li L, Bennett D, Guo Y, Key TJ, Bian Z et al. Fresh Fruit Consumption and Major Cardiovascular Disease in China. *N Engl J Med.* 2016;374:1332-43. doi: 10.1056/NEJMoal501451.
41. Nakagawa T, Lanaspá MA, Johnson RJ. The effects of fruit consumption in patients with hyperuricaemia or gout. *Rheumatology (Oxford).* 2019;58:1133-41. doi: 10.1093/rheumatology/kez128.
42. Loureiro G, Martel F. The effect of dietary polyphenols on intestinal absorption of glucose and fructose: Relation with obesity and type 2 diabetes. *Food Reviews International.* 2019;35:390-406. doi: 10.1080/87559129.2019.1573432.
43. Björck I, Granfeldt Y, Liljeberg H, Tovar J, Asp NG. Food properties affecting the digestion and absorption of carbohydrates. *Am J Clin Nutr.* 1994;59:699S-705S. doi: 10.1093/ajcn/59.3.699S.
44. Stein HB, Hasan A, Fox IH. Ascorbic acid-induced uricosuria. A consequence of megavitamin therapy. *Ann Intern Med.* 1976;84:385-8. doi: 10.7326/0003-4819-84-4-385.
45. Kuo CF, Grainger MJ, Zhang W, Doherty M. Global epidemiology of gout: prevalence, incidence and risk factors. *Nat Rev Rheumatol.* 2015;11:649-62. doi: 10.1038/nrrheum.2015.91.

46. Mundy HR, Lee PJ. Glycogenesis type I and diabetes mellitus: a common mechanism for renal dysfunction?. *Med Hypotheses*. 2002;59:110-4. doi: 10.1016/s0306-9877(02)00199-8.
47. Adamopoulos D, Vlassopoulos C, Seitanides B, Contoyiannis P, Vassilopoulos P. The relationship of sex steroids to uric acid levels in plasma and urine. *Acta Endocrinol (Copenh)*. 1977;85:198-208. doi: 10.1530/acta.0.0850198.

Not Proof Read

Table 1. Anthropometric characteristics of participants[†]

Parameters	Apple trial		Honey trial	
	Apple(n=18)	Fructose(n=18)	Honey(n=21)	Fructose(n=21)
Age (y)	22.5±2.2	22.5±2.2	22.8±1.7	22.8±1.7
women, n (%)	12 (66.67)	12 (66.67)	12 (57.14)	12 (57.14)
men, n (%)	6 (33.33)	6 (33.33)	9 (42.86)	9 (42.86)
BMI (kg/m ²)	20.72±1.72	20.72±1.72	20.16±2.05	20.16±2.05
Systolic blood pressure (mm Hg)	116.39±11.6	115.06±10.18	110.24±8.54	111.05±11.29
Diastolic blood pressure (mm Hg)	78.39±9.00	80.22±8.33	70.81±7.96	70.14±7.66
Blood glucose (mmol/L)	4.09±0.38	4.17±0.24	4.17±0.61	3.92±0.57

DBP: diastolic blood pressure; SBP: systolic blood pressure.

[†]Data were obtained at 0 min of dietary intervention and presented as mean±SD.

Table 2. Serum uric acid concentration at different time points after food intervention[†]

	Gender	SUA at 0min, $\mu\text{mol/L}$ [‡]		n	SUA test food –SUA reference food, $\mu\text{mol/L}$ [‡]					
		SUA test Food	SUA reference food		30 min	p [§]	60 min	p [§]	120 min	p [§]
Apple trail	women	286.67 \pm 37.00	307.92 \pm 75.81	24	-22.53 \pm 8.41*	0.01	-20.62 \pm 7.74*	0.02	-18.61 \pm 10.40	0.09
	men	342.67 \pm 36.84	357.83 \pm 45.84	12	-6.91 \pm 11.32	0.56	-18.22 \pm 9.03	0.07	-12.43 \pm 9.00	0.20
	All subjects	305.33 \pm 13.51	324.56 \pm 13.51	29	-35.17 \pm 22.50	0.54	-37.33 \pm 22.32	0.63	-36.33 \pm 22.68	0.51
Honey trail	women	293.00 \pm 40.79	299.75 \pm 36.76	24	-13.53 \pm 7.45	0.08	-11.99 \pm 7.97	0.15	2.80 \pm 6.48	0.67
	men	345.22 \pm 39.19	351.44 \pm 78.60	18	-12.58 \pm 16.92	0.47	-5.80 \pm 20.27	0.78	1.05 \pm 17.35	0.95
	All subjects	315.38 \pm 12.51	323.25 \pm 12.82	42	-22.33 \pm 21.09	0.88	-18.71 \pm 20.92	0.89	-6.39 \pm 21.26	1.00

[†]Participants with complete data (0 min, 30 min, 60 min, and 120 min) and normal uric acid concentration (men: <416.4 $\mu\text{mol/L}$; women: <356.9 $\mu\text{mol/L}$) were enrolled.

[‡]Data at 0 min were presented as mean \pm SD; results at 30 min, 60 min, and 120 min were presented as mean \pm SEM.

[§]General linear model: repeated measures analysis was used to compare the effects of dietary treatments. The treatments were considered as fixed effects, while the participants were considered as random effects; values at 0 min were included in the model as covariates.

* p <0.05.

Table 3. Comparison of SUA concentration at different time points after meal by gender[†]

	Foods	n	SUA men –SUA women at 30 min [‡]		p [§]	SUA men –SUA women at 60 min [‡]		p [§]	SUA men –SUA women at 120 min [‡]		p [§]
Apple trail	Apple	18	2.69 \pm 7.92	0.74	5.09 \pm 8.38	0.55	18.67 \pm 9.95	0.08			
	Fructose	18	-5.36 \pm 13.41	0.70	7.19 \pm 11.33	0.54	5.54 \pm 14.56	0.71			
Honey trail	Honey	21	13.02 \pm 16.11	0.43	12.54 \pm 19.03	0.52	5.30 \pm 16.54	0.75			
	Fructose	21	22.00 \pm 11.77	0.08	20.98 \pm 12.78	0.12	21.35 \pm 10.59*	0.04			

[†]Participants with complete data (0 min, 30 min, 60 min, and 120 min) and normal uric acid concentration (men: <416.4 $\mu\text{mol/L}$; women: <356.9 $\mu\text{mol/L}$) were enrolled.

[‡]Results at 30 min, 60 min, and 120 min were presented as mean \pm SEM.

[§]General linear model: repeated measures analysis was used to compare the SUA concentrations at different time points of food intake by gender. Gender was considered as fixed effects, while the participants were considered as random effects; values at 0 min were included in the model as covariates.

* p <0.05.

Table 4. 2h-AUC of SUA of apple, honey by gender[†]

	FOODS	2h-AUC men [‡]	2h-AUC women [‡]	n	<i>p</i> [§]
Apple trail	Apple	42595.00±4221.52**	35270.00±4561.66	18	0.005
	Fructose	45827.50±6044.01	39823.75±8747.86	18	0.153
Honey trail	Honey	44788.33±8285.04*	35898.75±5349.75	21	0.015
	Fructose	46273.33±10731.42*	37608.75±4064.67	21	0.047

[†]AUC of uric acid was calculated by GraphPad Prism software using the trapezoidal method.

[‡]All data were presented as mean±SD.

[§]The normality of data distribution was assessed using Shapiro-Wilk test. The data with normal and non-normal distribution were analyzed using one-way analysis of variance and non-parametric test, respectively.

p*<0.05, *p*<0.01 when compared to 2h-AUC women.

Table 5. The uric acid index of apple and honey

Gender	Testing Food	n	Uric acid index [†]	<i>p</i> [‡]
Woman	Apple	12	0.90±0.12	0.294
	honey	12	0.96±0.13	
Men	Apple	6	0.94±0.10	0.419
	honey	9	0.98±0.10	
All subjects	Apple	18	0.91±0.12	0.153
	honey	21	0.97±0.12	

All data were presented as mean±SD.

[†]Uric acid index was newly defined as the ratio of 2h-AUC of the test food (contain 25 g fructose) to that of the reference food (pure fructose powder, 25g).

[‡]The normality of data distribution was assessed using Shapiro-Wilk test. The data with normal and non-normal distribution were analyzed using independent-samples T-test and non-parametric test, respectively.




Figure 1. The Consolidated Standards of Reporting Trials diagram of the participant recruitment process. Sixty-four college students (apple trial, n=29; honey trial, n=35) aged 22–26 years were enrolled. Prior to the administration of the dietary intervention, seven (men) and three (two men and one woman) volunteers dropped out of apple trial and honey trial, respectively, due to unrelated personal reasons. Four participants (three men and one woman) in the apple trial and 11 participants (eight men and three women) in the honey trial were excluded from the analysis owing to the occurrence hyperuricemia. Overall, 78 full datasets were included in the study.




Figure 2. Serum uric acid (SUA) concentration after consuming 25 g of fructose derived from different sources. A: SUA concentration of all participants at 0 min, 30 min, 60 min, and 120 min. B: SUA of women at 0 min, 30 min, 60 min, and 120 min. C: SUA of men at 0 min, 30 min, 60 min, and 120 min. D: Area under the curve (AUC) for SUA in all participants. E: AUC for SUA in women and men. F: Uric acid index in all participants. G: Uric acid index in women and men. * $p < 0.05$.




Figure 3. Effect of apple, honey, and fructose consumption on blood glucose and blood pressure. A: Blood glucose concentration at 0 min, 30 min, 60 min, and 120 min. B: Area under the curve of blood glucose. C and D: Changes in systolic blood pressure and diastolic blood pressure between 0 min and 120 min. Values without a common superscript are significantly different; $p < 0.05$.




Figure 2. Conceptual diagram of various sources of fructose caused an elevation of serum uric acid concentration to different extent. Fructose readily absorbed can be rapidly metabolized by fructokinase in liver to produce fructose-1-phosphate, leading to a depletion in intracellular ATP and generation of IMP. Then IMP is further degraded to hypoxanthine and xanthine, that is used by xanthine oxidase to generate end-product uric acid. Abbreviations: KHK, ketohexokinase; ATP, adenosine triphosphate; ADP, adenosine diphosphate; AMP, adenosine monophosphate; SUA, serum uric acid; AUC, the areas under the curve.