

This author's PDF version corresponds to the article as it appeared upon acceptance. Fully formatted PDF versions will be made available soon.

Is abdominal adiposity in healthy Sri Lankan neonates different from the rest of the world?

doi: 10.6133/apjcn.202203/PP.0005

Published online: March 2022

Running title: Abdominal adiposity in Sri Lankan neonates

Marianne Nishani Lucas DCH, MD, MRCPCH 1,2, Pulani Lanerolle PhD1, Upul Senarath MD, PhD1, Andrew Peter Hills PhD1, Vithanage Pujitha Wickramasinghe DCH, MD, PhD¹

¹Department of Medicine, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand

²Gastroenterology Unit, Dunedin Hospital, Southern District Health Board, Dunedin, New Zealand

Authors' email addresses and contributions:

Marianne Nishani Lucas, nishani@pdt.cmb.ac.lk, contributed by developing the study design, executing the clinical data collection, data analysis and interpretation and preparation of the draft manuscript

Pulani Lanerolle, pulani@bmb.cmb.ac.lk contributed by developing the study concept and design, supervising data interpretation, providing critique, revision of the manuscript and overall scientific management

Upul Senarath, upul@commed.cmb.ac.lk contributed by developing the study concept and design, supervising statistical analysis and data interpretation, providing critique and revision of the manuscript

Andrew Peter Hills, BEd, MSc, PhD, andrew.hills@utas.edu.au contributed by developing the study concept and design, providing critique and revision of the manuscript

Vithanage Pujitha Wickramasinghe, pujitha@pdt.cmb.ac.lk contributed by developing the study concept and design, supervising data interpretation, providing critique, revision of the manuscript and overall scientific management

Corresponding Author: Dr Nishani Lucas, Department of Paediatrics, Faculty of Medicine, University of Colombo, Kynsey Road, Colombo 8, Sri Lanka. Tel: +94 777 745294. Email: nishani@pdt.cmb.ac.lk

ABSTRACT

Background and Objectives: Adiposity at birth is a predictor of childhood obesity. Abdominal circumference (AC) at birth has been shown to correlate well with visceral adipose tissue and abdominal subcutaneous adipose tissue. Adiposity differs according to ethnicity and geography. The aim of this study was to describe the anthropometry derived adiposity phenotype in neonates from Colombo, Sri Lanka and compare it with global data. **Methods and Study Design:** Birth anthropometry was performed within 12-24 hours by the same investigator as part of a prospective cohort study on healthy term babies, at a tertiary care hospital in Colombo, Sri Lanka, 2015-2019. The anthropometry derived adiposity phenotype was indicated by skinfold thickness, AC and upper arm fat area (UFA) derived from the mid-upper arm circumference (MUAC). **Results:** Sri Lankan neonates had a significantly lower weight with significantly higher AC ($n=337$, 2.9 ± 0.4 kg, 30.6 ± 2.3 cm) compared to Canadian ($n=389$, 3.5 ± 0.02 kg, 29.9 ± 2.1 cm; $p<0.001$) and Australian ($n=1270$, 3.4 ± 0.4 kg, 28.5 ± 1.9 cm; $p<0.001$) neonates. Anthropometry derived adiposity at birth showed a significant correlation with weight and BMI of both mother and father ($p<0.05$) as opposed to their income or education ($p>0.05$). **Conclusions:** Healthy neonates from Colombo, Sri Lanka demonstrated significantly higher AC despite significantly lower weight, indicating increased abdominal adiposity compared to neonates from high-income countries as well as Indian neonates with the thin-fat phenotype.

Key Words: anthropometry derived adiposity, abdominal circumference, Sri Lankan neonate, birth anthropometry, abdominal adiposity at birth

INTRODUCTION

Adiposity at birth is a significant predictor of childhood obesity from 2 to 6 years of age.¹ Abdominal circumference at birth has been shown to have a high correlation with ultrasound measured abdominal adipose tissue, both subcutaneous and visceral.² Although there are only few studies describing the abdominal circumference at birth, waist circumference in children has been shown to be the best predictor of visceral adipose tissue assessed by MRI.³ Abdominal obesity, based on waist circumference has also been reported as a better predictor of cardiovascular risk factors compared to BMI-defined obesity in children.⁴ Skinfold thickness (SFT) is widely used as an indicator of subcutaneous adipose tissue.⁵ Both waist circumference and SFT of subscapular and supra-iliac regions have been associated with increased risk of liver disease and metabolic dysregulation in children.⁶

Studies from New Zealand and the United Kingdom have revealed that Asian neonates have higher adiposity despite a smaller body size compared to Caucasians, thereby suggesting that adiposity differs according to ethnicity.^{7,8}

The ‘thin-fat phenotype’ was first described in Indian neonates by Yajnik et al, in 2003. These babies had higher subscapular SFT depicting preserved subcutaneous adipose tissue indicating the ‘fat’ of the ‘thin-fat’ phenotype and were smaller in size with lower weight, length, circumferences of head, arm and abdomen indicating the ‘thin’ of this phenotype, when compared to British Caucasian babies.⁹ The same ‘thin-fat’ phenotype was reported in South Asian neonates in the Netherlands when compared to Dutch Caucasian neonates as well as neonates of fourth to fifth generation Indian immigrants in Surinam, South America, South Indian neonates from Mysore, and in the Vellore birth cohort when compared to Caucasian babies born in Southampton, UK.¹⁰⁻¹³ Persistence of this thin-fat phenotype at 4 years of age with higher SFT compared to the UK population was shown in the Mysore study.¹² Indian babies had higher abdominal adiposity in both subcutaneous and intra-abdominal compartments when assessed by MRI, despite having a very small abdominal circumference compared to the British babies.¹⁴ The thin-fat phenotype has been described as fat being added to an already thin frame, where babies have a relatively high fat mass despite a low lean mass.⁹ Pomeroy’s analysis suggests that the low lean mass phenotype characteristic of the South Asian ancestry has existed for a minimum of 11,000 years.¹⁵ Low lean mass appears to have been favoured through evolution as an adaptation to the hot climate by maximising heat loss via an increased surface area to volume ratio. In addition, repeated famines causing high mortality in South Asia in the 19th and 20th centuries are also thought to have led to the selection of genes associated with low lean mass through intergenerational plasticity.¹⁵ Changing from hunter gatherers to agriculture during the Holocene era as well as the increase in sugar and fat content in the diet associated with reduced activity levels over the past 2 decades, are thought to result in a relatively high fat mass despite smaller anthropometric measurements.¹⁶

Although Sri Lanka is a low-middle-income country, it has excellent health indicators on par with middle and high-income countries.^{17,18} Colombo has the highest monthly household income and the lowest poverty head count index amongst the 25 administrative districts in Sri Lanka.¹⁹ Therefore, we were interested to know whether babies born in Colombo demonstrated the “thin-fat” phenotype described in Indian South Asians due to similar geography or if they differ based on better health indicators.

This is the first paper to describe anthropometry derived neonatal adiposity from Colombo, Sri Lanka. We also present a secondary analysis of our data to determine whether our study population demonstrates the “thin-fat” phenotype described in South Asian babies of Indian descent, and how the adiposity of our study population compares to published data from high-income countries.

MATERIALS AND METHODS

Study design and setting

Birth data was obtained as part of an observational, prospective, cohort study on infant body composition from birth to 2 years. The study was conducted at the Neonatal Unit of the University of Colombo, De Soyza Hospital for Women, Colombo, a tertiary care maternity hospital in Sri Lanka, from July 2015 to December 2019.

Study population

Trained research assistants visited the Obstetric Unit twice a day, on all weekdays and approached all the pregnant women admitted between 37 weeks and 41 weeks + 6 days period of amenorrhea (POA) and screened those who gave informed written consent to join the study. All consecutive mothers and babies fulfilling the inclusion criteria were selected via purposive sampling.

Inclusion criteria

Non-smoking mothers with a singleton pregnancy, between 37 weeks and 41 weeks + 6 days POA, aged more than 18 years, living in the study area (Colombo District), who intended to breastfeed and gave informed written consent were recruited prior to delivery. All newborns who did not have morbidity according to supplementary table 1 were included in the study.

Exclusion criteria

Women who lived outside the study area were excluded. Babies with an Apgar score <8 at 5 minutes of age or had congenital anomalies or required admission to the neonatal intensive care unit, or had morbidity were excluded.

Sample size calculation

Males vs females

The pooled standard deviation (SD) for upper arm fat area (UFA) was assumed to be 48 cm² according to the data given for Caucasian newborns.²⁰ A sample size of 161 per group (males

vs females) were required to detect a true difference of 15 cm² in the mean UFA between boys and girls to achieve a power of 80% and a level of significance of 5% (two-sided).²⁰

Comparison with global data

The pooled SD for mid upper arm circumference (MUAC) was 0.9 cm according to our data as well as for Pomeroy's study on Australian newborns.²¹ A sample size of 159 per group (each country) was required to detect a true difference of 0.4 cm in the mean MUAC between our study data and the Australian newborns, to achieve a power of 80% and significance of 5% (two-sided).²¹

Data collection

Data were collected using interview-administered questionnaires and data recording forms. Gestational age was derived by antenatal ultrasound scan measurement of crown rump length at 8-13 weeks period of amenorrhea (POA). If this was not available, an ultrasound scan of biparietal diameter at 13-20 weeks POA, was used. Last regular menstrual period was used if both the above-mentioned ultrasound scan measurements were not available.

Anthropometry

Babies' measurements were undertaken within 12-24 hours of birth. Weight was measured to the nearest 5 g, using an electronic weighing scale (Seca 334, Seca GmbH®, Hamburg, Germany) which was calibrated twice weekly. Length was measured using an infantometer (Seca 417®, Seca GmbH, Hamburg, Germany) to the nearest 1 mm, and circumferences were measured using a non-stretchable measuring tape to the nearest 1 mm (Seca 212®, Seca GmbH, Hamburg, Germany). Biceps, triceps, subscapular and supra-iliac SFT was measured using a Harpenden skinfold calliper to the nearest 0.2 mm (Baty International, Burgess Hill, West Sussex, UK). Each SFT was read after 2 seconds, consistent with the WHO Multicentre Growth Reference Study (MGRS).²² Head circumference and mid upper-arm circumference (MUAC) were measured according to WHO-MGRS methodology.²² Chest circumference was just below the nipple level) and abdominal circumference was measured at the level of the umbilicus.^{9,23}

Quality control was ensured using similar methodology to the WHO-MGRS study.²² All measurements were performed with the same instruments and by the same investigator who was trained by an International Society for the Advancement of Kinanthropometry (ISAK) Level 2 accredited anthropometrist.

Technical error of measurement (TEM) was calculated as $\sqrt{\sum (d_i^2 / 2n)}$, where d_i is the difference between the i^{th} participant's test and retest measurements and n is the number of participants.²⁴ TEM% was calculated by dividing the TEM by the variable average value, i.e., the arithmetic mean calculated by averaging the mean values for the 1st and 2nd measurements. Intra-observer TEM considered acceptable for a skilled anthropometrist was 5% for SFT and 1% for other measures whereas inter-observer TEMs were 7.5% for SFT and 1.5% for other measures.² Intra-observer TEM was 0.0005% for weight, 0.01% for length, 0.01% - 0.03% for circumferences and 0.2% - 0.4% for SFT.

Each measurement was repeated independently by a second trained anthropometrist and quality control and observer reliability monitored. Intra-observer TEM was 0.0005% for weight, 0.01% for length, 0.03% - 0.08% for circumferences and 0.8% - 0.9% for SFT. If the difference between two measurements exceeded 50 g (birth weight), 7 mm (length), 5 mm (circumferences) and 1 mm (SFT), both observers independently repeated the measurement and, if necessary, a third time.²⁴ Standardization of anthropometry was conducted every 6 months. The inter-observer TEM was 0.09% for weight, 0.3% for length, 0.6% - 1.2% for circumferences, and 4.5% - 6.7% for SFT measurements.

The upper-arm composition was assessed using upper-arm muscle area (UMA), upper-arm fat area (UFA), arm-fat index (AFI), upper-arm fat estimate (UFE) and upper-arm muscle estimate (UME), calculated from MUAC and triceps (TSF) SFT.^{26,27} Estimated muscle circumferences were calculated based on the equation of Gurney and Jelliffe.²⁸

Mid-arm muscle circumference (MAMC)(cm) = limb circumference (cm) – π (limb skinfold). Triceps SFT is considered as limb SFT (cm).

Upper limb fat area was calculated based on the equations of Frisancho as follows:²⁹

$$\text{Limb area (cm}^2\text{)} = (\text{limb circumference (cm)})^2 / 4\pi$$

$$\text{Upper limb muscle area (UMA) (cm}^2\text{)} = (\text{muscle circumference (cm)})^2 / 4\pi$$

$$\text{Upper limb fat area (UFA) (cm}^2\text{)} = \text{limb area (cm}^2\text{)} - \text{muscle area (cm}^2\text{)}.$$

Arm fat index (AFI), upper-arm fat estimate (UFE) and upper-arm muscle estimate (UME) were calculated as follows, as described by Jaswant, Rolland-Cachera and Frisancho.^{26,27,29}

$$\text{AFI} = \text{UFA} / \{(\text{MUAC})^2 / (4 \times \pi)\} \times 100$$

$$\text{UFE} = \text{MUAC} \times (\text{TSF} / 2)$$

$$\text{UME} = \{(\text{MUAC})^2 / (4 \times \pi)\} - \text{UFE}$$

The inter observer TEM for body mass index (BMI)=0.8%, ponderal index (PI)=1.4%, MAMC=1.7%, UMA=3.5%, UFA=6.5%, AFI=6%, UFE=7%, UME=3.8% and sum of SFT=4.2%.

Comparison of newborn anthropometry and derived adiposity to similar data from other countries

A PubMed and Google Scholar search on anthropometry and its derived adiposity at birth using the keywords “newborn”, “neonate”, “anthropometry”, “adiposity” identified papers which reported on newborn anthropometry derived adiposity. Studies available in English which included babies born between 37-42 weeks gestation, from singleton pregnancies, measured within 0 – 5 days of birth, which reported a measure of adiposity i.e., upper arm adiposity, abdominal circumference or SFT, were included in the study. Abdominal circumference was included, only if measured at the level of the umbilicus. The Southampton UK babies’ abdominal circumference was measured at the level of the xiphisternum and was not included. Anthropometric measurements were conducted by a trained person/s in all studies. Different instruments were used for measurement.

Ethics approval

The Ethics Review Committee of the Faculty of Medicine, University of Colombo reviewed the protocol and approved the study (EC-14-145).

Statistical analysis

Statistical analysis was completed using SPSS version 26.0 for Windows. T-test for independent samples were used for the comparison between males and females. Spearman correlation was used to determine the association between each anthropometric measure and parental factor, due to the non-normal distribution of the parental factors as assessed by the Shapiro Wilk test. Published data on newborn anthropometry and derived adiposity from other countries were compared to our study data, by conducting paired comparisons using the independent samples T test on the published summary data of each study versus our study findings.

RESULTS

All mothers admitted to the University Unit on weekdays during the study period were screened (n=4140), however only 427 mothers fulfilled the inclusion criteria prior to delivery.

A significant proportion of mothers had come from different parts of the island, to deliver at a tertiary care centre and did not qualify based on living outside the study area. Only 344 mothers consented to participate in the study after delivery, as many had difficulties in committing to the monthly follow-up required for the longitudinal study. Seven records were not included, due to incomplete birth data, resulting in a study population of 337 newborns. Socio demographic characteristics of the study population are described in Table 1.

The majority of parents were married (98.8%) and the mean age of mother and father was 29 and 32 years, respectively. Both parents had a mean duration of 11 years of education. All except 3 fathers were employed whereas the majority (61.7%) of mothers were housewives. The median family monthly income was Sri Lankan Rupees 30,000 (200 USD). Many babies were delivered vaginally (62%). The mean gestational age was 39 weeks and 51% were males. All babies received the first breastfeed within the first hour and were exclusively breastfed at the time of discharge. Differences in anthropometry and derived adiposity at birth between girls and boys are given in Table 2 as mean and SD.

Boys were bigger than girls at birth, with a significantly higher ($p < 0.05$) weight, length, head and chest circumference. However, there was no significant difference in BMI, ponderal index, measures of muscle mass or fat mass, except for a marginally higher abdominal circumference / length ratio in girls ($p = 0.051$).

Associations between anthropometric measures of neonatal adiposity and parental factors in our study population revealed that abdominal circumference showed a positive correlation with maternal weight ($r = 0.154$, $p = 0.004$), maternal BMI ($r = 0.126$, $p = 0.018$) and paternal BMI ($r = 0.243$, $p = 0.036$). Similarly, abdominal circumference / length was correlated with maternal weight ($r = 0.118$, $p = 0.028$), maternal BMI ($r = 0.109$, $p = 0.042$), paternal weight ($r = 0.237$, $p = 0.038$) and paternal BMI ($r = 0.280$, $p = 0.015$). Subscapular SFT showed a positive correlation with maternal weight ($r = 0.205$, $p < 0.001$), maternal BMI ($r = 0.135$, $p = 0.012$) and maternal height ($r = 0.105$, $p = 0.049$). Monthly income, parental age and parental education was not significantly correlated ($p > 0.05$) with any of the anthropometric measures of adiposity.

Anthropometry derived adiposity in our study population of urban Sri Lankan neonates were compared with similar published data from around the world which were based on neonates born as the outcome of a singleton pregnancy, at a gestation of 37 weeks or more. The timing of measurements varied from 0 to 5 days. Abdominal circumference was taken for comparison only when measured at the level of the umbilicus.

Table 3 shows the comparison of anthropometric data of our study population, denoted as the Sri Lankan data, with published data from other countries.

Our study population UFA and SFT were significantly ($p<0.001$) higher than the US neonates despite smaller size with significantly lower weight and length. Abdominal circumference was not measured in the US population.²⁰

Our study population had no significant difference in subscapular SFT despite smaller size as evidenced by significantly lower weight, length, head circumference and triceps SFT compared to the Canadian Caucasians and Canadian South Asians. Amongst the Canadian neonates, Canadian South Asians showed a higher waist circumference, higher triceps and subscapular SFT despite being smaller in size (lower weight, length and head circumference), when compared to the Canadian Caucasians.³⁰

Our study population was significantly bigger in size than rural Indian neonates, in Pune in all parameters including SFT and abdominal circumference and had significantly higher subscapular SFT than the Caucasian babies in Southampton, UK, despite being significantly smaller in size with lower weight, length, and head circumference.⁹ Abdominal circumference could not be compared due to the difference in the site of measurement.⁹ Similarly, significantly higher triceps and subscapular SFT was reported despite being smaller in size with significantly lower weight and head circumference compared to the 4th generation Indian immigrants in Surinam, South America. Abdominal circumference was not significantly different in our study population and the 4th generation Indian neonates, in Surinam.¹¹

Compared to Australian neonates, our study population showed significantly higher abdominal circumference with no significant difference in the subscapular SFT, despite being significantly smaller with lower weight, head circumference, mid-upper-arm-circumference and triceps SFT.²¹

DISCUSSION

Female neonates in our cohort were smaller than their male counterparts with significantly lower weight, length and head circumference, similar to the WHO growth standards. Despite a smaller body size, female babies had a marginally increased abdominal circumference / length ratio ($p=0.051$) with no significant differences in other fat mass indicators, suggestive of increased abdominal adiposity, compared to males. Increased adiposity in female neonates, was also described using UFA at birth in USA, using SFT at birth in UK, intrahepatocellular lipid deposition in MRI at 13 days of life in UK, and using 18O dilution method at 4-6 months of age in Sri Lanka.^{20,31,32,33}

Neonatal adiposity has been found to be associated with parental adiposity with evidence of intergenerational transmission.^{34,35,36} This led us to look for an association between the anthropometry derived adiposity in neonates with parental anthropometry. We found that both maternal and paternal anthropometry were associated with different aspects of neonatal adiposity. Maternal weight and BMI correlated with subscapular skinfold thickness, abdominal circumference and UFA in the offspring suggesting that maternal anthropometry affected both central and peripheral fat deposition in the offspring and contributed to both subcutaneous and visceral adipose tissue. Paternal weight and BMI correlated with neonatal abdominal circumference suggesting that paternal anthropometry predominantly affected abdominal adiposity. Ornellas et al. and Catalano et al. reported similar findings where both paternal obesity and maternal BMI were found to be strong predictors for childhood obesity.^{37,38} The lasting effect of paternal obesity on their offspring was demonstrated by its correlation with the offspring's growth hormone levels at 2 years of age.³⁹ Intergenerational transmission of adiposity has been reported as the most likely explanation for the effect of parental adiposity on neonatal adiposity.³⁷

Our study population also demonstrated significantly higher abdominal circumference suggestive of increased abdominal adiposity and relatively preserved or increased subscapular SFT, despite significantly smaller size, compared to American, British, Canadian and Australian neonates.^{9,20,21,30} Similar findings were noted in the comparison of Canadian South Asians to Canadian Caucasian neonates by Anand et al.³⁰ Higher subscapular SFT at birth in African-Americans, Hispanics and Asians compared to Caucasians and higher abdominal adiposity as quantified by MRI in South Asian neonates compared to Chinese neonates in Singapore are also in agreement with our study findings.^{40,41}

Significantly higher abdominal circumference, indicating abdominal adiposity was a more consistent finding in our study population, compared to skinfold thickness indicating higher subcutaneous adipose tissue. Higher abdominal circumference together with higher skinfold thickness, indicating higher visceral and subcutaneous adipose tissue, in our study population from Colombo Sri Lanka, differ from the thin-fat phenotype described by Yajnik in the Pune study, where the smaller Indian baby demonstrated a smaller abdominal circumference with higher skinfold thickness implying increased subcutaneous but not visceral adipose tissue.⁹ This implies that our study population from Colombo Sri Lanka with a significantly higher abdominal circumference may have a higher amount of abdominal adiposity with higher risk of obesity associated complications compared to the Indian baby who was also reported to have increased subcutaneous and visceral fat via MRI compared to their Caucasian

counterparts, despite a smaller abdominal circumference.¹⁴ Sri Lanka's excellent health indicators, which have surpassed the rest of South Asia and are on par with high and upper middle-income countries, may indicate a greater lifestyle change in Sri Lanka, that is closer to that of high-income countries, with higher consumption of refined sugars and fats along with lower activity levels. This life-style change may increase the risk of abdominal adiposity in Sri Lankans compared to other South Asians in neighbouring countries and explain the higher abdominal circumference in our study population that makes them different from the thin-fat phenotype described in Indian newborns.

Healthy neonates born in Colombo Sri Lanka had a significantly higher abdominal circumference, indicating higher abdominal adiposity despite smaller size compared to babies from high income countries. Childhood adiposity as early as 3 years was associated with non-alcoholic fatty liver disease (NAFLD), in 17-year-old adolescents.⁴² Sri Lanka has a high prevalence of NAFLD in adulthood (34%) compared to Africa (13.8%), Middle East (31%), and India (16.6%).^{43,44} Further studies are needed to establish if babies with higher abdominal circumference at the time of birth is linked to the increased incidence of NAFLD in Sri Lanka.

Strengths of our study include the availability of both parent's anthropometric data and anthropometry being conducted by a single trained investigator while adhering to strict quality control measures, as evidenced by good intra-observer and inter-observer TEM. Limitations of the study include not having a direct measure for abdominal adiposity and using abdominal circumference as a proxy. We were limited to the use of published summary data as raw data was not available for the global comparisons.

Healthy neonates from Colombo Sri Lanka, had significantly higher abdominal adiposity despite being significantly smaller in size than neonates in high-income countries. Female neonates demonstrated higher abdominal adiposity than males with marginally higher abdominal circumference / length ratio despite being significantly smaller at birth. The phenotype of Sri Lankan neonates differed from the thin-fat phenotype described in Indian babies with significantly lower abdominal circumference, implying that Sri Lankan babies are at increased risk of central adiposity compared to Indian babies. Parental weight and BMI correlated with neonatal adiposity in our study population in contrast to family income implying that intergenerational transmission may have a larger effect on neonatal adiposity compared to socioeconomic status.

Further studies are indicated to measure body composition which should be correlated with abdominal circumference which can be used as a screening tool to assess abdominal adiposity from birth. Further studies are also required to establish whether higher abdominal adiposity

as indicated by abdominal circumference at the time of birth, is linked to the increased incidence of metabolic syndrome and NAFLD in Sri Lanka. A conceptual diagram summarising how anthropometry predicts adiposity including the geo-ethnic difference is shown in Figure 1.

ACKNOWLEDGEMENTS

We are indebted to the research assistants M Paranchothy, M Muthuthamby, R Muthuthamby, R Munasinghe, V Mahendralingam, R Balasubramaniam, S Bandara, S Hasarangi, S Maduwanthi, K Tesni, AGP Sathsaranie, AVS Tennekooni, KHC Jayashani, WMPM Wijeratne and SASN Solangarachchi who supported data collection, data management and data analysis, staff of De Soysa Hospital, Colombo, staff of the Faculty of Medicine, University of Colombo and the study participants who contributed to this study in different ways.

CONFLICT OF INTEREST AND FUNDING DISCLOSURE

Conflict of interest

This work was supported by the International Atomic Energy Agency under Doctoral contract [18559].

REFERENCES

1. Winter JD, Langenberg P, Krugman SD. Newborn adiposity by body mass index predicts childhood overweight. *Clin Pediatr (Phila)*. 2010;49:866-70. doi: 10.1177/0009922810369698.
2. Ferreira AP, da Silva Junior JR, Figueiroa JN, Alves JG. Abdominal subcutaneous and visceral fat thickness in newborns: correlation with anthropometric and metabolic profile. *J Perinatol*. 2014; 34:932-35. doi: 10.1038/jp.2014.110.
3. Brambilla P, Bedogni G, Moreno LA, Goran MI, Gutin B, Fox KR, Peters DM, Barbeau P, De Simone M, Pietrobelli A. Cross validation of anthropometry against magnetic resonance imaging for the assessment of visceral and subcutaneous adipose tissue in children. *Int J Obes (Lond)*. 2006;30:23-30. doi: 10.1038/sj.ijo.0803163
4. Hassapidou M, Tzotzas T, Makri E, Pagkalos I, Kaklamanos I, Kapantais E, Abrahamian A, Polymeris A, Tziomalos K. Prevalence and geographic variation of abdominal obesity in 7- and 9-year-old children in Greece; World Health Organization Childhood Obesity Surveillance Initiative 2010. *BMC Public Health*. 2017;17:126. doi: 10.1186/s12889-017-4061-x
5. Eaton-Evans J. Nutritional Assessment: Anthropometry. In: Caballero B, Allen L, Prentice A (eds). *Encyclopedia of Human Nutrition*. 3rd ed. Oxford, UK Academic Press, 2013. pp 227-32. doi: 10.1016/B978-0-12-375083-9.00197-5

6. Mager DR, Yap J, Rodriguez-Dimitrescu C, Mazurak V, Ball G, Gilmour S. Anthropometric measures of visceral and subcutaneous fat are important in the determination of metabolic dysregulation in boys and girls at risk for non-alcoholic fatty liver disease. *Nutr Clin Pract*. 2013;28:101-11. doi: 10.1177/0884533612454884
7. Alexander T, Conlon CA, Gamble G, von Hurst PR, van Dorp L, Ichhpuniani B, Housh R, Ke P, Bloomfield FH. Body composition of New Zealand-born term babies differs by ethnicity, gestational age and sex. *Early Hum Dev* 2019; 140; 104924. Online ahead of print. doi: 10.1016/j.earlhumdev.2019.104924
8. Stanfield KM, Wells JC, Fewtrell MS, Frost C, Leon DA. Differences in body composition between infants of South Asian and European ancestry: the London Mother and Baby Study. *Int J Epidemiol*. 2012; 41:1409-18.
<https://doi.org/10.1093/ije/dys139>
9. Yajnik C, Fall, C, Coyaji K, Hirve SS, Rao S, Barker DJP, Joglekar C, Kellingray S. Neonatal anthropometry: the thin-fat Indian baby. The Pune Maternal Nutrition Study. *Int J Obes*. 2003; 27:173-80. doi: 10.1038/sj.ijo.802219
10. Karamali NS, Ariëns GA, Kanhai HH, de Groot CJ, Tamsma JT, Middelkoop BJ. Thin-fat insulin-resistant phenotype also present in South Asian neonates born in the Netherlands. *J Dev Orig Health Dis*. 2015;6:47-52. doi: 10.1017/S204017441400052X
11. Van Steijn L, Karamali NS, Kanhai HH, Ariens GAM, Fall CHD, Yajnik CS, Middlekoop BJC, Tamsma JT. Neonatal anthropometry: thin-fat phenotype in fourth to fifth generation South Asian neonates in Surinam. *Int J Obes (Lond)*. 2009;33:1326-9. doi: 10.1038/ijo.2009.154
12. Krishnaveni GV, Hill JC, Veena SR, Leary SD, Saperia J, Chachyamma KJ, Karat SC, Fall CHD. Truncal adiposity is present at birth and in early childhood in south Indian children. *Indian Paediatrics* 2005;42:527-38. doi: 10.1007/s13312-018-1244-1
13. Antonisamy B, Raghupathy P, Christopher S, Richard J, Rao PS, Barker DJ, Fall CHD. Cohort Profile: The 1969-73 Vellore birth cohort study in South India. *Int J Epidemiol*. 2009;38:663-9. doi: 10.1093/ije/dyn159
14. Modi N, Thomas EL, Uthaya SN, Umranikar S, Bell JD, Yajnik CS. Whole body magnetic resonance imaging of healthy newborn infants demonstrates increased central adiposity in Asian Indians. *Pediatr Res*. 2009;65:584-7. doi: 10.1203/PDR.0b013e31819d98be
15. Pomeroy E, Mushrif-Tripathy V, Cole TJ, Wells JCK, Stock JT. Ancient origins of low lean mass among South Asians and implications for modern type 2 diabetes susceptibility. *Sci Rep*. 2019;9:10515.
16. Kurpad AV, Varadharajan KS, Aeberli I. The thin-fat phenotype and global metabolic disease risk. *Curr Opin Clin Nutr Metab Care*. 2011;14:542-7. doi: 10.1097/MCO.0b013e32834b6e5e
17. World Health Organization. World health statistics 2015. World Health Organization. 2015. Available at: <https://apps.who.int/iris/handle/10665/170250>

18. de Silva A, Ranasinghe T, Abeykoon P. Universal health coverage and the health Sustainable Development Goal: achievements and challenges for Sri Lanka. *WHO South-East Asia Journal of Public Health*. 2016;5:1-7. doi: 10.4103/2224-3151.206257
19. Department of Census and Statistics. Percentage of households and share of income to total household income by national household income decile and sector – 2016. Sri Lanka Household Income and Expenditure Survey 2016. Ministry of National Policies and Economic Affairs Sri Lanka; 2018; 1: 16-41. Available at: <http://www.statistics.gov.lk/Resource/en/IncomeAndExpenditure/HouseholdIncomeandExpenditureSurvey2016FinalResults.pdf>
20. Brandt JM, Allen GA, Butler MG. Normative standards and patterning of fat and muscle in white and black newborn infants. *Dysmorphol Clin Genet*. 1991;5:8896. doi: 10.1177/000992286600500102
21. Pomeroy E, Stock JT, Cole TJ, O'Callaghan M, Wells JCK. Relationships between neonatal weight, limb lengths, skinfold thicknesses, body breadths and circumferences in an Australian cohort. *PLoS One*. 2014;9:e105108. doi: 10.1038/s41598-019-46960-9
22. de Onis M, Onyango AW, den Broeck JV, Chumlea WC, Martorell R. The WHO Multicentre Growth Reference Study (MGRS): Measurement and standardization protocols for anthropometry used in the construction of a new international growth reference. *Food Nutr Bull*. 2004;25(Suppl 1): S27-36. doi: 10.1177/15648265040251S105
23. NHS. Procedure for making circumference measurement of infants. Standard Operating Procedure from NIHR Southampton Biomedical Research Centre. 2015. [accessed 2020/10/10]; Available from: <http://www.uhs.nhs.uk/Media/Southampton-Clinical-Research/Procedures/BRCProcedures/Procedure-for-circumference-measurements-of-children.pdf>.
24. Cheikh Ismail L, Knight H, Ohuma E, Hoch L, Chumlea WC. Anthropometric standardisation and quality control protocols for the construction of new, international, fetal and newborn growth standards: the INTERGROWTH-21st Project. *BJOG*. 2013;120:4855. doi: 10.1111/1471-0528.12127
25. Perrini TA, de Oliveira GL, Ornellas JS, Oliveira FP. Technical error of measurement in anthropometry. *Rev Bras Med Esporte*. 2005;11:86-90. English version available at https://www.scielo.br/pdf/rbme/v11n1/en_24109.pdf (accessed 2021/02/18).
26. Jaswant S, Nitish M. Use of upper-arm anthropometry as measure of body-composition and nutritional assessment in children and adolescents (6-20 years) of Assam, Northeast India. *Ethiop J Health Sci*. 2014;24:243-52. doi: 10.4314/ejhs.v24i3.8
27. Rolland-Cachera MF, Brambilla P, Manzoni P, Akrouf M, Sironi S, Del Maschio A, Chiumello G. Body composition assessed on the basis of arm circumference and triceps skinfold thickness: a new index validated in children by magnetic resonance imaging. *Am J Clin Nutr*. 1997;65:1709-13. doi: 10.1093/ajcn/65.6.1709
28. Gurney JM, Jelliffe DB. Arm anthropometry in nutritional assessment: Nomogram for rapid calculation of muscle circumference and cross-sectional muscle and fat areas. *Am J Clin Nutr*. 1973; 26:912-5. doi: 10.1093/ajcn/26.9.912

29. Frisancho AR. New norms of upper limb fat and muscle areas for assessment of nutritional status. *Am J Clin Nutr.* 1981;34:2540-5. doi: 10.1093/ajcn/34.11.2540
30. Anand SS, Gupta MK, Schulze KM, Desai D, Abdalla N, Wahi G et al. What accounts for ethnic differences in newborn skinfold thickness comparing South Asians and White Caucasians? Findings from the START and FAMILY Birth Cohorts. *Int J Obes (Lond).* 2016;40:239-44. doi: 10.1038/ijo.2015.171
31. Shields BM, Knight BA, Powell RJ, Hattersley AT, Wright DE. Assessing newborn body composition using principal components analysis: differences in the determinants of fat and skeletal size. *BMC Pediatrics.* 2006;6:24. doi: 10.1186/1471-2431-6-24
32. Gale C, Logan KM, Jeffries S, Parkinson JRC, Santhakumaran S, Uthaya S, Durighel G, Alavi A, Thomas EL, Bell JD, Modi N. Sexual dimorphism in relation to adipose tissue and intrahepatocellular lipid deposition in early infancy. *Int J Obes.* 2015;39:629-32. doi: 10.1038/ijo.2015.4
33. Bandara T, Hettiarachchi M, Liyanage C, Amarasena S. Body composition among Sri Lankan infants by O dilution method and the validity of anthropometric equations to predict body fat against O dilution. *BMC Pediatrics.* 2015;15:52. doi: 10.1186/s12887-015-0371-2
34. Patro B, Liber A, Zalewski B, Poston L, Szajewska H, Koletzko B. Maternal and paternal body mass index and offspring obesity: a systematic review. *Ann Nutr Metab.* 2013;63:32-41. doi: 10.1159/000350313
35. Diaz EC, Cleves MA, DiCarlo M, Sobik SR, Ruebel ML, Thakali KM et al. Parental adiposity differentially associates with newborn body composition. *Pediatr Obes.* 2020;15:e12596. doi: 10.1111/ijpo.12596
36. Jääskeläinen A, Pussinen J, Nuutinen O, Schwab U, Pirkola J, Kolehmainen M, Jarvelin MR, Laitinen J. Intergenerational transmission of overweight among Finnish adolescents and their parents: a 16-year follow-up study. *Int J Obes.* 2011;35:1289-94. doi: 10.1038/ijo.2011.150
37. Ornellas F, Carapeto P, Mandarim-de-Lacerda CA, Marcia BA. Obese fathers lead to an altered metabolism and obesity in their children in adulthood: review of experimental and human studies. *Jornal de Pediatria (Versão em Português).* 2017;93:551-9. doi: 10.1016/j.jpdp.2017.08.006
38. Catalano PM, Farrell K, Thomas A, Huston-Presley L, Mencin P, de Mouzon SH, Amini SB. Perinatal risk factors for childhood obesity and metabolic dysregulation. *Am J Clin Nutr.* 2009;90:1303-13. doi: 10.3945/ajcn.2008.27416
39. Fuentes ED, Cleves M, Sims C, Shankar K, Borsheim E, Andres A. Parental obesity differentially associates with newborn adiposity. Is programming of the GH-IGF axis involved? *Curr Dev Nutr.* 2019; 3(Suppl 1):nzz041. doi: 10.1093/cdn/nzz041.OR09-05-19
40. Paley C, Hull H, Ji Y, Toro-Ramos T, Thornton T, Bauer J, Matthews P, Yu A, Navder K, Dorsey K, Gallagher D. Body fat differences by self-reported race/ethnicity in healthy term newborns. *Pediatr Obes.* 2016;11:361-8. doi: 10.1111/ijpo.12072

41. Tint MT, Fortier MV, Godfrey KM, Shuter B, Kapur J, Rajadurai VS et al. Abdominal adipose tissue compartments vary with ethnicity in Asian neonates: Growing Up in Singapore Toward Healthy Outcomes birth cohort study. *Am J Clin Nutr.* 2016;103:1311-7. doi: 10.3945/ajcn.115.108738
42. Ayonrinde OT, Olynyk JK, Marsh JA, Beilin LJ, Mori TA, Oddy WH, Adams LA. Childhood adiposity trajectories and risk of nonalcoholic fatty liver disease in adolescents. *J Gastroenterol Hepatol.* 2015;30:163-71. doi: 10.1111/jgh.12666
43. Amarapurkar D, Kamani P, Patel N, Gupte P, Kumar P, Agal S, Baijal R, Lala S, Chaudhary D, Deshpande A. Prevalence of non-alcoholic fatty liver disease: population based study. *Ann Hepatol.* 2007;6:161-3. doi: 10.1016/S1665-2681(19)31922-2
44. Dassanayake AS. Nonalcoholic fatty liver disease: Identifying the disease burden in Sri Lanka. *Euroasian J Hepatogastroenterol.* 2018;8:69-72. doi: 10.5005/jp-journals-10018-1263.

Table 1. Socio demographic data of our study population

Characteristic	N=337
Maternal Age (years) (mean±SD, range)	29±6 (19-44)
Maternal education	
Years of formal education (mean ± SD, range)	11±3 (0-17)
School not attended	4 (1.1%)
Primary	12 (3.6%)
Secondary	302 (89.7%)
University	6 (1.7%)
Professional	13 (3.9%)
Maternal occupation	
Housewife	208 (61.7%)
Skilled manual work	82 (24.4%)
Unskilled manual work	7 (2.2%)
Managerial	31 (9.2%)
Clerical support	9 (2.5%)
Maternal BMI (kg/m ²) (mean ± SD, range)	24.5± 5.1 (12.1-39.9)
Paternal age (years) (mean ± SD, range)	32±6 (19-52)
Paternal education	
Years of formal education (mean ± SD, range)	11±2 (0-17)
School not attended	5 (1.4%)
Primary	57 (17.0%)
Secondary	260 (77.3%)
University	7 (2.0%)
Professional	8 (2.3%)
Paternal occupation	
Housework	1 (0.3%)
Student	2 (0.6%)
Skilled manual work	195 (57.8%)
Unskilled manual work	2 (0.6%)
Managerial	85 (25.2%)
Clerical support	52 (15.5%)
Paternal BMI (kg/m ²) (mean ± SD, range)	24.6± 5.9 (15.2-44.1)
Marital status	
Married	334 (99%)
Unmarried	3(1%)
Parity (mean ± SD, range)	2±1 (1-6)
Monthly family income (SLR) (1 USD=150SLR)	
Median (Inter quartile range)	30,000 (25,000-45,000)
1 st quintile (Less than 23,518)	75 (22.4%)
2 nd quintile (23,519 – 36,445)	144 (42.7%)
3 rd quintile (36,446 – 51,862)	65 (19.3%)
4 th quintile (51,863 – 81,371)	33 (9.7%)
5 th quintile (81,371 upwards)	20 (5.9%)

†Japanese, Thai, Korean, Philippine.

Table 2. Comparison of birth anthropometry between males and females

Parameter	Male (n=174)	Female (n=163)	p value
Weight (kg)	2.98±0.5	2.84±0.4	0.004
Length (cm)	49.0±2.4	47.9±2.1	<0.001
HC (cm)	34.1±1.3	33.5±1.2	<0.001
CC (cm)	32.7±2.6	32.1±1.9	0.010
Body Mass Index (BMI)(kg/m ²)	12.4±1.2	12.3±1.5	0.985
Ponderal index (PI) (kg/m ³)	25.2±2.5	25.8±3.6	0.083
<i>Fat mass</i>			
AC (cm)	30.8±2.4	30.5±2.1	0.26
AC/length	0.63±0.0	0.64±0.0	0.051
UFA (cm ²)	2.4±0.7	2.3±0.5	0.319
AFI	26.0±4.5	26.1±3.7	0.804
UFE (cm ²)	2.6±0.8	2.5±0.6	0.321
<i>Skinfolds</i>			
Biceps (mm)	4.4±0.9	4.4±0.9	0.682
Triceps (mm)	4.8±1.1	4.7±0.9	0.577
Subscapular (mm)	5.1±1.4	5.0±1.2	0.697
Supra-iliac (mm)	4.2±1.1	4.2±1.0	0.958
Sum of skinfolds (mm)	18.5±3.8	18.4±3.4	0.866
<i>Muscle mass</i>			
MUAC (cm)	10.7±0.9	10.5±0.9	0.17
MAMC (cm)	9.2±0.8	9.1±0.8	0.163
UMA (cm ²)	6.8±1.2	6.6±1.1	0.165
UME (cm ²)	6.6±1.1	6.4±1.1	0.179

All values are given as mean and standard deviations.

HC: head circumference; CC: chest circumference; AC: abdominal circumference; UFA: upper arm fat area; AFI: arm fat index; UFE: upper arm fat estimate; MUAC: mid upper arm circumference; MAMC: mid arm muscle circumference; UMA: upper arm muscle area; UME: upper arm muscle estimate.

Table 3. Comparison of anthropometric parameters and its derivatives of our study with other published studies

	SL M=174 F=163	USA.C M=40 F=40	USA.Afri M=30 F=30	Canada.C A=389	Canada.A A=400	Pune A=631	UK A=338	Surinam A=39	Australia M=668 F=602
W (kg)									
A	2.91±0.4	3.3±0.4 (<i>p</i> <0.001)		3.52±0.02 (<i>p</i> <0.001)	3.28±0.02 (<i>p</i> <0.001)	2.66±0.35 (<i>p</i> <0.001)	3.49±0.48 (<i>p</i> <0.001)	3.16±0.5 (<i>p</i> =0.004)	
M	2.98±0.5	3.35±0.4	3.31±0.4						3.50±0.4 (<i>p</i> <0.001)
F	2.84±0.4	3.40±0.5	3.28±0.5						3.38±0.4 (<i>p</i> <0.001)
L (cm)									
A	48.5±2.3	50±1.7 (<i>p</i> <0.001)		50.2±0.1 (<i>p</i> <0.001)	52.1±0.1 (<i>p</i> <0.001)	47.7±2 (<i>p</i> <0.001)	49.8±1.9 (<i>p</i> <0.001)	48.7±2.1 (<i>p</i> =0.579)	
M	49.0±2.4	51.2±2.7	50.0±1.7						
F	47.9±2.1	51.2±2.3	50.1±2.1						
HC (cm)									
A	33.8±1.3			34.9±0.1 (<i>p</i> <0.001)	34.1±0.1 (<i>p</i> <0.001)	31±1.2 (<i>p</i> <0.001)	35.2±1.3 (<i>p</i> <0.001)	34.3±1.3 (<i>p</i> =0.027)	
M	34.1±1.3								35.4±1.3 (<i>p</i> <0.001)
F	33.5±1.2								34.8±1.1 (<i>p</i> <0.001)
PI kg/m ³									
A	25.5 ±3			29.9 ±0.2 (<i>p</i> <0.001)	23.3±0.2 (<i>p</i> <0.001)	24.5±2.5 (<i>p</i> <0.001)	28.2±2.3 (<i>p</i> <0.001)	27.2±2.6 (<i>p</i> <0.001)	
M	25±3								
F	26±3								
MUAC (cm)									
M	10.7±0.9	10.0±0.8 (<i>p</i> <0.001)	10.2±0.9 (<i>p</i> =0.006)			9.7±0.9 (<i>p</i> <0.001)	11.5±1 (<i>p</i> <0.001)	10.6±1.2	10.7±0.9 (<i>p</i> <0.001)
F	10.5±0.9	10±0.8 (<i>p</i> =0.001)	10.1±1.0 (<i>p</i> =0.038)						10.9±0.9 (<i>p</i> <0.001)
Abd (cm)									
A	30.6±2.3					28.6±1.9 (<i>p</i> <0.001)		31.1±2.4 (<i>p</i> =0.221)	
M	30.8±2.4								28.8±1.9 (<i>p</i> <0.001)
F	30.5±2.1								28.3±1.98 (<i>p</i> <0.001)

W: weight; L: length, HC: head circumference; PI: ponderal index; MUAC: mid-upper arm circumference; Abd: abdominal circumference; B: biceps skinfold thickness; T: triceps skinfold thickness; SUB: subscapular skinfold thickness; SI: suprailiac skinfold thickness; UFA: upper arm fat area; SL: Sri Lanka, USA: United States of America; C: Caucasian; Afri: African; A: Asian; UK: United Kingdom; Pune: Indian; A: all; M: male; F: female.
Significance shown in comparison to the Sri Lankan data via independent sample, unequal variance, T test.

Not Proof Read

Table 3. Comparison of anthropometric parameters and its derivatives of our study with other published studies

	SL M=174 F=163	USA.C M=40 F=40	USA.Afri M=30 F=30	Canada.C A=389	Canada.A A=400	Pune A=631	UK A=338	Surinam A=39	Australia M=668 F=602
Waist (cm)									
A				29.9±2.1	31.1±0.1				
B (mm)									
M	4.4±1.0	3.1±0.8 (<i>p</i> <0.001)	3.1±0.5 (<i>p</i> <0.001)						
F	4.4±1.0	3.2±0.7 (<i>p</i> <0.001)	3.1±0.7 (<i>p</i> <0.001)						
T (mm)									
A	4.8±1.0			5.4±0.1 (<i>p</i> <0.001)	6.2±0.1 (<i>p</i> <0.001)	4.2 (<i>p</i> <0.001)		4.6 (<i>p</i> <0.001)	
M	4.8±1.9	3.5±0.6 (<i>p</i> <0.001)	3.7±0.6 (<i>p</i> <0.001)						4.9±0.9 (<i>p</i> <0.001)
F	4.7±0.9	3.8±0.9 (<i>p</i> <0.001)	3.7±0.9 (<i>p</i> <0.001)						5.0±0.9 (<i>p</i> <0.001)
SUB (mm)									
A	5.1±1.3			5.2±0.1 (<i>p</i> =0.144)	5.6±0.1 (<i>p</i> =0.144)	4.2 (<i>p</i> <0.001)	4.6 (<i>p</i> <0.001)	4.8 (<i>p</i> <0.001)	
M	5.1±1.4	4±0.9 (<i>p</i> <0.001)	4.5±0.8 (<i>p</i> =0.002)						5.3±1.0 (<i>p</i> <0.001)
F	5.0±1.2	4.7±0.9 (F) (<i>p</i> =0.044)	4.9±1.4 <i>p</i> =0.574						5.5±1.4 (<i>p</i> <0.001)
SI (mm)									
M	4.2±1.0	3.6±0.6 (<i>p</i> <0.001)	3.6±0.9 <i>p</i> =0.002						
F	4.2±1.0	4.0±0.8 (<i>p</i> =0.097)	4.0±1.4 <i>p</i> =0.638						
T+ SUB (mm)									
A	9.8±2.0			10.6±0.1 (<i>p</i> <0.001)	11.7±0.1 (<i>p</i> <0.001)				
UFA (cm ²)									
M	240±67	166±46 (<i>p</i> <0.001)	180±37.7 <i>p</i> <0.001						
F	233±54	181±48 (<i>p</i> <0.001)	177±51.5 (<i>p</i> <0.001)						

W: weight; L: length; HC: head circumference; PI: ponderal index; MUAC: mid-upper arm circumference; Abd: abdominal circumference; B: biceps skinfold thickness; T: triceps skinfold thickness; SUB: subscapular skinfold thickness; SI: suprailiac skinfold thickness; UFA: upper arm fat area; SL: Sri Lanka; USA: United States of America; C: Caucasian; Afri: African; A: Asian; UK: United Kingdom; Pune: Indian; A: all; M: male; F: female
Significance shown in comparison to the Sri Lankan data via independent sample, unequal variance, T test.

Not Proof Read




Figure 1. Conceptual diagram of how anthropometry can predict adiposity and the geo-ethnic difference.