

This author's PDF version corresponds to the article as it appeared upon acceptance. Fully formatted PDF versions will be made available soon.

## **Malnutrition and its risk factors in a home for Seniors in Shanghai**

doi: 10.6133/apjcn.202211/PP.0011

Published online: November 2022

**Running title:** Malnutrition risk among Shanghainese seniors

Jianqin Sun MD<sup>1†</sup>, Wuke Yuan BS<sup>2†</sup>, Min Chen MS<sup>1</sup>, Yanqiu Chen MD<sup>1</sup>, Danfeng Xu PhD<sup>1</sup>, Zhen Li BS<sup>1</sup>, Huijing Bai PhD<sup>1</sup>, Qi Xu PhD<sup>2</sup>, Yuanrong Jiang PhD<sup>3</sup>, Jie Gu PhD<sup>3</sup>, Shengqi Li PhD<sup>3</sup>, Chenxi Su MS<sup>3</sup>, Lili Gu BS<sup>4</sup>, Jiaxin Fang BS<sup>4</sup>, Xinyao Zhu BS<sup>4</sup>

<sup>1</sup>Clinical Nutrition Center, Fudan University Affiliated to Hua Dong Hospital, Shanghai, China

<sup>2</sup>School of Public Health, Shanghai University of Traditional Chinese Medicine, Shanghai, China

<sup>3</sup>Wilmar (Shanghai) Biotechnology Research & Development Center Co., Ltd., Shanghai, China

<sup>4</sup>Shanghai XiJiao Union Retirement Center, Shanghai, China

<sup>†</sup>Both authors contributed equally to this manuscript

**Authors' email addresses and contributions:**

JS: jianqins@163.com

Contribution: conception and design, doing the field, data collection, data analysis and interpretation, preparation of draft manuscript, doing revisions or providing critique, and overall and sectional scientific management.

WY: 15078800806@163.com

Contribution: doing the field, data collection, data analysis and interpretation, preparation of draft manuscript, and doing revisions.

MC: chenmin1892@hotmail.com

Contribution: conception and design, doing the field, and data collection.

YC: yqchen0808@126.com

Contribution: conception and design, doing the field, and data collection.

DX: danfeng.xu@qq.com

Contribution: conception and design, doing the field, and data collection.

ZL: 1647914519@qq.com

Contribution: conception and design, doing the field, and data collection.

HB: 109642976@qq.com

Contribution: conception and design, doing the field, and data collection.

QX: isuxuqi@163.com

Contribution: preparation of draft manuscript, and revised and critiqued the draft manuscript.

YJ: jiangyuanrong@cn.wilmar-intl.com

Contribution: data collection, and overall and sectional scientific management.

JG: gujie@cn.wilmar-intl.com

Contribution: data collection, revised and critiqued the draft manuscript, and overall and sectional scientific management.

SL: shengqili@outlook.com

Contribution: data collection, and overall and sectional scientific management.

CS: 568106486@qq.com

Contribution: doing the field, and data collection.

LG: 157135937@qq.com

Contribution: overall and sectional scientific management.

JF: 642584915@qq.com

Contribution: overall and sectional scientific management.

XZ: 415271986@qq.com

Contribution: doing the field, data collection, and overall and sectional scientific management.

**Corresponding Author:** Dr Jianqin Sun, Huadong Hospital, Shanghai Medical College, Fudan University, West Yan-An Road 221#, Shanghai 200040, China. Tel: 021-62483180. Fax: 021-62483696. Email: jianqins@163.com

Not Proof Read

## ABSTRACT

**Background and Objectives:** Older adults residing in senior homes are at a high risk of malnutrition. In this study, we investigated the nutritional status of these individuals and factors associated with malnutrition in this population. **Methods and Study Design:** This cross-sectional study (September 2020–January 2021) included a total of 583 older adults residing in a senior home in Shanghai (mean age, 85.0±6.6 years). The Mini Nutritional Assessment Short Form (MNA-SF) questionnaire was administered to assess the nutritional status of the participants. Patients with possible sarcopenia were identified according to the guidelines recommended by the Asian Working Group for Sarcopenia in its 2019 consensus (AWGS 2019). Moreover, the factors influencing malnutrition were determined through multivariate analyses. **Results:** The likelihoods of having malnutrition and being at a risk of malnutrition were noted in 10.5% and 37.4% of the participants, respectively. In both male and female participants, handgrip strength (HGS) and calf circumference (CC) increased significantly with increasing scores on the aforementioned questionnaire ( $p<0.001$ ). Among the participants, 44.6% had  $\geq 3$  chronic diseases and 48.2% used multiple medicines. Multivariate analyses revealed that dysphagia (OR, 3.8; 95% CI, 1.7–8.5), possible sarcopenia (OR, 3.6; 95% CI, 2.2–5.6), and dementia (OR, 4.5; 95% CI, 2.8–7.0) were correlated with a relatively high prevalence of malnutrition/malnutrition risk. Exercise (at least thrice a week) reduced malnutrition risk. **Conclusions:** Malnutrition is common among older adults residing in senior homes; therefore, the associated factors must be identified, and appropriate interventions should be administered.

**Key Words:** nutritional status, chronic disease, senior home, older adult, malnutrition

## INTRODUCTION

Adequate nutrition is essential for healthy aging. Malnutrition is associated with numerous adverse outcomes, such as frailty, decreased immunity, and increased chronic disease and mortality risks. The risk of malnutrition increases with increasing age and decreasing functional ability.<sup>1</sup> In older adults, appetite loss is common and can be attributed to underlying acute or chronic medical conditions or adverse effects of medications.<sup>2</sup> Moreover, taste and smell deficits may adversely affect food choice and intake in older adults. During the coronavirus disease 2019 pandemic, malnutrition emerged as a major concern, which can be attributed to individuals' inactivity and difficulty in obtaining fresh food items because of social distancing and lockdown.<sup>3</sup> The Seventh National Census of China revealed that

Chinese individuals aged  $\geq 60$  years accounted for 18.7% of the total population in 2020; the corresponding value was 23.4% in Shanghai.<sup>4</sup> The trend toward an increasingly older population will inevitably lead to increased incidence rates of not only chronic diseases but also disability and malnutrition due to multitudinous etiologies.

Older adults residing in senior homes constitute a unique population. The nutritional status of these individuals may be affected by various medical conditions, including mobility issues and systemic and chronic health conditions.<sup>5-7</sup> Consequently, the risk of malnutrition in older adults residing in senior homes is higher than that in the general population. Active nutritional surveys help prevent the deterioration of nutritional status and diseases in older adults; hence, such surveys are essential for promoting healthy aging and achieving the national goal of “Healthy China 2030.”<sup>8</sup> The Seventh National Census of China indicated that the aging population in Shanghai has been gradually increasing in number; accordingly, the number of older adults residing in senior homes has been steadily increasing.<sup>4</sup>

The Mini Nutritional Assessment Short Form (MNA-SF) is a valid nutritional screening tool for detecting malnutrition and malnutrition risk in older adults.<sup>9</sup> To the best of our knowledge, few studies have focused on the status of malnutrition in Chinese older adults residing in senior homes and the factors associated with malnutrition in this population; furthermore, only a few surveys have been conducted using the MNA-SF questionnaire to detect malnutrition and its risk in older adults. Thus, in the present study, we investigated the prevalence of malnutrition in older adults residing in senior homes in Shanghai as well as the factors associated with malnutrition in these individuals.

## **MATERIALS AND METHODS**

### ***Participants***

This cross-sectional study was conducted between September 2020 and January 2021 and included all residents of a senior home in Changning, Shanghai (Figure 1). The only inclusion criterion was an age of  $\geq 60$  years. The study protocol was approved by the Research Ethics Committee of Huadong Hospital affiliated to Fudan University (approval number: 2020K178). All participants provided written informed consent for participation.

### ***General data collection***

Relevant data were collected through questionnaire surveys and face-to-face interviews. These data included the participants’ demographics, exercise frequency (per week), diet type, nutritional supplement use, nursing care requirement, medical history, and medication use.

Diet was divided into the following two categories: general and nongeneral. Nutritional supplements included protein powders, mineral and vitamin supplements, and oral nutritional supplements.

The care requirement of the participants was assessed by professionals on a semiannual basis in accordance with the DB31/T684-2013 Standard issued by the Shanghai Municipal Bureau of Quality and Technical Supervision (2013).<sup>10</sup> The assessment included the evaluation of various dimensions of self-care ability, cognitive ability, emotion, and vision. On the basis of the assessment results, the nursing care requirements of older adults were classified into four levels. Higher levels indicated the individuals' poorer ability to live independently. Levels 3 and 4 indicated the need for high levels of nursing care.

Medical history included the history of tumors, stroke, dysphagia, Alzheimer's disease, osteoporosis, chronic bronchitis, lumbar disease, Parkinson's disease, fractures, chronic obstructive pulmonary disorder, and renal insufficiency. Taking  $\geq 5$  medications per day was regarded as polypharmacy.<sup>11</sup>

### ***Nutritional status***

Nutritional status was assessed using the MNA-SF questionnaire.<sup>9</sup> The questionnaire provides information on appetite and food intake, weight loss, mobility, psychological stress or acute disease in the past 3 months, mental status, and body mass index (calf circumference [CC] is used if height and weight cannot be measured). The total score is 14; scores of  $\geq 12$ , 8–11, and  $< 8$  indicate normal nutrition, malnutrition risk, and malnutrition, respectively.

### ***Sarcopenia screening***

The concept of possible sarcopenia in Asian populations was introduced by the Asian Working Group for Sarcopenia in its 2019 consensus (AWGS 2019). This serves as a basis for the practical and feasible diagnosis of sarcopenia.<sup>12</sup> Decreased muscle strength is a key symptom of sarcopenia; it is central to sarcopenia diagnosis and helps predict adverse outcomes.<sup>13</sup> Using the AWGS 2019 data, individuals with possible sarcopenia were identified as follows: screening in addition to CC measurements and the simple five-item questionnaire (SARC-F) or SARC-F combined with CC (SARC-CalF) questionnaire during the case finding stage, followed by the assessment of handgrip strength (HGS) in the screen-positive population to identify patients with possible sarcopenia.

The circumference of the widest area of the calf was measured using a nonelastic tape with the participants in the sitting position and their knee flexed at 90°. The SARC-F questionnaire

is used for the rapid screening of sarcopenia. The scores on this questionnaire range from 0 to 10; scores of  $\geq 4$  indicate an abnormal status.<sup>14</sup> SARC-F involves the following five items: lifting and carrying a 10-lb weight, walking across a room, getting up from a bed or chair, climbing 10 stair steps, and the number of falls in the previous year. The SARC-CalF questionnaire incorporates CC into the SARC-F questionnaire; the scores on SARC-CalF range from 0 to 20; scores of  $\geq 11$  indicate an abnormal status.<sup>15</sup>

HGS was measured using a digital hand dynamometer (CAMRY EH101, Guangdong, China).<sup>16</sup> The participants were in the standing position with their arms drooping naturally and elbows extended, and they were instructed to use their dominant hand to squeeze the handle of the dynamometer using their maximum strength. The test was performed at least twice, and the higher of the two readings was used in analysis. If the participants could not stand independently, they could sit during the test.

As per the recommendations of the AWGS 2019 consensus, sarcopenia is indicated by a CC of  $< 34$  cm for men, CC of  $< 33$  cm for women, SARC-F score of  $\geq 4$ , or SARC-CalF score of  $\geq 11$ .<sup>12</sup> Low muscle strength is indicated by an HGS of  $< 28$  kg for men and  $< 18$  kg for women.<sup>12</sup>

### ***Statistical analysis***

Statistical analyses were performed using SPSS Statistics (version 26.0; IBM Corporation, Armonk, NY, USA). Data are presented in terms of mean  $\pm$  standard deviation or number and percentage values. Continuous variables were analyzed using Student's t or Mann–Whitney U tests, whereas categorical variables were analyzed using Pearson's chi-square or Fisher's exact tests. The correlations between variables were analyzed using Spearman's correlation coefficients. A two-tailed  $p$  value of  $< 0.05$  indicated statistical significance. The factors associated with malnutrition were assessed using the OR for binary logistic regression and 95% CI. Variables with a  $p$  value of  $< 0.1$  in univariate analysis (Pearson's chi-square test) were included in a binary logistic regression model (forward layer recurrent). The *Hosmer–Lemeshow* (H–L) test was used to evaluate the goodness of fit of the binary logistic regression model.

## **RESULTS**

### ***Nutritional status and chronic diseases***

This study included a total of 583 older adults (mean age,  $85.96 \pm 6.58$  years); among them, 204 (35.0%) were men (mean age,  $85.25 \pm 7.81$  years) and 333 (65.0%) were women (mean

age,  $86.35 \pm 5.79$  years). The mean MNA-SF scores of all participants was  $11.1 \pm 2.4$  (Table 1). The difference between the two sexes in terms of the prevalence of malnutrition was nonsignificant ( $p > 0.05$ ). On the basis of the participants' scores on the MNA-SF questionnaire, 61 residents (10.5%) were regarded as malnourished individuals, 218 (37.4%) were identified to be at a risk of malnutrition, and 304 (52.1%) were regarded as well-nourished individuals (Figure 2).

Among the diseases surveyed, the top five chronic diseases with the highest prevalence were hypertension (69.6%), possible sarcopenia (65.9%), stroke (46.5%), coronary heart disease (40.3%), and dementia (31.6%). Of the participants, 90.7% had  $\geq 2$  chronic diseases, 44.6% had  $\geq 3$  chronic diseases, and 48.2% used multiple medicines. Significant differences were noted between the sexes in terms of the prevalence of osteoporosis and coronary heart disease ( $p < 0.05$ ).

### ***Risk factors for malnutrition***

Table 2 summarizes the results of the univariate analysis. Significant differences were observed between participants with malnutrition or those at malnutrition risk and participants with normal nutritional status in terms of nursing care requirement, exercise frequency, diet type, protein powder use, mineral and vitamin supplement use, comorbidities ( $n \geq 3$ ), and polypharmacy ( $p < 0.05$ ). The prevalence of dysphagia, stroke, dementia, fracture history, and possible sarcopenia was higher in participants with malnutrition and those at malnutrition risk than in participants with normal nutritional status ( $p < 0.05$ ).

The correlations of nutrition status with HGS and CC were analyzed using Spearman's correlation coefficients (Figure 3). Significant positive correlations were noted between HGS and MNA-SF scores in both men ( $R_s = 0.54$ ;  $p < 0.001$ ) and women ( $R_s = 0.45$ ;  $p < 0.001$ ). Similarly, significant positive correlations were observed between CC and MNA-SF scores in both men ( $R_s = 0.56$ ;  $p < 0.001$ ) and women ( $R_s = 0.47$ ;  $p < 0.001$ ). Both HGS and CC increased significantly with increasing MNA-SF scores in both men and women ( $p < 0.001$ ).

### ***Independent factors influencing malnutrition risk***

Variables with a  $p$  value of  $< 0.10$  in the univariate analysis and age were included in the multivariate analysis model. The  $p$  value (0.582) for the multivariate analysis model was obtained using the H-L test. Table 3 presents the correlation between the prevalence of malnutrition or malnutrition risk and dysphagia (OR=3.8; 95% CI=1.7–8.5). Possible sarcopenia exacerbated malnutrition. The likelihood of having malnutrition or being

malnutrition risk (OR=3.6; 95% CI=2.2–5.6) was higher in participants with possible sarcopenia than in non-sarcopenia. Furthermore, dementia was identified to be a risk factor for having malnutrition or being at malnutrition risk (OR=4.5; 95% CI=2.8–7.0). Participants who exercised at least thrice a week exhibited a lower likelihood of having malnutrition or being at malnutrition risk (OR=0.3; 95% CI=0.2–0.5) than did those who exercised <3 times a week.

## DISCUSSION

The residents of senior homes are older and have poor health. Malnutrition is a major concern for these older adults. However, the symptoms of malnutrition in these individuals are often overshadowed by aging-related health problems. Thus, relevant diagnosis and treatment are only possible when their nutritional status and disease deteriorate. Active nutritional surveys and interventions may help prevent and treat malnutrition and improve the health of older adults. MNA-SF scores reflect the nutritional status of individuals; the MNA-SF scores of Chinese older adults are positively correlated with their health-related quality of life.<sup>17</sup> In the present study, 10.5% of the participants had malnutrition and 37.4% of the participants were at a risk of malnutrition. Our findings are consistent with those reported by a cross-sectional study conducted in Shanghai, in which the MNA-SF tool was used to survey a total of 600 older adults (aged  $\geq 65$  years) residing in long term care institutions; the findings of the earlier study revealed that 43.7% of the participants were at a risk of malnutrition and 10.7% of the participants had malnutrition.<sup>18</sup> In the present study, 90.7% and 44.6% of the participants had  $\geq 2$  and  $\geq 3$  chronic diseases, respectively. These findings corroborate those of a cross-sectional study conducted in Nanjing, in which 90.3% of a total of 918 participants (mean age, 86.2 years) had  $\geq 2$  chronic diseases.<sup>19</sup>

Adequate nutritional intake is necessary to ensure a good nutritional status. Dysphagia impairs individuals' ability to eat and drink, thus reducing the dietary intake of energy, water, and other nutrients, ultimately causing malnutrition (if corrective measures are not adopted).<sup>20</sup> In the present study, dysphagia increased the likelihood of having malnutrition or being at a risk of malnutrition by 3.8 times (OR=1.8; 95% CI=1.7–8.5). In a cross-sectional study conducted in Finland, the MNA questionnaire was used to survey a total of 2,114 nursing home residents (mean age, 82 years); dysphagia was identified to be an independent risk factor for malnutrition (OR=3.0; 95% CI=2.1–4.4). This finding is consistent with that of our study.<sup>21</sup> Cognitive changes affect the nutritional status of individuals. According to the European Society for Clinical Nutrition and Metabolism, the nutritional status and weight of

patients with dementia decline with disease progression; this decline may be associated with various factors, such as appetite loss; olfactory and taste dysfunction; inflammatory response; depression; medications' adverse effects; and reduced ability to remember to eat, recognize food, and eat independently. In the present study, patients with dementia (OR=4.5; 95% CI=2.8–7.0) exhibited a higher likelihood of having malnutrition or being at a risk of malnutrition than did those without dementia. A meta-analysis that included six longitudinal studies to explore the factors associated with malnutrition in older adults identified dementia to be a risk factor for malnutrition (OR=2.1;  $p<0.001$ ).<sup>22</sup> The effects of dementia on the likelihood of having malnutrition or being at risk of malnutrition were stronger in our study than in the aforementioned study; this difference was probably because the two studies differed in terms of design and participants.

Physical activity is essential for improved energy metabolism, appetite, bowel movement, and nutritional status. In the present study, participants who exercised at least thrice a week exhibited a lower likelihood of having malnutrition or being at a risk of malnutrition (OR=0.3; 95% CI=0.2–0.5) than did those who exercised <3 times a week. The positive effects of increased physical activity on the nutritional status of older adults was also demonstrated in a randomized controlled trial, in which a total of 55 older adults (mean age, 82 years) were divided into two groups (SGB group, 6 weeks of group exercise + 6 weeks of home exercise, and LGB group, 12 weeks of group exercise); the findings revealed that exercise helped improved the participants' MNA scores (baseline vs posttest: SGB group:  $22.1\pm 2.6$  vs.  $25\pm 3.6$ ,  $p<0.001$ ; LGB group:  $21.1\pm 3.7$  vs  $23.9\pm 2.5$ ,  $p<0.01$ ).<sup>23</sup> In the present study, possible sarcopenia was found to increase the risk of malnutrition (OR=3.6; 95% CI=2.2–5.6). Skeletal muscles are crucial for physical movement, posture, and breathing. A less common but pivotal role of these muscles involves the metabolism of energy, glucose, and protein.<sup>24</sup> Low amounts of skeletal muscle mass is a major factor assessed using the MNA-SF questionnaire and the Global Leadership Initiative on Malnutrition framework to diagnose malnutrition.<sup>9,25</sup> Therefore, sarcopenia is a key risk factor for the loss of skeletal muscles, which leads to malnutrition. We noted significantly positive correlations of HGS and CC with MNA-SF scores in both men and women ( $p<0.001$ ). The evaluation of HGS and CC, which are the potential indicators of the nutritional status of older adults, is rapid and convenient.<sup>26,27</sup>

The strengths of our study are as follows. We included all residents (aged  $\geq 60$  years) of a senior home in Shanghai; thus, the findings appear to reflect real situations. Furthermore, several factors associated with malnutrition were identified in this study; these findings may

serve as a reference for future studies aimed at the prevention and treatment of malnutrition in older adults residing in nursing homes.

The present study has some limitations, such as the cross-sectional design and small sample size; thus, future prospective studies and clinical trials with larger sample sizes are warranted to investigate the causes of malnutrition in older adults.

In conclusion, we noted a high prevalence of malnutrition in older adults residing in a senior home in Shanghai. Several factors were found to be independently associated with malnutrition, such as dysphagia, possible sarcopenia, and dementia. Furthermore, exercise (at least thrice a week) was noted to be negatively associated with malnutrition. Systematic screening and personalized interventions should be developed to prevent and treat malnutrition in this population.

## **ACKNOWLEDGEMENTS**

The authors gratefully acknowledge support from Xiuli Yu, Guizhen Ren, Beibei Deng, Jiashuo Lu, Ziyi Han, Xingyu Duan, and Lingling Jiang.

## **CONFLICT OF INTEREST AND FUNDING DISCLOSURE**

The authors declare no personal or financial conflicts of interest.

This work was supported by the National Nutrition Science Research Grant (grant number: CNS-NNSRG2019-253) and the Shanghai Municipal Health System Important and Weak Discipline (Clinical Nutrition) Project (grant number: 2019ZB0102).

## **REFERENCES**

1. Norman K, Hass U, Pirlich M. Malnutrition in older adults-recent advances and remaining challenges. *Nutrients*. 2021;13:2464. doi: 10.3390/nu13082764.
2. Cox NJ, Morrison L, Ibrahim K, Robinson SM, Sayer AA, Roberts HC. New horizons in appetite and the anorexia of ageing. *Age Ageing*. 2020;49:526-34. doi: 10.1093/ageing/afaa014.
3. Chen LK, Arai H, Assantachai P, Akishita M, Chew STH, Dumlao LC, Duque G, Woo J. Roles of nutrition in muscle health of community-dwelling older adults: evidence-based expert consensus from Asian Working Group for Sarcopenia. *J Cachexia Sarcopenia Muscle*. 2022;13:1653-72. doi: 10.1002/jcsm.12981.
4. Tu WJ, Zeng X, Liu Q. Aging tsunami coming: the main finding from China's seventh national population census. *Aging Clin Exp Res*. 2022;34:1159-63. doi: 10.1007/s40520-021-02017-4.
5. Bell CL, Lee AS, Tamura BK. Malnutrition in the nursing home. *Curr Opin Clin Nutr Metab Care*. 2015;18:17-23. doi: 10.1097/MCO.0000000000000130.

6. Fayemendy P, Mabiama G, Vernier T, Massoulard-Gainant A, Villemonteix C, Desport JC, Jesus P. Nutritional status, dementia, and mobility among nursing home's residents: First exhaustive cross-sectional study in Limousin territory (France). *PLoS One*. 2021;16:e0250595. doi: 10.1371/journal.pone.0250595.
7. van Nie-Visser NC, Meijers J, Schols J, Lohrmann C, Bartholomeyczik S, Spreeuwenberg M, Halfens R. Which characteristics of nursing home residents influence differences in malnutrition prevalence? An international comparison of The Netherlands, Germany and Austria. *Br J Nutr*. 2014;111:1129-36. doi: 10.1017/S0007114513003541.
8. Tan X, Zhang Y, Shao H. Healthy China 2030, a breakthrough for improving health. *Glob Health Promot*. 2019;26:96-9. doi: 10.1177/1757975917743533.
9. Kaiser MJ, Bauer JM, Ramsch C, Uter W, Guigoz Y, Cederholm T, Thomas DR, Anthony P, Charlton KE, Maggio M, Tsai AC, Grathwohl D, Vellas B, Sieber CC; MNA-International Group. Validation of the Mini Nutritional Assessment short-form (MNA-SF): a practical tool for identification of nutritional status. *J Nutr Health Aging*. 2009;13:782-8. doi: 10.1007/s12603-009-0214-7.
10. Shanghai Municipal Bureau of Quality and Technical Supervision. Assessment for Elderly Care. 2013/2/8 [cited 2021/11/7]; Available from: <http://scjgj.sh.gov.cn/912/20210122/2c9bf2f675e566b40175e57eb4c8397d.html>.
11. Pazan F, Wehling M. Polypharmacy in older adults: a narrative review of definitions, epidemiology and consequences. *Eur Geriatr Med*. 2021;12:443-52. doi: 10.1007/s41999-021-00479-3.
12. Chen LK, Woo J, Assantachai P, Auyeung TW, Chou MY, Iijima K et al. Asian Working Group for Sarcopenia: 2019 Consensus update on sarcopenia diagnosis and treatment. *J Am Med Dir Assoc*. 2020;21:300-7.e2. doi: 10.1016/j.jamda.2019.12.012.
13. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*. 2019;48:16-31. doi: 10.1093/ageing/afy169.
14. Malmstrom TK, Morley JE. SARC-F: a simple questionnaire to rapidly diagnose sarcopenia. *J Am Med Dir Assoc*. 2013;14:531-2. doi: 10.1016/j.jamda.2013.05.018.
15. Barbosa-Silva TG, Menezes AM, Bielemann RM, Malmstrom TK, Gonzalez MC. Enhancing SARC-F: Improving sarcopenia screening in the clinical practice. *J Am Med Dir Assoc*. 2016;17:1136-41. doi: 10.1016/j.jamda.2016.08.004.
16. Bai HJ, Sun JQ, Chen M, Xu DF, Xie H, Yu ZW et al. Age-related decline in skeletal muscle mass and function among elderly men and women in Shanghai, China: a cross sectional study. *Asia Pac J Clin Nutr*. 2016;25:326-32. doi: 10.6133/apjcn.2016.25.2.14.
17. Liu H, Jiao J, Zhu C, Zhu M, Wen X, Jin J et al. Associations between nutritional status, sociodemographic characteristics, and health-related variables and health-related quality of life among Chinese elderly patients: A multicenter prospective study. *Front Nutr*. 2020;7:583161. doi: 10.3389/fnut.2020.583161.

18. Wu CF, Lu YW, Li H, Liu J, Shen JP, Ge LP. Dietary intake and nutritional status of 600 elderly in 3 long-term care institutions in Shanghai. *Geriatr Health Care (in Chinese)*. 2018;24:30-3.
19. Yang X, Chen YX, Zhang Y, Shi ZR, Wan WH, Zhang XH. Surveying and characteristics of comorbidities in senile people: report of 918 cases. *Chin J Heal Care Med (in Chinese)*. 2020;22:364-8. doi: 10.3969/j. issn.1674-3245.2020.04.006.
20. Wirth R, Dziewas R, Beck AM, Clavé P, Hamdy S, Heppner HJ et al. Oropharyngeal dysphagia in older persons - from pathophysiology to adequate intervention: a review and summary of an international expert meeting. *Clin Interv Aging*. 2016;11:189-208. doi: 10.2147/cia.S97481.
21. Suominen M, Muurinen S, Routasalo P, Soini H, Suur-Uski I, Peiponen A, Finne-Soveri H, Pitkala KH. Malnutrition and associated factors among aged residents in all nursing homes in Helsinki. *Eur J Clin Nutr*. 2005;59:578-83. doi: 10.1038/sj.ejcn.1602111.
22. Favaro-Moreira NC, Krausch-Hofmann S, Matthys C, Vereecken C, Vanhauwaert E, Declercq A, Bekkering GE, Duyck J. Risk factors for malnutrition in older adults: A systematic review of the literature based on longitudinal data. *Adv Nutr*. 2016;7:507-22. doi: 10.3945/an.115.011254.
23. Echeverria I, Amasene M, Urquiza M, Labayen I, Anaut P, Rodriguez-Larrad A, Irazusta J, Besga A. Multicomponent physical exercise in older adults after hospitalization: A randomized controlled trial comparing short- vs. long-term group-based interventions. *Int J Environ Res Public Health*. 2020;17:666. doi: 10.3390/ijerph17020666.
24. Argiles JM, Campos N, Lopez-Pedrosa JM, Rueda R, Rodriguez-Manas L. Skeletal muscle regulates metabolism via interorgan crosstalk: Roles in health and disease. *J Am Med Dir Assoc*. 2016;17:789-96. doi: 10.1016/j.jamda.2016.04.019.
25. Barazzoni R, Jensen GL, Correia MITD, Gonzalez MC, Higashiguchi T, Shi HP et al. Guidance for assessment of the muscle mass phenotypic criterion for the Global Leadership Initiative on Malnutrition (GLIM) diagnosis of malnutrition. *Clin Nutr*. 2022;41:1425-33. doi: 10.1016/j.clnu.2022.02.001.
26. Kizilarlanoglu MC, Kilic MK, Gokce D, Sakalar T, Ulger Z. Is it possible using handgrip strength instead of body mass index in MNA-SF test to assess the nutritional status of geriatric patients? *J Nutr Health Aging*. 2017;21:579-84. doi: 10.1007/s12603-016-0744-8.
27. Deutz NEP, Ashurst I, Ballesteros MD, Bear DE, Cruz-Jentoft AJ, Genton L, Landi F, Laviano A, Norman K, Prado CM. The underappreciated role of low muscle mass in the management of malnutrition. *J Am Med Dir Assoc*. 2019;20:22-7. doi: 10.1016/j.jamda.2018.11.021.

**Table 1.** Nutritional status and chronic diseases in participants

Characteristic	Overall (n=583)	Men (n=204)	Women (n=379)	<i>p</i>
MNA-SF, (mean±sd)	11.1±2.4	11.0±2.4	11.1±2.4	0.563
Malnutrition, n (%)	61 (10.5)	21 (10.3)	40 (10.6)	0.405
At risk of malnutrition, n (%)	218 (37.4)	84 (41.2)	134 (35.4)	
Normal nutritional status, n (%)	304 (52.1)	99 (48.5)	205 (54.1)	
Number of comorbidities, n (%)				0.163
<3	323 (55.4)	12 (59.3)	202 (53.3%)	
≥3	260 (44.6)	83 (40.7)	177 (46.7%)	
Polypharmacy, n (%)	281 (48.2)	103 (50.5)	178 (47.0)	0.417
Hypertension, n (%)	406 (69.6)	140 (68.6)	266 (70.2)	0.697
“Possible sarcopenia”, n (%)	384 (65.9)	134 (65.7)	250 (66.0)	0.946
Stroke, n (%)	271 (46.5)	106 (52.0)	165 (43.5)	0.520
Coronary disease, n (%)	235 (40.3)	71 (34.8)	164 (43.3)	0.047
Dementia, n (%)	184 (31.6)	54 (26.5)	130 (34.3)	0.520
Osteoporosis, n (%)	141 (24.2)	29 (14.2)	112 (29.6)	<0.001
Diabetes, n (%)	131 (22.5)	45 (22.1)	86 (22.7)	0.861
Dysphagia, n (%)	72 (12.3)	22 (10.8)	50 (13.2)	0.399
Hyperlipidemia, n (%)	71 (12.2)	23 (11.3)	48 (12.7)	0.642
Chronic bronchitis, n (%)	65 (11.1)	22 (10.8)	43 (11.3)	0.837
Lumbar disease, n (%)	57 (9.8)	15 (7.4)	42 (11.1)	0.148
History of tumors, n (%)	35 (6.0)	15 (7.4)	20 (5.3)	0.314
Renal insufficiency, n (%)	32 (5.5)	14 (6.9)	18 (4.7)	0.285
COPD, n (%)	21 (3.6)	9 (4.4)	12 (3.2)	0.441
Parkinson’s syndrome, n (%)	20 (3.4)	10 (4.9)	10 (2.6)	0.152

MNA-SF: Mini Nutritional Assessment-short form; COPD: chronic obstructive pulmonary disorder.

**Table 2.** Characteristics of the participants


Characteristic	Normal nutritional (n=304)	Malnutrition / malnutrition risk (n=279)	<i>p</i>
Age (year) (mean±sd)	85.7±6.3	86.2±6.8	0.332
Age (year) [n (%)]			0.369
60 ~ <80	41 (13.5)	45 (16.1)	
≥80	263 (86.5)	234 (83.9)	
Gender [n (%)]			0.200
Men	99 (32.6)	105 (37.6)	
Women	205 (67.4)	174 (62.4)	
High level of nursing care [n (%)]	233 (76.6)	260 (93.2)	<0.001
Exercise frequency (number of times per week; n (%))			
<3	65 (21.4)	182 (65.2)	<0.001
≥3	239 (78.6)	97 (34.8)	
General dietary [n (%)]	64 (21.1)	118 (42.3)	<0.001
Protein powder [n (%)]	22 (7.2)	40 (14.3)	0.005
Minerals and vitamin supplements [n (%)]	52 (17.1)	28 (10.0)	0.013
ONS, n (%)	35 (11.5)	32 (11.5)	0.987
Polypharmacy [n (%)]	130 (42.8)	151 (54.1)	0.006
Dysphagia [n (%)]	9 (3.0)	63 (22.6)	<0.001
History of tumors [n (%)]	19 (6.3)	16 (5.7)	0.794
Stroke [n (%)]	126 (41.4)	145 (52.0)	0.011
Dementia, n(%)	40 (13.2)	144 (51.6)	<0.001
Osteoporosis [n (%)]	64 (21.1)	77 (27.6)	0.065
Hyperlipidemia [n (%)]	41 (13.5)	30 (10.8)	0.313
Chronic bronchitis, n(%)	30 (9.9)	35 (12.5)	0.305
Lumbar disease, n (%)	27 (8.9)	30 (10.8)	0.447
Parkinson's syndrome, n (%)	9 (3.0)	11 (3.9)	0.515
History of fractures [n (%)]	7 (2.3)	31 (11.1)	<0.001
COPD, n (%)	7 (2.3)	14 (5.0)	0.079
Renal insufficiency, n (%)	14 (4.6)	18 (6.5)	0.328
Number of comorbidities ≥3, n (%)	80 (26.3)	180 (64.5)	<0.001
Hypertension, n (%)	222 (73.0)	184 (65.9)	0.063
Coronary disease, n (%)	115 (37.8)	120 (43.0)	0.203
Diabetes, n (%)	66 (21.7)	65 (23.3)	0.647
“Possible sarcopenia”, n (%)	145 (47.7)	239 (85.7)	<0.001

ONS: oral nutritional supplement; COPD: chronic obstructive pulmonary disorder.

**Table 3.** Results of multivariate logistic regression<sup>†</sup>

Variables	OR (95% CI)	<i>p</i>
Dysphagia		
No	1	
Yes	3.8 (1.7-8.5)	0.001
Exercise frequency (number of times per week)		
<3	1	
≥3	0.3 (0.2-0.5)	<0.001
Possible sarcopenia		
No	1	
Yes	3.6 (2.2-5.6)	<0.001
Dementia		
No	1	
Yes	4.5 (2.8-7.0)	<0.001

<sup>†</sup>Adjusted for age, nursing care level, exercise frequency, diet categories (general and nongeneral), polypharmacy, dysphagia, stroke, dementia, osteoporosis, history of fractures, chronic obstructive pulmonary disorder, comorbidities and possible sarcopenia



**Figure 1.** Flowchart for participant inclusion. MNA-SF: Mini Nutritional Assessment-short form; AWGS 2019: Asian Working Group for Sarcopenia 2019 Consensus.




Figure 2. Nutritional status of the participants (determined using the Mini Nutritional Assessment-short form questionnaire).




Figure 3. Correlations of nutrition status with handgrip strength and calf circumference.